

# My Math Notes

John Lockwood \*

June 2022

## 1 Introduction

Math Quantum Clep Calculus Youtube Study Guide

What I want to be able to do is to keep important math notes handy. The goal of all this is to take the Calculus CLEP Exam. One approach to this is the Modern States Course

## 2 Modern States Notes

### 2.1 Module 1, Limits

$$\frac{d}{dx}e^x = e^x$$

Continuity is essential to calculus

$$f(x) = \frac{3x^2 + 2x + 9}{4}$$

Epsilon/delta formal def of Limits

Limits are central to def of both derivative and integrals.

Limits happen at a certain point, not for whole function.

Technical (epsilon/delta) definition of limits given here

Limits might not exist

Function is continuous if  $f(x)$  exists  $\lim_{y \rightarrow x} f(y) = f(x)$

---

\*Special thanks to my lovely wife

He's not going to do left and right limits.

To get limit, evaluating function only works for continuous functions.

Polynomials are continuous, rational functions are more tricky. For example:

$$\lim_{x \rightarrow -1} \frac{x^2 + 2x + 1}{x + 1}$$

limit by subst. is  $0/0$ , which could be zero, one, infinity. Indeterminate form.

Can do it by factoring out  $x+1$ , and cancel one out.

If get zero over zero, need to factor.

Another example:  $\lim_{x \rightarrow 0} \frac{1}{x}$

Limit does not exist! DNE.

Watch out too for absolute value of  $x$  over  $x$  – no limit, because diff for pos and neg values.

Note not all piecewise functions are discontinuous.

General rules for continuous functions:

Polynomials, exponential functions, and sin and cos are continuous. Rational functions are continuous except at points where denominator is zero. Logarithm is continuous, because domain is only  $(0, \infty)$  Tangent function is discontinuous

Classic function that uses the squeeze theorem (memorize):  $\lim_{x \rightarrow 0} x \frac{\sin(x)}{x}$   
Answer ends up being one c.f. (Here's a proof.)

## 2.2 Theory of the Derivative

Derivative is slope of tangent line. A tangent line intersects function at only one point. Tangent lines – are constructed of the limit of secant lines (a line that intersects at exactly two points)

Definition of derivative:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

So derivative is defined in terms of a limit.

As we shrink average rate of change to zero, we get derivative, which is instantaneous rate of change.

## 3 Things to Research, Things To Memorize, Preliminaries

TODO: Add formulas for the area or volume of anything, area of circle, square (you know these), triangle, conic section, ellipse. Volume of sphere, cone, pyramid, cube (you know) etc.

### 3.1 Things to Research

Inverse functions

### 3.2 Things To Memorize

### 3.3 Limits of Trigonometric Functions

The proofs of the three identities here is given in Professor Leonard Calc 1, Video 1.2.

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \tag{1}$$

$$\lim_{x \rightarrow 0} \frac{1 - \cos x}{x} = 0 \tag{2}$$

$$\lim_{x \rightarrow 0} \frac{\tan x}{x} = 1 \tag{3}$$

### 3.4 Calculus Derivative Rules

#### 3.4.1 Derivatives of log functions

Derivative of Natural log of x:

$$\frac{d}{dx}[\ln x] = \frac{1}{x}$$

Derivative of log in another base is

$$\frac{d}{dx}[\log_a x] = \frac{1}{(\ln a)x}$$

For example:

$$\frac{d}{dx}[\log_{10} x] = \frac{1}{(\ln 10)x}$$

For video showing this, see Derivative of log at any base of x

### 3.4.2 Derivatives of inverse functions:

(Notes on Kahn academy video.)

Given:  $f(x)$  and  $g(x)$  are inverses, i.e.

$$g(x) = f^{-1}(x)$$

then the derivative of the inverse of  $f(x)$  is given as follows:

$$g'(x) = \frac{1}{f'(g(x))}$$

"This comes straight out of the chain rule." Proof:

$$f(g(x)) = x$$

(by inverse definition), therefore we can derivative of both sides:

$$\frac{d}{dx}f(g(x)) = \frac{d}{dx}x$$

by Chain rule we get:

$$f'(g(x)) * g'(x) = 1$$

therefore, just divide both sides by  $f'(x)$  to get

$$g'(x) = \frac{1}{f'(g(x))}$$

### 3.4.3 Derivatives of basic trig functions

$$\frac{d}{dx}\sin(x) = \cos(x)$$

$$\frac{d}{dx}\cos(x) = -\sin(x)$$

$$\frac{d}{dx}\tan(x) = \sec^2(x) = \frac{1}{\cos^2(x)}$$

### 3.4.4 Derivative of inverse sin of x

$$\frac{d}{dx}[\sin^{-1}(x)] = \frac{1}{\sqrt{1-x^2}}$$

This is the same as the derivative of arcsin of x, which is mentioned explicitly as "included" on the CLEP handout.

### 3.4.5 Derivative of inverse cos of x

$$\frac{d}{dx}[\cos^{-1}(x)] = -\frac{1}{\sqrt{1-x^2}}$$

This is same as derivative of arccosin


### 3.4.6 Derivative of inverse tan of x

$$\frac{d}{dx}[\tan^{-1}(x)] = \frac{1}{1+x^2}$$

This is the same as the derivative of arctan of x, which is mentioned explicitly as "included" on the CLEP handout.

### 3.4.7 Derivative of inverse tan of x

### 3.4.8 Unit Circle



### 3.4.9 Trigonometric Identities

$$\csc x = \frac{1}{\sin x}$$

$$\sec x = \frac{1}{\cos x}$$

$$\cot x = \frac{1}{\tan x}$$

$$\tan x = \frac{\sin x}{\cos x}$$

$$\cot x = \frac{\cos x}{\sin x}$$

$$\sin^2 x + \cos^2 x = 1$$

## 4 Misc

### 4.1 LaTeX examples

$$\left(\frac{f(x)}{g(x)}\right)$$