

# Monte Carlo Optimization

*Jonathan Navarrete*

*June 3, 2017*

## Introduction

This section will cover topics to optimization problems, and solutions Monte Carlo methods provide. Topics covered include

1. Stochastic Search and Simulated Annealing
2. EM Algorithm and MC EM

## Light bulb example

This exercise is taken from Flury and Zoppe, 2000, see Exercises in EM.

Below is the setup for the first exercise.

## The First Exercise

Suppose there are two light bulb survival experiments. In the first, there are  $N$  bulbs whose exact lifetimes  $y_i$  for  $i \in \{1, \dots, N\}$  are recorded. The lifetimes have an exponential distribution, such that  $y_i \sim \text{Exp}(\theta)$ . In the second experiment, there are  $M$  bulbs. After some time  $t > 0$ , a researcher walks into the room and only records how many lightbulbs are still burning out of  $M$  bulbs. Depending on whether the lightbulbs are still burning or out, the results from the second experiment are right- or -left-censored. There are indicators  $E_1, \dots, E_M$  for each of the bulbs in the second experiment. If the bulb is still burning,  $E_i = 1$ , else  $E_i = 0$ .

Given this information, our task is to solve for an MLE estimator for  $\theta$ .

Our first step in solving this is finding the joint likelihood for the observed and unobserved data (i.e. complete-data likelihood).

Let  $X_1, \dots, X_M$  be the (unobserved) lifetimes for the second experiment, and let  $Z = \sum_{i=1}^M E_i$  be the number of light bulbs still burning. Thus, the observed data from both the experiments combined is  $\mathcal{Y} = (Y_1, \dots, Y_N, E_1, \dots, E_M)$  and the unobserved data is  $\mathcal{X} = (X_1, \dots, X_M)$ .

The complete data log-likelihood is obtained by

$$\begin{aligned} L(\theta|X, Y) &= \prod_{i=1}^N \frac{1}{\theta} e^{y_i/\theta} \times \prod_{i=1}^M \frac{1}{\theta} e^{x_i/\theta} \\ &= \theta^{-N} e^{N\bar{y}/\theta} \times \theta^{-M} e^{\sum_{i=1}^M x_i/\theta} \end{aligned}$$

And log-likelihood is obtained by

$$\begin{aligned} \log(L(\theta)) &= N \times \log(\theta) + N\bar{y}/\theta + M \times \log(\theta) + \sum_{i=1}^M x_i/\theta \\ &= N(\log(\theta) + \bar{y}/\theta) + M \times \log(\theta) + \sum_{i=1}^M x_i/\theta \end{aligned}$$

(1)

$$\log^c(L(\theta|\mathcal{Y}, \mathcal{X})) = -N(\log(\theta) + \bar{Y}/\theta) - \sum_{i=1}^M (\log(\theta) + X_i/\theta)$$

which is linear for unobserved  $X_i$ . But

(2)

$$E[X_i|\mathcal{Y}] = E[X_i|E_i] = \begin{cases} t + \theta & \text{if } E_i = 1 \\ \theta - t \frac{e^{-t/\theta}}{1 - e^{-t/\theta}} & \text{if } E_i = 0 \end{cases}$$

and therefore the  $j$ th step consists of replacing  $X_i$  in (1) by its expected value (2), using the current numerical parameter value  $\theta^{(j-1)}$ . The result is

(3)

$$\log(L(\theta)) = -(N + M)\log(\theta) - \frac{1}{\theta}[N\bar{Y} + Z(t + \theta^{(j-1)}) + (M - Z)(\theta^{(j-1)} - tp^{(j-1)})]$$

where

$$p^{(j)} = \frac{e^{-t/\theta^{(j)}}}{1 - e^{-t/\theta^{(j)}}}$$

There is more (not shown here) in the paper, but my main concern is how  $p^{(j)}$  is obtained.