

# Monte Carlo Optimization

*Jonathan Navarrete*

*June 3, 2017*

## Introduction

This section will cover topics to optimization problems, and solutions Monte Carlo methods provide. Topics covered include

1. Stochastic Search and Simulated Annealing
2. EM Algorithm and MC EM

## Light bulb example

This exercise is taken from Flury and Zoppe, 2000, see Exercises in EM.

Below is the setup for the first exercise.

## The First Exercise

Suppose there are two light bulb survival experiments. In the first, there are  $N$  bulbs whose exact lifetimes  $y_i$  for  $i \in \{1, \dots, N\}$  are recorded. The lifetimes have an exponential distribution, such that  $y_i \sim \text{Exp}(\theta)$ . In the second experiment, there are  $M$  bulbs. After some time  $t > 0$ , a researcher walks into the room and only records how many lightbulbs are still burning out of  $M$  bulbs. Depending on whether the lightbulbs are still burning or out, the results from the second experiment are right- or -left-censored. There are indicators  $E_1, \dots, E_M$  for each of the bulbs in the second experiment. If the bulb is still burning,  $E_i = 1$ , else  $E_i = 0$ .

Given this information, our task is to solve for an MLE estimator for  $\theta$ .

Our first step in solving this is finding the joint likelihood for the observed and unobserved data (i.e. complete-data likelihood).

Let  $X_1, \dots, X_M$  be the (unobserved) lifetimes for the second experiment, and let  $Z = \sum_{i=1}^M E_i$  be the number of light bulbs still burning. Thus, the observed data from both the experiments combined is  $\mathcal{Y} = (Y_1, \dots, Y_N, E_1, \dots, E_M)$  and the unobserved data is  $\mathcal{X} = (X_1, \dots, X_M)$ .

The complete data log-likelihood is obtained by

$$\begin{aligned} L(\theta|X, Y) &= \prod_{i=1}^N \frac{1}{\theta} e^{y_i/\theta} \times \prod_{i=1}^M \frac{1}{\theta} e^{x_i/\theta} \\ &= \theta^{-N} e^{-N\bar{y}/\theta} \times \theta^{-M} e^{-\sum_{i=1}^M x_i/\theta} \end{aligned}$$

And log-likelihood is obtained by

$$\begin{aligned} \log(L(\theta)) &= -N \times \log(\theta) - N\bar{y}/\theta - M \times \log(\theta) + \sum_{i=1}^M x_i/\theta \\ &= -N(\log(\theta) + \bar{y}/\theta) - M \times \log(\theta) + \sum_{i=1}^M x_i/\theta \end{aligned}$$

Or as written by Flury and Zoppe,

$$\log^c(L(\theta|\mathcal{Y}, \mathcal{X})) = -N(\log(\theta) + \bar{Y}/\theta) - \sum_{i=1}^M (\log(\theta) + X_i/\theta)$$

The next step, is to take the expectation of  $\log(L(\theta))$  with respect to observed data.

$$\begin{aligned} E[\log(L(\theta))|\mathcal{Y}, \mathcal{X}] &= E[-N(\log(\theta) + \bar{Y}/\theta) - \sum_{i=1}^M (\log(\theta) + X_i/\theta)|\mathcal{Y}, \mathcal{X}] \\ &= -N(\log(\theta) + \bar{Y}/\theta) - E[\sum_{i=1}^M (\log(\theta) + X_i/\theta)|\mathcal{Y}, \mathcal{X}] \\ &= -N(\log(\theta) + \bar{Y}/\theta) - M \times \log(\theta) + E[\frac{1}{\theta} \sum_{i=1}^M X_i|\mathcal{Y}, \mathcal{X}] \\ &= -N(\log(\theta) + \bar{Y}/\theta) - M \times \log(\theta) + \frac{1}{\theta} \sum_{i=1}^M E[X_i|\mathcal{Y}, \mathcal{X}] \\ &= -N(\log(\theta) + \bar{Y}/\theta) - M \times \log(\theta) + \frac{1}{\theta} \sum_{i=1}^M E[X_i|E_i] \end{aligned}$$

which is linear for unobserved  $X_i$ . But

(2)

$$E[X_i|\mathcal{Y}] = E[X_i|E_i] = \begin{cases} t + \theta & \text{if } E_i = 1 \\ \theta - t \frac{e^{-t/\theta}}{1 - e^{-t/\theta}} & \text{if } E_i = 0 \end{cases}$$

For the first case,  $E_i = 1$ , so

$$\begin{aligned} E[x_i|x_i > t] &= E[x_i + t] \\ &= t + E[x_i] \\ &= t + \theta \end{aligned}$$

For the second case,  $E_i = 0$ , then

$$\int_0^t P(X_i > x|X_i < t) dx = \int_0^t \frac{P(x < X_i < t)}{P(X_i < t)} dx$$

For the denominator, we get

$$\begin{aligned} P(X_i < t) &= \int_0^t \frac{1}{\theta} e^{-x_i/\theta} dx \\ &= \frac{1}{\theta} (-\theta e^{-x_i/\theta}) \Big|_0^t \\ &= 1 - e^{-t/\theta} \end{aligned}$$

and for the numerator we obtain

$$\begin{aligned} P(x < X_i < t) &= \int_x^t \frac{1}{\theta} e^{-x_i/\theta} dx \\ &= \frac{1}{\theta} (-\theta e^{-x_i/\theta}) \Big|_x^t \\ &= e^{-x/\theta} - e^{-t/\theta} \end{aligned}$$

Altogether, we obtain

$$\begin{aligned}
\int_0^t P(X_i > x | X_i < t) dx &= \int_0^t \frac{P(x < X_i < t)}{P(X_i < t)} dx \\
&= \int_0^t \frac{e^{-x/\theta} - e^{-t/\theta}}{(1 - e^{-t/\theta})} dx \\
&= \frac{1}{(1 - e^{-t/\theta})} \int_0^t (e^{-x/\theta} - e^{-t/\theta}) dx \\
&= \frac{1}{(1 - e^{-t/\theta})} \left( \int_0^t e^{-x/\theta} dx - \int_0^t e^{-t/\theta} dx \right) \\
&= \frac{1}{(1 - e^{-t/\theta})} (\theta(1 - e^{-t/\theta}) - x \times e^{-t/\theta} \Big|_0^t) \\
&= \theta - t \times \frac{e^{-t/\theta}}{1 - e^{-t/\theta}}
\end{aligned}$$

In order to calculate EM estimates for  $\theta$ , we will plug in the expected values

$$E[X_i | \mathcal{Y}] = E[X_i | E_i] = \begin{cases} t + \theta & \text{if } E_i = 1 \\ \theta - t \frac{e^{-t/\theta}}{1 - e^{-t/\theta}} & \text{if } E_i = 0 \end{cases}$$

into the log-likelihood

$$\begin{aligned}
\log(L(\theta)) &= -N(\log(\theta) + \bar{y}/\theta) - M \times \log(\theta) + \sum_{i=1}^M x_i/\theta \\
&= -N \times \log(\theta) - N\bar{y}/\theta - M \times \log(\theta) + \sum_{i=1}^M x_i/\theta \\
&= -(N + M) \times \log(\theta) - N\bar{y}/\theta + \sum_{i=1}^M x_i/\theta \\
&= -(N + M) \times \log(\theta) - \frac{1}{\theta} (N\bar{y} + \sum_{i=1}^M x_i) \\
&= -(N + M) \log(\theta) - \frac{1}{\theta} [N\bar{Y} + Z(t + \theta) + (M - Z)(\theta - t \times \frac{e^{-t/\theta}}{1 - e^{-t/\theta}})]
\end{aligned}$$

As we iterate through estimates of  $\theta$ , we will use conditioned estimates of  $\theta$  given previous estimates of  $\theta$ . Such that the  $j$ th step consists of replacing  $X_i$  in (1) by its expected value (2), using the current numerical parameter value  $\theta^{(j-1)}$ .

(3)

$$\log(L(\theta)) = -(N + M) \log(\theta) - \frac{1}{\theta} [N\bar{Y} + Z(t + \theta^{(j-1)}) + (M - Z)(\theta^{(j-1)} - t p^{(j-1)})]$$

where

$$p^{(j)} = \frac{e^{-t/\theta^{(j)}}}{1 - e^{-t/\theta^{(j)}}}$$

Once we take the derivative of the log-likelihood and set it to zero, we will come up with an estimate for  $\theta$

$$\begin{aligned}\frac{d}{dx} \ln(L(\theta)) &= 0 \\ 0 &= -\frac{(N+M)}{\theta} + \frac{1}{\theta^2} [N\bar{Y} + Z(t+\theta) + (M-Z)(\theta - t \times \frac{e^{-t/\theta}}{1-e^{-t/\theta}})] \\ \frac{(N+M)}{\theta} &= \frac{1}{\theta^2} [N\bar{Y} + Z(t+\theta) + (M-Z)(\theta - t \times \frac{e^{-t/\theta}}{1-e^{-t/\theta}})] \\ \theta &= [N\bar{Y} + Z(t+\theta) + (M-Z)(\theta - t \times \frac{e^{-t/\theta}}{1-e^{-t/\theta}})] / (N+M)\end{aligned}$$

Thus, for each  $j$ th M-step, we will calculate

$$\begin{aligned}\theta^{(j)} &= f(\theta^{(j-1)}) \\ \theta &= [N\bar{Y} + Z(t+\theta^{(j-1)}) + (M-Z)(\theta^{(j-1)} - t \times \frac{e^{-t/\theta^{(j-1)}}}{1-e^{-t/\theta^{(j-1)}}})] / (N+M)\end{aligned}$$

```
set.seed(5678)
theta = 5
rate = 1/theta

t = 5
N = 100
M = 50
y = rexp(n = N, rate = rate)
x = rexp(n = M, rate = rate)
x = sort(x)
E = as.integer(x > t)

N.ybar = sum(y)
Z = sum(E)
t = 5

theta.j = 0.1
theta.jp1 = 0.5
for(i in 1:10){
  theta.j = theta.jp1
  p = (exp(-t/theta.j)/(1-exp(-t/theta.j)))
  theta.jp1 = (N.ybar + Z*( t + theta.j) + (M-Z)*(theta.j - t*p) ) / (N+M)
  print(theta.jp1)
}

## [1] 4.624345
## [1] 5.366158
## [1] 5.445061
## [1] 5.45323
## [1] 5.454073
## [1] 5.45416
## [1] 5.454169
## [1] 5.45417
## [1] 5.45417
## [1] 5.45417

## compare against MLE from observed data
mean(y)
```

```
## [1] 6.036602
```

```
## note, results will vary if you remove seed
```