



# High Dimensional Space


[longhuan@sjtu.edu.cn](mailto:longhuan@sjtu.edu.cn)



# Word Vector Model



## Web Page Model



- Nearest neighbor query
- Information retrieval
- Web page rank
- Online recommendation
- .....

The law of Large numbers

Properties of High-Dimensional space,  
unit ball

Generating points uniformly at random  
from a ball

Gaussians in High Dimension

Random Projection and Johnson-  
Lindenstrauss Lemma

Seperating Gaussians

# Normal distribution (Gauss Distribution)

$X \sim N(\mu, \sigma^2)$ , with density function:

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad -\infty < x < +\infty$$

**Variance**

$$\begin{aligned} \text{Var}(X) &= E((X - E[X])^2) \\ &= E(X^2 + E[X]^2 - 2XE[X]) \\ &= E(X^2 - E[X]^2) \\ &= E[X^2] - E[X]^2 \end{aligned}$$

## Chebyshev's Inequality

$$\forall a > 0, \Pr(|X - E(X)| \geq a) \leq \frac{\text{Var}[X]}{a^2}$$

# Law of Large Numbers

- In probability theory, the **law of large numbers (LLN)** is a theorem that describes the result of performing the same experiment a large number of times.
- According to the law, the average of the results obtained from a large number of trials should be close to the expected value, and will tend to become closer as more trials are performed.

# Law of large numbers

Let  $x_1, x_2, \dots, x_n$  be  $n$  independent samples of a random variable  $x$ , then

$$\Pr\left(\left|\frac{x_1 + x_2 + \dots + x_n}{n} - E(x)\right| \geq \epsilon\right) \leq \frac{\text{Var}(x)}{n\epsilon^2}$$

**Proof.** (Chebychev's Inequality)

$$\begin{aligned}\Pr\left(\left|\frac{x_1 + x_2 + \dots + x_n}{n} - E(x)\right| \geq \epsilon\right) &\leq \frac{\text{Var}\left(\frac{x_1 + x_2 + \dots + x_n}{n}\right)}{\epsilon^2} \\ &= \frac{\text{Var}(x_1 + x_2 + \dots + x_n)}{n^2 \epsilon^2} \\ &= \frac{\text{Var}(x)}{n \epsilon^2}\end{aligned}$$

# Application

- $\mathbf{x}$  be a  $d$  –dimensional random point whose coordinates are each selected from  $N\left(0, \frac{1}{2\pi}\right)$ ,
- i.e.  $\mathbf{x} = [x_1, x_2, \dots, x_d]$  with  $x_i \sim N\left(0, \frac{1}{2\pi}\right)$
- **By LLN:**  $|\mathbf{x}|^2 = \sum_{i=1}^d x_i^2 = \frac{d}{2\pi} = \Theta(d)$  with high probability.
- **The probability that point  $\mathbf{x}$  lie in the unit ball is *vanishingly small*.**

# Application

- $\mathbf{x}, \mathbf{y} : [z_1, z_2, \dots, z_d]$  with  $z_i \sim N(0, 1)$
- $|\mathbf{x}|^2 \approx d, |\mathbf{y}|^2 \approx d,$
- $|\mathbf{x} - \mathbf{y}|^2 \approx ?$



# Application

- $\mathbf{x}, \mathbf{y} : [z_1, z_2, \dots, z_d]$  with  $z_i \sim N(0, 1)$
- $|\mathbf{x}|^2 \approx d, |\mathbf{y}|^2 \approx d,$
- $|\mathbf{x} - \mathbf{y}|^2 = \sum_{i=1}^d (x_i - y_i)^2$   
$$E(x_i - y_i)^2 = E(x_i^2) + E(y_i^2) - 2E(x_i y_i)$$
$$= 1 + 1 - 2E(x_i)E(y_i) = 2.$$


# Application

- $\mathbf{x}, \mathbf{y} : [z_1, z_2, \dots, z_d]$  with  $z_i \sim N(0, 1)$
- $|\mathbf{x}|^2 \approx d, |\mathbf{y}|^2 \approx d,$
- $|\mathbf{x} - \mathbf{y}|^2 = \sum_{i=1}^d (x_i - y_i)^2 = 2d$   
$$E(x_i - y_i)^2 = E(x_i^2) + E(y_i^2) - 2E(x_i y_i)$$
$$= 1 + 1 - 2E(x_i)E(y_i) = 2.$$
- $|\mathbf{x} - \mathbf{y}|^2 \approx |\mathbf{x}|^2 + |\mathbf{y}|^2$
- **Pythagorean theorem**  $\Rightarrow$  random  $d$  –dimensional  $\mathbf{x}, \mathbf{y}$  are approximately orthogonal.

# Application

- $\mathbf{x}, \mathbf{y} : [z_1, z_2, \dots, z_d]$  with  $z_i \sim N(0, 1)$
- **Pythagorean theorem**  $\Rightarrow$  random  $d$  –dimensional  $\mathbf{x}, \mathbf{y}$  are **approximately orthogonal**.

- If we scale these random points to be unit length and call  $\mathbf{x}$  the **North Pole**, *much of the surface area of the unit ball must lie near the equator.*



# Table of tail bounds

	Condition	Tail bound
Markov	$x \geq 0$	$\text{Prob}(x \geq a) \leq \frac{E(x)}{a}$
Chebychev	Any $x$	$\text{Prob}( x - E(x)  \geq a) \leq \frac{\text{Var}(x)}{a^2}$
Chernoff	$x = x_1 + x_2 + \dots + x_n$ $x_i \in [0, 1]$ i.i.d. Bernoulli;	$\text{Prob}( x - E(x)  \geq \varepsilon E(x)) \leq 3e^{-c\varepsilon^2 E(x)}$
Higher Moments	$r$ positive even integer	$\text{Prob}( x  \geq a) \leq E(x^r)/a^r$
Gaussian Annulus	$x = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}$ $x_i \sim N(0, 1); \beta \leq \sqrt{n}$ indep.	$\text{Prob}( x - \sqrt{n}  \geq \beta) \leq 3e^{-c\beta^2}$
Power Law for $x_i$ ; order $k \geq 4$	$x = x_1 + x_2 + \dots + x_n$ $x_i$ i.i.d ; $\varepsilon \leq 1/k^2$	$\text{Prob}( x - E(x)  \geq \varepsilon E(x)) \leq (4/\varepsilon^2 kn)^{(k-3)/2}$

The law of Large numbers

Properties of High-Dimensional space,  
unit ball


Generating points uniformly at random  
from a ball

Gaussians in High Dimension

Random Projection and Johnson-  
Lindenstrauss Lemma


Seperating Gaussians

# Relationship between the sphere and cube



The difference between the volume of a **cube** with unit-length sides and the volume of a unit-radius **sphere** at the dimensions: 2, 4 and  $d$ .

# Conceptual drawing of a sphere and a cube



For large  $d$ , almost all the volume of the cube is located outside the sphere.

# Geometry of High Dimensions

- Most of the volume of the high-dimensional objects is near the surface:


$$\frac{\text{Volume}((1 - \epsilon)A)}{\text{Volume}(A)} = (1 - \epsilon)^d \leq e^{-\epsilon d}$$

Fix  $\epsilon$  and letting  $d \rightarrow \infty$ , the above quantity rapidly approaches zero.

- Application:**

$S$  be the unit ball in  $d$  –dimensions (i.e., the set of points within distance 1 of the origin). Then  $1 - e^{-\epsilon d}$  fraction of the volume is in  $S \setminus (1 - \epsilon)S$ .

Especially, consider  $\epsilon = \frac{1}{d}$ .





# Unit ball in $d$ – dimensions


- **Surface:**  $A(d) = \frac{2\pi^{\frac{d}{2}}}{\Gamma(\frac{d}{2})}$ , **Volume:**  $V(d) = \frac{2}{d} \frac{\pi^{\frac{d}{2}}}{\Gamma(\frac{d}{2})}$ .

- $V(2) = \pi, V(3) = \frac{4}{3}\pi$ ,  $\lim_{n \rightarrow \infty} V(d) = 0$ .

随着维度增加，单位球的体积会无限趋近于0

- **Most of the volume of a unit ball in high dimensions is concentrated near its equator no matter which direction is defined to be the North Pole.**

**Theorem:** For  $c \geq 1$  and  $d \geq 3$ , at least a  $1 - \frac{2}{c} e^{-c^2/2}$  fraction of the volume of the  $d$  – dimensional unit ball has  $|x_1| \leq \frac{c}{\sqrt{d-1}}$ .



The law of Large numbers

Properties of High-Dimensional space,  
unit ball

Generating points uniformly at random  
from a ball


Gaussians in High Dimension

Random Projection and Johnson-  
Lindenstrauss Lemma

Seperating Gaussians


# Generating points uniformly at random on the surface of the unit ball

- $d = 2$




# Generating points uniformly at random on the surface of the unit ball

- $d = 2$ 
  - Generate  $x_i, y_i$  u.a.r from the interval  $[-1,1]$ ;




# Generating points uniformly at random on the surface of the unit ball

- $d = 2$ 
  - Generate  $x_i, y_i$  u.a.r from the interval  $[-1,1]$ ;




# Generating points uniformly at random on the surface of the unit ball

- $d = 2$ 
  - Generate  $x_i, y_i$  u.a.r from the interval  $[-1,1]$ ;
  - Discard the points outside the unit circle;



# Generating points uniformly at random on the surface of the unit ball

- $d = 2$ 
  - Generate  $x_i, y_i$  u.a.r from the interval  $[-1,1]$ ;
  - Discard the points outside the unit circle;
  - Project the remaining points onto the circle.
- How about  $d$  is large?
  - The above strategy would fail. (why?)
  - ① **Surface:** Spherical normal distribution + Normalizing.
  - ② **Surface+interior:** Scale the point on the surface.



The law of Large numbers

Properties of High-Dimensional space,  
unit ball

Generating points uniformly at random  
from a ball


**Gaussians in High Dimension**

Random Projection and Johnson-  
Lindenstrauss Lemma

Seperating Gaussians



- 1-dimensional Gaussian



- $d$  –dimensional spherical Gaussian with 0 means and variance  $\sigma^2$  in each coordinate has density function:

$$p(x) = \frac{1}{(2\pi)^{d/2} \sigma^d} \exp\left(-\frac{|x|^2}{2\sigma^2}\right)$$

- Integrate the PDF over a **unit ball** centered at the origin will cover **almost 0 mass**, for the volume of such a ball is negligible.
- The radius of the ball need to be nearly  $\sqrt{d}$  before there is a **significant volume** and hence significant probability mass.

# Gaussian Annulus Theorem

- For a  $d$  –dimensional spherical Gaussian with unit variance in each direction ,for any  $\beta \leq \sqrt{d}$ , all but at most  $3e^{-c\beta^2}$  of the probability mass lies within the annulus

$$\sqrt{d} - \beta \leq |x| \leq \sqrt{d} + \beta$$

where  $c$  is a fixed positive constant.

The law of Large numbers

Properties of High-Dimensional space,  
unit ball

Generating points uniformly at random  
from a ball

Gaussians in High Dimension

Random Projection and Johnson-  
Lindenstrauss Lemma

Separating Gaussians

# Database query: Nearest neighbor search

$n$  points from  $R^d$ : 
$$\begin{bmatrix} v_{11} & v_{21} & \vdots & v_{n1} \\ v_{12} & v_{22} & \vdots & v_{n2} \\ \vdots & \vdots & \vdots & \vdots \\ v_{1d} & v_{2d} & \vdots & v_{nd} \end{bmatrix}$$

- **Nearest neighbor search:** find the nearest or approximately nearest database point to the query point.
- When  $d$  is large, it could cost more than expected.
- **Dimension reduction :** *Project* the database points to a  $k$  dimensional space with  $k \ll d$ . It will work so long as the relative distances between points are approximately preserved.

# Projection function

- Pick  $k$  vectors  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ , independently from the Gaussian distribution

$\frac{1}{(2\pi)^{d/2} \sigma^d} \exp\left(-\frac{|\mathbf{x}|^2}{2\sigma^2}\right)$ , for any vector  $\mathbf{v}$ , the projection  $f: \mathbb{R}^d \rightarrow \mathbb{R}^k$  is:

$$f(\mathbf{v}) = (\mathbf{u}_1 \cdot \mathbf{v}, \mathbf{u}_2 \cdot \mathbf{v}, \dots, \mathbf{u}_k \cdot \mathbf{v})$$

# Projection function

Pick  $k$  vectors

$\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ , independent  
ly from the Gaussian  
distribution

$\frac{1}{(2\pi)^{d/2} \sigma^d} \exp\left(-\frac{|\mathbf{x}|^2}{2\sigma^2}\right)$ , for  
any vector  $\mathbf{v}$ , the  
projection  $f: R^d \rightarrow R^k$  is:

$$f(\mathbf{v}) = (\mathbf{u}_1 \cdot \mathbf{v}, \mathbf{u}_2 \cdot \mathbf{v}, \dots, \mathbf{u}_k \cdot \mathbf{v})$$

- $f(\mathbf{v}_1 - \mathbf{v}_2) = f(\mathbf{v}_1) - f(\mathbf{v}_2)$
- $|f(\mathbf{v})| \approx \sqrt{k}|\mathbf{v}|$  w.h.p.
- To estimate  $|\mathbf{v}_1 - \mathbf{v}_2|$ , it suffices to compute  $|f(\mathbf{v}_1) - f(\mathbf{v}_2)|$

# Random Projection Theorem

- Let  $v$  be a fixed vector in  $R^d$  and let  $f$  be defined as above. Then there exists constant  $c > 0$  such that for  $\epsilon \in (0,1)$

$$\Pr \left( \left| \|f(v) - \sqrt{k}|v|\right| \geq \epsilon \sqrt{k}|v| \right) \leq 3e^{-ck\epsilon^2}$$



# Johnson-Lindenstrass Lemma

- For any  $0 < \epsilon < 1$  and any integer  $n$ , let  $k \geq \frac{3}{c\epsilon^2} \ln n$  for  $c$  as in the Gaussian Annulus theorem, for any set of  $n$  points in  $R^d$ , the random projection  $f$  defined above has the property that **for all pairs of points**  $v_i$  and  $v_j$ , with probability at least  $1 - \frac{1.5}{n}$   
 $(1 - \epsilon)\sqrt{k}|v_i - v_j| \leq |f(v_i) - f(v_j)| \leq (1 + \epsilon)\sqrt{k}|v_i - v_j|.$

# Comments

- JL lemma works for all pairs of points,
- $k$  depends on  $\ln n$ ,
- To the database, JL Lemma says the algorithm will yield the right answer with high probability whatever the query is.

The law of Large numbers


Properties of High-Dimensional space,  
unit ball

Generating points uniformly at random  
from a ball


Gaussians in High Dimension

Random Projection and Johnson-  
Lindenstrauss Lemma


Separating Gaussians



- Mixtures of Gaussians
- Parameter estimation problem



- When  $\Delta \in \Omega(d^{1/4})$



- Algorithm for separating points from two Gaussians:** Calculate all pairwise distance between points. The cluster of smallest pairwise distances must come from a single Gaussian. Remove these points. The remaining points come from the second Gaussian.