



# Combinatorial Counting

[longhuan@sjtu.edu.cn](mailto:longhuan@sjtu.edu.cn)





# Let's Count!



# $n$ balls are put into $m$ bins

balls per bin	unrestricted	$\leq 1$	$\geq 1$
$n$ distinct balls, $m$ distinct bins.			
$n$ identical balls, $m$ distinct bins.			
$n$ distinct balls, $m$ identical bins.			
$n$ identical balls, $m$ identical bins.			

# $n$ balls are put into $m$ bins

balls per bin	unrestricted	$\leq 1$	$\geq 1$
$n$ distinct balls, $m$ distinct bins.			
$n$ identical balls, $m$ distinct bins.			
$n$ distinct balls, $m$ identical bins.		$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	
$n$ identical balls, $m$ identical bins.		$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	


Basic counting

Binomial theorem

Generalized Binomial theorem Some

special numbers

We will start with  
counting the *ordered*  
objects.




Ordered sequence




- **Problem1:** How many 5-letter words are there(using the 26-letter English alphabet)?  
e. g. abcde, sssdd, ...
- **Problem2:** How many **distinct** 5-letter words are there(using the 26-letter English alphabet) ?  
e. g. abcde, ~~sssdd~~, ...

# 5-letter words




$$26 \times 26 \times 26 \times 26 \times 26 = 26^5$$


# Distinct 5-letter words



# Distinct 5-letter words



# Distinct 5-letter words



$$26 \times 25 \times 24 \times 23 \times 22$$

# Proof by induction


**Goal:** show that  $P(x)$  is true for any  $x \in \omega$

- ① Check that  $P(0)$  is true;
- ② Suppose that  $P(k)$  is true; // Induction hypothesis
- ③ Prove that  $P(k + 1)$  is true.

# The generalization of Problem 1

- **Proposition 1:** Let  $N$  be an  $n$ -element set, and  $M$  be an  $m$ -element set, with  $n \geq 0, m \geq 1$ . Then the number of all possible mappings  $f: N \rightarrow M$  is  $m^n$ .
- Proof: ( By induction on  $n$ )
  - $n = 0$ :  $f = \emptyset$ ;  $m^0 = 1$  .
  - Suppose the results works for  $n = k$ ;
  - If  $n = k + 1$  :

$n = k + 1$  , take any  $a \in N$ :



$$m \cdot m^{n-1} = m^n$$



# $n$ balls are put into $m$ bins

balls per bin	unrestricted	$\leq 1$	$\geq 1$
$n$ distinct balls, $m$ distinct bins.	$m^n$		
$n$ identical balls, $m$ distinct bins.			
$n$ distinct balls, $m$ identical bins.		$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	
$n$ identical balls, $m$ identical bins.		$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	

# The generalization of Problem 2


- **Proposition 2:** Let  $N$  be an  $n$ -element set, and  $M$  be an  $m$ -element set, with  $n, m \geq 0$ . Then there exist exactly

$$m(m-1) \dots (m-n+1) = \prod_{i=0}^{n-1} (m-i)$$

**one-to-one** mappings from  $N$  into  $M$ .

- **Proof:** ( By induction on  $n$ )
  - $n = 0$ :  $f = \emptyset$ . The value of an empty product is defined as 1.
  - Suppose the results works for  $n = k$ ;

– for  $n = k + 1$ , take any  $a \in N$ :



$$m(m-1) \dots (m-n+1)$$

# Falling factorial notation

$$\begin{aligned} & (x)_n \\ &= x^{\underline{n}} \\ &= x(x-1)\cdots(x-n+1) \end{aligned}$$

# $n$ balls are put into $m$ bins

balls per bin	unrestricted	$\leq 1$	$\geq 1$
$n$ distinct balls, $m$ distinct bins.	$m^n$	$(m)_n$	
$n$ identical balls, $m$ distinct bins.			
$n$ distinct balls, $m$ identical bins.		$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	
$n$ identical balls, $m$ identical bins.		$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	

# Application 1: Counting the different subsets

Given set  $X$ ,  $|X| = n$ , then  $X$  has exactly  $2^n$  subsets ( $n \geq 0$ ).


- Proof<sup>1</sup>: By induction on  $n$ . (Exercise)
- Proof<sup>2</sup>:

for any  $A \subseteq X$ , define  $f_A: X \rightarrow \{0,1\}$  as

$$f_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$

# Characteristic function

$$f_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$



There exists a **bijective** relation between the subsets of  $X$  and  $f: X \rightarrow \{0,1\}$  (Recall: Equinumerous).

# Application2: Counting the permutations

- ***Permutation***: A bijective mapping of a finite set  $X$  to itself is called a permutation of the set  $X$ .
- Recall: Bijective functions.




# Counting permutations-**Factorial**

Given set  $X$  ,  $|X| = n$ , then there are  $n \cdot (n - 1) \cdot \dots \cdot 2 \cdot 1$  different permutations on set  $X$ .


***n factorial:***

$$n! = n \cdot (n - 1) \cdot \dots \cdot 2 \cdot 1 = \prod_{i=1}^n i.$$

- So far, we considered **ordered** sequences.
- What about the **un-ordered** occasion?



Ordered sequence



Un-ordered set

### Problem 3: counting $k$ -element subsets

Given set  $X$ ,  $|X| = n$ ,  $n \geq k \geq 0$ , how many different subsets of  $X$  contains exactly  $k$  elements?

e. g.  $X = \{a, b, c\}$ ,  $k = 2$ .

Then:  $\{a, b\}$ ,  $\{a, c\}$ ,  $\{b, c\}$ . Three 2-size subsets.

**Convention:**  $\binom{X}{k}$  VS.  $|\binom{X}{k}|$

e. g.  $\binom{X}{k} = \{\{a, b\}, \{a, c\}, \{b, c\}\}$ ,  $|\binom{X}{k}| = 3$ .

- **Proposition:** For any finite set  $X$  with  $|X| = n$ , the number of all  $k$ -element subsets is

$$\left| \binom{X}{k} \right| = \frac{n(n-1)(n-2)\dots(n-k+1)}{k(k-1)\dots 2 \cdot 1}.$$


- Proof: (Double counting!)

# Binomial coefficients

- $\boxed{\binom{n}{k}} = \left| \binom{X}{k} \right| = \frac{n(n-1)(n-2)\dots(n-k+1)}{k(k-1)\dots 2 \cdot 1}$   
 $= \frac{\prod_{i=0}^{k-1} (n-i)}{k!}$   
 $= \frac{n(n-1)(n-2)\dots(n-k+1) \cdot (n-k) \cdot \dots \cdot 1}{k(k-1)\dots 2 \cdot 1 \cdot (n-k) \cdot \dots \cdot 1}$   
 $= \boxed{\frac{n!}{k! \cdot (n-k)!}}$

Application: counting non-negative solutions.

$m \geq r \geq 0$ , the equation  $x_1 + x_2 + \cdots + x_r = m$  has  $\binom{m+r-1}{r-1}$  non-negative integer solutions of the form  $(x_1, x_2, \dots, x_r)$ .




$$x_1 = 3, x_2 = 0, x_3 = 2, x_4 = 0, \dots, x_{r-1} = 4, x_r = 2$$




Application: counting **non-negative** solutions.

$m \geq r \geq 0$ , the equation  $x_1 + x_2 + \cdots + x_r = m$  has  $\binom{m+r-1}{r-1}$  non-negative integer solutions of the form  $(x_1, x_2, \dots, x_r)$ .




Application: counting **non-negative** solutions.

$m \geq r \geq 0$ , the equation  $x_1 + x_2 + \cdots + x_r = m$  has  $\binom{m+r-1}{r-1}$  non-negative integer solutions of the form  $(x_1, x_2, \dots, x_r)$ .




Question: counting **positive** solutions.

$m \geq r \geq 0$ , the equation  $x_1 + x_2 + \cdots + x_r = m$  has \_\_\_\_\_ **positive** integers solutions of the form  $(x_1, x_2, \dots, x_r)$  .



**Question:** counting **positive** solutions.

$m \geq r \geq 0$ , the equation  $x_1 + x_2 + \cdots + x_r = m$  has  $\binom{m-1}{r-1}$  **positive** integers solutions of the form  $(x_1, x_2, \dots, x_r)$  .




# Basic Properties

$$\binom{n}{k} = \binom{n}{n-k}$$

- Proof<sup>1</sup>:

- Proof<sup>2</sup>:




# Pascal's Identity:


$$\binom{n-1}{k-1} + \binom{n-1}{k} = \binom{n}{k}$$

- Proof:



$$\binom{n-1}{k-1}$$



$$\binom{n-1}{k}$$



# Pascal's Triangle (1654) / 杨辉三角 (1261)



# Exercise


$$\sum_{k=0}^n \binom{k}{m} = \binom{n+1}{m+1}$$

$$\sum_{k=0}^n \binom{m+k-1}{k} = \binom{n+m}{n}$$



$$\sum_{i=0}^n \binom{n}{i}^2 = \binom{2n}{n}$$

- Proof:  $\sum_{i=0}^n \binom{n}{i}^2 = \sum_{i=0}^n \binom{n}{i} \binom{n}{n-i}$



# Vandermonde's identity/convolution

$$\binom{m+n}{r} = \sum_{k=0}^r \binom{m}{k} \binom{n}{r-k}$$

The general form

$$\binom{n_1 + \dots + n_p}{m} = \sum_{k_1 + \dots + k_p = m} \binom{n_1}{k_1} \binom{n_2}{k_2} \dots \binom{n_p}{k_p}$$

# $n$ balls are put into $m$ bins

balls per bin	unrestricted	$\leq 1$	$\geq 1$
$n$ distinct balls, $m$ distinct bins.	$m^n$	$(m)_n$	
$n$ identical balls, $m$ distinct bins.	$\binom{n+m-1}{m-1}$	$\binom{m}{n}$	$\binom{n-1}{m-1}$
$n$ distinct balls, $m$ identical bins.		$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	
$n$ identical balls, $m$ identical bins.		$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	

# Multiset Coefficient

- The number of multisets of cardinality  $k$ , with elements taken from a finite set of cardinality  $n$ , is called the **multiset coefficient** or **multiset number**.

- $$\left( \binom{n}{k} \right) = \binom{n+k-1}{n-1} = \binom{n+k-1}{k}$$

转化为 $x_1+x_2+\dots+x_n=k$ 的非负整数解

$$= \frac{n(n+1)(n+2)\cdots(n+k-1)}{k!} = \frac{n^{\overline{k}}}{k!}$$

# $n$ balls are put into $m$ bins

balls per bin	unrestricted	$\leq 1$	$\geq 1$
$n$ distinct balls, $m$ distinct bins.	$m^n$	$(m)_n$	
$n$ identical balls, $m$ distinct bins.	$\binom{m+n-1}{n}$	$\binom{m}{n}$	$\binom{n-1}{m-1}$
$n$ distinct balls, $m$ identical bins.		$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	
$n$ identical balls, $m$ identical bins.		$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	

Basic counting

Binomial theorem

Generalized Binomial theorem

Some special numbers

# Binomial theorem

- **Binomial Theorem:** for any non-negative integer  $n$ , we have

$$(1 + x)^n = \sum_{k=0}^n \binom{n}{k} x^k$$

- Proof: Exercise
- Applications:


$$- \binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \cdots + \binom{n}{n} = 2^n \quad (\text{take } x = 1)$$

可用来证明任意有限集合的 odd subset 和 even subset 一样多


$$- \binom{n}{0} + \binom{n}{1} + \binom{n}{2} - \binom{n}{3} \cdots = \sum_{k=0}^n \binom{n}{k} (-1)^k = 0$$

$$- 2 \left[ \binom{n}{0} + \binom{n}{2} + \binom{n}{4} + \cdots \right] = 2^n$$


# Pascal's Triangle (1654) / 杨辉三角 (1261)







(Un-)Ordered sequence




Ordered sequence

- With 5 different red balls, 3 different yellow balls, 4 different blue balls, we can get  $(5 + 3 + 4)! = 12!$  different sequences.



- Question:** With 5 equal red balls, 3 equal yellow balls, 4 equal blue balls, how many different sequences can we get?



- **Theorem:** if we have objects of  $m$  kinds,  $k_i$  indistinguishable objects of  $i$ th kind, where  $k_1 + k_2 + \cdots + k_m = n$ , then the number of distinct arrangements of the objects in a row is  $\frac{n!}{k_1!k_2!\cdots k_m!}$ . Usually written  $\binom{n}{k_1, k_2, \dots, k_m}$ .



$$\frac{12!}{5!3!4!} \text{ 种}$$

- **Multinomial Theorem:** For arbitrary real number  $x_1, x_2, \dots, x_m$  and any natural number  $n \geq 1$ , the following equality holds:

$$(x_1 + x_2 + \dots + x_m)^n = \sum_{\substack{k_1 + \dots + k_m = n \\ k_1, \dots, k_m \geq 0}} \binom{n}{k_1, k_2, \dots, k_m} x_1^{k_1} x_2^{k_2} \dots x_m^{k_m}.$$

- e. g. In  $(x + y + z)^{10}$  the coefficient of  $x^2 y^3 z^5$  is  $\binom{10}{2, 3, 5} = 2520$ .

Basic counting

Binomial theorem

Generalized Binomial theorem

Some special numbers

## Newton(1665)'s generalized binomial theorem

Let  $\binom{r}{k} = \frac{r(r-1)\cdots(r-k+1)}{k!} = \frac{(r)_k}{k!}$  where  $r$  is arbitrary,  
 $k > 0$  is an integer

If  $x$  and  $y$  are real numbers with  $|x| > |y|$

$$(x + y)^r = \sum_{k=0}^{\infty} \binom{r}{k} x^{r-k} y^k$$

$$\begin{aligned} &= x^r + rx^{r-1}y + \frac{r(r-1)}{2!}x^{r-2}y^2 \\ &\quad + \frac{r(r-1)(r-2)}{3!}x^{r-3}y^3 + \dots \end{aligned}$$

$$\sqrt{1+x} = 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{16}x^3 - \frac{5}{128}x^4 + \frac{7}{256}x^5 - \dots$$

$$(1+x)^{-1} = \frac{1}{1+x} = 1 - x + x^2 - x^3 + x^4 - x^5 + \dots$$

Generally:  $r = -s$

$$\frac{1}{(1-x)^s} = \sum_{k=0}^{\infty} \binom{s+k-1}{k} x^k$$

$$\frac{1}{\sqrt{1+x}} = 1 - \frac{1}{2}x + \frac{3}{8}x^2 - \frac{5}{16}x^3 + \frac{35}{128}x^4 - \frac{63}{256}x^5 + \dots$$



Basic counting

Binomial theorem

Generalized Binomial theorem

Some special numbers


# Stirling subset numbers

- The **second Stirling Numbers**  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ : The number of ways to partition a set of  $n$  things into  $k$  **nonempty subsets**.

- e.g.  $\left\{ \begin{smallmatrix} 4 \\ 2 \end{smallmatrix} \right\} = 7$


- $\left\{ \begin{smallmatrix} n \\ 2 \end{smallmatrix} \right\} = 2^{n-1} - 1$  why?

注意 $2^{n-1}$ 的前面没有系数 $n$ , 这是因为最开始选择的红色圆点虽然有 $n$ 种选法, 但是后面再和其他子集合并的时候, 由于集合元素的无序性, 会导致重复, 所以前面的系数应该去掉



# Stirling subset numbers


- The second Stirling Numbers  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ : The number of ways to partition a set of  $n$  things into  $k$  nonempty subsets.
- e.g.  $\left\{ \begin{smallmatrix} 4 \\ 2 \end{smallmatrix} \right\} = 7$
- $\left\{ \begin{smallmatrix} n \\ 2 \end{smallmatrix} \right\} = 2^{n-1} - 1$  why?



# Stirling subset numbers

- The second Stirling Numbers  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ : The number of ways to partition a set of  $n$  things into  $k$  nonempty subsets.
- e.g.  $\left\{ \begin{smallmatrix} 4 \\ 2 \end{smallmatrix} \right\} = 7$
- $\left\{ \begin{smallmatrix} n \\ 2 \end{smallmatrix} \right\} = 2^{n-1} - 1$  why?

选取一个元素 $a$ ，在剩下的 $n-1$ 个元素里面任意取一个子集，将其与 $a$ 合并，作为一个子集，剩下的作为另一个集合，这样就得到两个子集，但是在剩下的 $n-1$ 里面不能全选，否则实际上只有一个集合，没有划分为两个  
所以是 $2^{n-1} - 1$



# Stirling subset numbers

- The second Stirling Numbers  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ : The number of ways to partition a set of  $n$  things into  $k$  **nonempty subsets**.
- e.g.  $\left\{ \begin{smallmatrix} 4 \\ 2 \end{smallmatrix} \right\} = 7$
- $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} = k \left\{ \begin{smallmatrix} n-1 \\ k \end{smallmatrix} \right\} + \left\{ \begin{smallmatrix} n-1 \\ k-1 \end{smallmatrix} \right\}$

先选取一个元素 $a$ ，现在分成两种情况：

1.  $a$ 单独作为一个集合，此时需要将剩下的 $n-1$ 元素划分为 $k-1$ 个
2.  $a$ 被合并到某一个集合里面，那么剩下的 $n-1$ 个元素就需要被分成 $k$ 个，最后 $a$ 在这 $k$ 个里面选一个加进去

# Stirling **cycle** numbers

- The first Stirling Numbers  $\begin{bmatrix} n \\ k \end{bmatrix}$ : The number of ways to partition a set of  $n$  things into  $k$  **nonempty cycles**.

- $\begin{bmatrix} n \\ k \end{bmatrix} \geq \left\{ \begin{matrix} n \\ k \end{matrix} \right\}$ , e.g.  $\begin{bmatrix} 4 \\ 2 \end{bmatrix} = 11$

- $\begin{bmatrix} n \\ 1 \end{bmatrix} = (n - 1)!$

相当于 $n$ 个人围成一圈，可以认为先 $n!$ 排列，在围成圈，这样就不用管谁先开始

证明思路： $n!$ 其实是 $\{1, 2, \dots, n\}$ 到自身的不同的双射函数的个数，所以我们只要证明任意一个斯特灵数和一个这样的双射函数——对应就可以了

- $\sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} = n!$  where  $n \in \mathbb{Z}^+$ .

系数 $n-1$ 是因为插入的选择有 $n-1$ 个

1 2 3 4  
1 4 2 3

唯一决定了 $\{1\} \{2 \rightarrow 4 \rightarrow 3 \rightarrow 2\}$ 这样两个圈

- $\begin{bmatrix} n \\ k \end{bmatrix} = (n - 1) \cdot \begin{bmatrix} n - 1 \\ k \end{bmatrix} + \begin{bmatrix} n - 1 \\ k - 1 \end{bmatrix}$

Why?

# $n$ balls are put into $m$ bins

balls per bin	unrestricted	$\leq 1$	$\geq 1$
$n$ distinct balls, $m$ distinct bins.	$m^n$	$(m)_n$	先把 $n$ 个球分成 $m$ 份, 再对这 $m$ 份全排列 $m! \left\{ \begin{smallmatrix} n \\ m \end{smallmatrix} \right\}$
$n$ identical balls, $m$ distinct bins.	非负整数解 $\binom{n+m-1}{m-1}$	$\binom{m}{n}$	正整数解 $\binom{n-1}{m-1}$
$n$ distinct balls, $m$ identical bins.	$\sum_{k=1}^m \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$	$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	$\left\{ \begin{smallmatrix} n \\ m \end{smallmatrix} \right\}$
$n$ identical balls, $m$ identical bins.		$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	

# Partition of a number

- $P_k(n)$ : number of partition the positive integer  $n$  into  $k$  parts.
- e.g.  $P_2(7) = 3 \quad \{\{1,6\}, \{2,5\}, \{3,4\}\}$   
 $P_6(7) = 1 \quad \{\{1,1,1,1,1,2\}\}$
- Number of integral solutions to
 
$$\begin{cases} x_1 + x_2 + \cdots + x_k = n \\ x_1 \geq x_2 \geq \cdots \geq x_k \geq 1 \end{cases}$$
 这个序关系排除了多余的解
- $P_k(n) = P_{k-1}(n-1) + P_k(n-k)$  why?

如果 $x_k = 1$ , 那么就转化为了右边的第一部分, 否则可以在第一个式子左右都减去 $k$



# $n$ balls are put into $m$ bins

balls per bin	unrestricted	$\leq 1$	$\geq 1$
$n$ distinct balls, $m$ distinct bins.	$m^n$	$(m)_n$	$m! \left\{ \begin{matrix} n \\ m \end{matrix} \right\}$
$n$ identical balls, $m$ distinct bins.	$\binom{n+m-1}{m-1}$	$\binom{m}{n}$	$\binom{n-1}{m-1}$
$n$ distinct balls, $m$ identical bins.	$\sum_{k=1}^m \left\{ \begin{matrix} n \\ k \end{matrix} \right\}$	$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	$\left\{ \begin{matrix} n \\ m \end{matrix} \right\}$
$n$ identical balls, $m$ identical bins.	$\sum_{k=1}^m p_k(n)$	$\begin{cases} 1 & n \leq m \\ 0 & n > m \end{cases}$	$p_m(n)$



# Partition of a number

- $P_k(n)$ : number of partition the positive integer  $n$  into  $k$  parts.

- $\sum_{k=1}^m p_k(n) = p_m(n + m)$  why?

理解思路：也是构造一个左右各项——对应的双射

等式的右边相当于先 $m$ 个盒子里面每个里面放一个球，再将剩下的 $n$ 个球分成 $m$ 个正整数解，这样其实和左边是一个意思

# Twelfold way

The twelve combinatorial objects and their enumeration formulas.

<i>f</i> -class	Any <i>f</i>	Injective <i>f</i>	Surjective <i>f</i>
<i>f</i>	<i>n</i> -sequence in <i>X</i> $x^n$	<i>n</i> -permutation in <i>X</i> $x^n$	composition of <i>N</i> with <i>x</i> subsets $x! \left\{ \begin{smallmatrix} n \\ x \end{smallmatrix} \right\}$
<i>f</i> ∘ <b>S</b> <sub><i>n</i></sub>	<i>n</i> -multisubset of <i>X</i> $\binom{x+n-1}{n}$	<i>n</i> -subset of <i>X</i> $\binom{x}{n}$	composition of <i>n</i> with <i>x</i> terms $\binom{n-1}{n-x}$
<b>S</b> <sub><i>x</i></sub> ∘ <i>f</i>	partition of <i>N</i> into ≤ <i>x</i> subsets $\sum_{k=0}^x \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$	partition of <i>N</i> into ≤ <i>x</i> elements $[n \leq x]$	partition of <i>N</i> into <i>x</i> subsets $\left\{ \begin{smallmatrix} n \\ x \end{smallmatrix} \right\}$
<b>S</b> <sub><i>x</i></sub> ∘ <i>f</i> ∘ <b>S</b> <sub><i>n</i></sub>	partition of <i>n</i> into <i>x</i> non-negative parts $p_x(n+x)$	partition of <i>n</i> into ≤ <i>x</i> parts 1 $[n \leq x]$	partition of <i>n</i> into <i>x</i> parts $p_x(n)$

[https://en.wikipedia.org/wiki/Twelfold\\_way](https://en.wikipedia.org/wiki/Twelfold_way)