ACM SMART CARD

JOINT MULTI-FUNCTIONAL STRIKE BATTALION

SELF AID · BUDDY AID

LEVEL 1 · THE 3 BS

There is no first aid in a firefight. Win the fight, then provide aid.



03 AFTER THE FIGHT

Cover first. Tourniquet, pressure bandage, elastic wrap, paracetamol. Back in the fight.

01 MINIMAL SELF AID

Check yourself fast. Can you still fight? Do only what it takes to keep shooting.

02 COMMUNICATE

Tell your team you are hit, how bad, and where you are. Then keep fighting.

Example: "Smith, RED, by the water tower"

SELF AID

1. Win the fight. Return fire, get to cover, then look at yourself.
2. Tourniquet any limb with many wounds.
3. Pressure bandage every other wound.
4. Elastic wrap once the bleeding has stopped.
5. Paracetamol for pain. 1 to 2 pills, works in about 4 minutes, lasts about 50.
6. Call CLS if you cannot stop the bleeding or a limb will not work.
7. Get back in the fight.

CALL CLS FOR

- Any unconscious casualty
- Any chest wound
- Bleeding you cannot stop
- A limb that will not work

BUDDY AID

1. Win the fight. Do not go to a casualty under fire.
2. Drag your buddy to cover. Treat behind cover.
3. Stop the bleeding. Head wounds first, then limbs, then the rest.
4. **Responsive?** Yes: finish bandaging, wrap, paracetamol, done. No: call CLS and keep going.
5. **Breathing?** No: turn the head to clear the mouth and check again.
6. **Pulse at the neck?** No: CPR. Yes: recovery position, stay with them.
7. Tell CLS what happened, how long they have been down, where the tourniquets are, pulse and breathing, and how many paracetamol.

YOU DO NOT

Airway adjuncts, chest seals, needles, IVs, any medication other than paracetamol.

1 BLEEDING

- **Limb with many wounds:** tourniquet. High and tight, above the wounds.
- **Every other wound:** pressure bandage. One per wound.
- **Bleeding stopped:** elastic wrap, so wounds do not reopen.
- Elastic wrap does not stop a bleeding wound on its own.

2 BREATHING

CHECK BREATHING	DO
Not breathing	Head turn, check pulse, call CLS
Shallow	Call CLS
Slow or rapid	Tell CLS
Normal	Recovery position

Blue skin on the face and arms: short of oxygen. Tell CLS.

3 BEATING

1. Check the pulse at the neck.
2. No pulse: CPR for 2 minutes.
3. Check the pulse again.
4. Repeat until the pulse returns or CLS takes over.

No pulse, no recovery position. It blocks CPR.

Weak or missing pulse at the arms or legs: heavy blood loss. Tell CLS.