ACM SMART CARD

JOINT MULTI-FUNCTIONAL STRIKE BATTALION

CLS

LEVEL 2 · MARCH

Treat in MARCH order. Do not move to the next letter until the current one is controlled.

M MASSIVE HEMORRHAGE

- **Limb, many wounds:** tourniquet.
- **Body part, many wounds:** emergency trauma dressing. Not on yourself.
- **Single wound:** pressure bandage.
- **Once stable:** elastic wrap over bandaged and clotted wounds.
- **Bruising, or bleeding that will not stop:** call the Medic for TXA.
- Tourniquet off once the wounds under it are bandaged.

A AIRWAY

PROBLEM	DO
Tongue blocks airway	NPA or OPA, head-tilt chin-lift, or recovery position
Vomit or blood	Head turn if mild. Suction bag
Collapse, long unconscious	Head-tilt chin-lift or recovery position. Medic for i-gel
Chemical inflammation or spasm	Medic. Needs a surgical airway

Suction first. NPA and OPA only go in a clear airway.

R RESPIRATION

INSPECT CHEST	MEANS	DO
Sides uneven	Pneumothorax	Chest seal
Tracheal deviation	Tension pneumothorax	Chest seal, then NCD
Bruising on upper chest	Hemothorax	Chest seal, call Medic
No movement	Not breathing	Check airway and pulse

PULSE OXIMETER

UNDER 80	UNDER 67	UNDER 55
Unconscious	Cardiac arrest	Death

With systolic under 90 the reading can be off by up to 30%.

C CIRCULATION

CHECK	RESULT	DO
Carotid pulse	None	Cardiac arrest drill
Radial or femoral	Weak or none	Medic for fluids
Body parts	Bruising	Medic for TXA

CARDIAC ARREST DRILL

1. Fix what you can: tension pneumothorax (seal and decompress), blocked airway (clear it).
2. Morphine or fentanyl on board: naloxone now.
3. Connect the AED, select AED mode.
4. CPR for 2 minutes or until the AED prompts. Let it re-analyze. Repeat.
5. No AED: CPR for 2 minutes, carotid pulse check, repeat.

H HEAD AND HYPOTHERMIA

- **Unconscious, stable vitals:** ammonia inhalant.
- **Still unconscious:** go back through M, A, R, C.
- **Conscious:** pain, then fractures.

FRACTURES

SAM splint, then elastic wrap over it. Significant swelling is a complex fracture: evacuate. Realignment is the Medic's.

NOT CLS

IV or IO, fluids, blood, vial medications, i-gel, surgical airway, thoracostomy, manual AED, fracture realignment, casualty conversion.

MEDS

MED	FOR	DOSE	ONSET	LASTS	WATCH
Paracetamol	Mild to moderate pain	1 to 2 pills	Under 4 min	About 50 min	
Penthrox inhaler	Moderate to severe pain, short term	1 to 5 uses, 8 per inhaler	Under 3 s	About 4 min	Not with chest injury, slow breathing, low blood volume or altered mental status
Naloxone spray	Opioid overdose	1 to 2 uses	Under 3 s	About 6 min	Opioids last longer. Watch and re-dose
Ammonia inhalant	Unconscious with stable vitals	1 use, 8 per inhalant	Under 3 s	Under 40 s	
ATNA autoinjector	Nerve agent	2 to 3 injections to start	Under 15 s	About 15 min	
Midazolam autoinjector	Seizures from nerve agent	1 injection	Under 30 s	About 20 min	Lowers heart rate and blood pressure