- 1. Let X and Y be two decision problems. Suppose we know that X reduces to Y. Which of the following can we infer? Explain
 - a. If Y is NP-complete then so is X.
 - b. If X is NP-complete then so is Y.
 - c. If Y is NP-complete and X is in NP then X is NP-complete.
 - d. If X is NP-complete and Y is in NP then Y is NP-complete.
 - e. X and Y can't both be NP-complete.
 - f. If X is in P, then Y is in P.
 - g. If Y is in P, then X is in P.
- 2. Consider the problem COMPOSITE: given an integer y, does y have any factors other than one and itself? For this exercise, you may assume that COMPOSITE is in NP, and you will be comparing it to the well-known NP-complete problem SUBSET-SUM: given a set S of n integers and an integer target t, is there a subset of S whose sum is exactly t? Clearly explain whether or not each of the following statements follows from that fact that COMPOSITE is in NP and SUBSET-SUM is NP-complete:
 - a. SUBSET-SUM ≤p COMPOSITE.
 - b. If there is an $O(n^3)$ algorithm for SUBSET-SUM, then there is a polynomial time algorithm for COMPOSITE.
 - c. If there is a polynomial algorithm for COMPOSITE, then P = NP.
 - d. If $P \neq NP$, then **no** problem in NP can be solved in polynomial time.
- 3. Two well-known NP-complete problems are 3-SAT and TSP, the traveling salesman problem. The 2-SAT problem is a SAT variant in which each clause contains at most two literals. 2-SAT is known to have a polynomial-time algorithm. Is each of the following statements true or false? Justify your answer.
 - a. $3-SAT \le p TSP$.
 - b. If $P \neq NP$, then 3-SAT $\leq_{p} 2$ -SAT.
 - c. If $P \neq NP$, then no NP-complete problem can be solved in polynomial time.
- 4. LONG-PATH is the problem of, given (G, u, v, k) where G is a graph, u and v vertices and k an integer, determining if there is a simple path in G from u to v of length at least k. Show that LONG-PATH is NP-complete.

CS 325 - Homework 4

- 5. Graph-Coloring. Mapmakers try to use as few colors as possible when coloring countries on a map, as long as no two countries that share a border have the same color. We can model this problem with an undirected graph G = (V,E) in which each vertex represents a country and vertices whose respective countries share a border are adjacent. A k-coloring is a function $c: V \to \{1, 2, ..., k\}$ such that $c(u) \neq c(v)$ for every edge $(u,v) \in E$. In other words the number 1, 2, ..., k represent the k colors and adjacent vertices must have different colors. The graph-coloring problem is to determine the minimum number of colors needed to color a given graph.
 - a. State the graph-coloring problem as a decision problem K-COLOR. Show that your decision problem is solvable in polynomial time if and only of the graph-coloring problem is solvable in polynomial time.
 - b. It has been proven that 3-COLOR is NP-complete by using a reduction from SAT. Use the fact that 3-COLOR is NP-complete to prove that 4-COLOR is NP-complete.