

Relative-fit and nested difference tests for estimated-weight moment-quadratic estimators

June 20, 2026

Purpose

The estimated-weight sandwich (companion note `weighted_moment_ij_grid`) corrects the parameter covariance by adding a weight-derivative term. This note asks whether the same correction extends to model-comparison tests, and whether the nested difference test follows as a corollary. Three answers, stated up front.

1. The correction is proportional to the pseudo-true residual r_* (Proposition 1). It is identically zero for the goodness-of-fit null, where the focal model is assumed correct, and present for every test whose null does not assume correctness.
2. For distinguishable models the relative-fit (Vuong) variance is a clean corollary: $\omega^2 = c^\top \Gamma c$ with the weight channel carried by an explicit term in c (Proposition 2).
3. The nested, degenerate case is *not* a clean corollary. The weight channel and the bread curvature both deform Satorra's projection, so the naive substitution $\Gamma \mapsto (W - W_r)\Gamma(W - W_r)^\top$ into the U -matrix is wrong (Proposition 3). It needs its own derivation.

1 Recap of the corrected influence

The estimator minimises $F_N(\theta) = \frac{1}{2}r(\theta)^\top \widehat{W}r(\theta)$ with residual $r(\theta) = \eta(\theta) - u$, Jacobian $\Delta = \partial\eta/\partial\theta^\top$, and weight $\widehat{W} = W(u)$ built from the sample moments u . Write g_i for the moment influence function, $\Gamma = \mathbb{E}[g_i g_i^\top]$, and

θ_* , $r_* = \eta(\theta_*) - u_0$ for the pseudo-true value and residual. The estimator is asymptotically linear with influence function $\phi_i = H^{-1}\Delta^\top v_i$, where

$$H = A + C, \quad A = \Delta^\top W \Delta, \quad C = \sum_k (W r_*)_k \nabla^2 \eta_k(\theta_*), \quad v_i = (W - W_r) g_i, \quad (1)$$

and $W_r = [\partial W / \partial u_1 r_*, \dots, \partial W / \partial u_m r_*]$ is the residual-contracted weight derivative. The corrected moment-space meat is $\Omega = (W - W_r) \Gamma (W - W_r)^\top$.

2 The residual governs the correction

Proposition 1. *Every estimated-weight correction in (1), namely the curvature term C and the weight term W_r , is linear in the pseudo-true residual r_* . Hence both vanish exactly when $r_* = 0$. For a test whose null fixes the focal model as correctly specified the correction is identically zero; for a test whose null leaves the focal model misspecified the correction is present.*

Proof. C contracts $W r_*$ and W_r contracts r_* columnwise, so both are $O(\|r_*\|)$ and vanish at $r_* = 0$. The goodness-of-fit null sets $\eta(\theta_*) = u_0$, hence $r_* = 0$. The relative-fit and the not-assuming-correct nested nulls leave $r_* \neq 0$. $\square$

A second, sharper reason closes the goodness-of-fit case. At that null the fitted residual is $r(\hat{\theta}) = O_p(N^{-1/2})$, so the weight fluctuation enters the residual only through $(\hat{W} - W)r(\hat{\theta}) = O_p(N^{-1})$, which is negligible. The estimated weight is therefore asymptotically irrelevant to the fit statistic, which is why the fixed-weight projection theory of [1] and the scaled statistic implemented in practice are already correct under the null.

3 Goodness of fit: the projection, unchanged

For a fixed weight, expand the fitted residual at the goodness-of-fit null. With $H = A$ and $W_r = 0$ there,

$$\sqrt{N} r(\hat{\theta}) = -M \sqrt{N}(u - u_0) + o_p(1), \quad M = I - \Delta A^{-1} \Delta^\top W,$$

so $\text{Var}(\sqrt{N} r(\hat{\theta})) = M \Gamma M^\top$ and the fit statistic $N r^\top W r$ converges to $\sum_j \lambda_j \chi_{1,j}^2$ with $\lambda_j = \text{eig}(U \Gamma)$ and $U = W M = W - W \Delta (\Delta^\top W \Delta)^{-1} \Delta^\top W$ [1]. When $W = \Gamma^{-1}$ the product $U \Gamma$ is idempotent and the law collapses to χ_{df}^2 ; for $W \neq \Gamma^{-1}$ (DWLS, ULS) it is a genuine mixture, the reason the scaled statistic [3] exists. By Proposition 1 the estimated weight contributes nothing here.

4 Relative fit: a clean corollary

Compare two structural models A and B on the same indicators. The weight $\widehat{W} = W(u)$ is a function of the sample moments alone, so it is shared, $W_A = W_B = W$. Let $D = F_A(\widehat{\theta}_A) - F_B(\widehat{\theta}_B)$ be the sample discrepancy difference and r_*^A, r_*^B the two pseudo-true residuals.

Proposition 2. *The discrepancy difference is asymptotically linear, $\sqrt{N} D = N^{-1/2} \sum_i c^\top g_i + o_p(1)$, with*

$$c = -W(r_*^A - r_*^B) + \frac{1}{2}(b^A - b^B), \quad b_l^A = (r_*^A)^\top \frac{\partial W}{\partial u_l} r_*^A, \quad (2)$$

and b_l^B defined likewise. For distinguishable models $\omega^2 = c^\top \Gamma c > 0$ and $\sqrt{N} D / \widehat{\omega} \rightarrow \mathcal{N}(0, 1)$.

Proof. View F_A as a functional of the sampling distribution through u , the minimiser $\widehat{\theta}_A(u)$, and the weight $W(u)$. The minimiser is interior, so the envelope theorem removes the $\widehat{\theta}_A$ channel and

$$\text{IF}_i(F_A) = \left(\frac{\partial F_A}{\partial u} + \frac{\partial F_A}{\partial W} DW(u_0) \right) g_i = (-W r_*^A + \frac{1}{2} b^A)^\top g_i,$$

using $\partial F_A / \partial u = -r^\top W$ and $\partial(\frac{1}{2} r^\top W r) / \partial W$ contracted with $DW[g_i] = \sum_l (\partial W / \partial u_l) g_{i,l}$. Subtracting $\text{IF}_i(F_B)$ gives (2); the central limit theorem gives the normal law. $\square$

The term $-W(r_*^A - r_*^B)$ is the difference of weighted residuals that a fixed-weight account would give. The term $\frac{1}{2}(b^A - b^B)$ is the weight channel, quadratic in each model's own residual and contracted with the weight derivative. It is the testing analogue of W_r and is the new ingredient. Every quantity is already available: r_*^A, r_*^B from the two fits, $\partial W / \partial u$ from the same derivative the standard error uses, and Γ from the stage-one moments. The relative-fit test is therefore a corollary of the standard-error machinery, with no projection involved because the envelope theorem has removed the parameter channel.

5 The nested case does not commute

When B is nested in A the discrepancy difference degenerates under the null that the restriction holds, $r_*^A = r_*^B$ in the relevant directions, so $c = 0$ and $\omega^2 = 0$. The first-order term vanishes and the limiting law is the second-order quadratic form, which is where the projection returns. The question

is whether that quadratic form is Satorra's U -difference with the meat Γ simply replaced by the corrected Ω .

Proposition 3. *Off the goodness-of-fit null the fitted residual fluctuation is*

$$\sqrt{N}(r_A(\hat{\theta}_A) - r_*^A) = -\widetilde{M}_A \sqrt{N}(u - u_0) + o_p(1), \quad \widetilde{M}_A = I - \Delta_A H_A^{-1} \Delta_A^\top (W - W_r^A),$$

with $H_A = A_A + C_A$. The nested difference statistic converges to $\sum_j \lambda_j \chi_{1,j}^2$ with $\lambda_j = \text{eig}[(\widetilde{M}_A^\top W \widetilde{M}_A - \widetilde{M}_B^\top W \widetilde{M}_B)\Gamma]$. The generalised annihilator $\widetilde{M}_A$ carries both the bread curvature C_A and the weight channel W_r^A inside the projector, so it does not equal the fixed-weight annihilator $M_A = I - \Delta_A A_A^{-1} \Delta_A^\top W$. Consequently the law is not $\text{eig}[(U_B - U_A)\Omega]$, and the substitution $\Gamma \mapsto \Omega$ into the U -matrix is incorrect.

Proof. Expand $r_A(\hat{\theta}_A) = r_*^A + \Delta_A(\hat{\theta}_A - \theta_*^A) - (u - u_0) + o_p(N^{-1/2})$ and substitute the corrected influence $\sqrt{N}(\hat{\theta}_A - \theta_*^A) = H_A^{-1} \Delta_A^\top (W - W_r^A) \sqrt{N}(u - u_0) + o_p(1)$ from (1). Collecting terms gives the stated $\widetilde{M}_A$. The second-order term of $N(F_B - F_A)$ is the difference of the quadratic forms $z^\top \widetilde{M}_A^\top W \widetilde{M}_A z$ and $z^\top \widetilde{M}_B^\top W \widetilde{M}_B z$ with $z \sim \mathcal{N}(0, \Gamma)$, whence the eigenvalue characterisation. The fixed-weight reduction sets $C_A = 0$ and $W_r^A = 0$, returning $\widetilde{M}_A = M_A$ and $\lambda_j = \text{eig}[(U_B - U_A)\Gamma]$ of [2]. $\square$

The correction does not factor out of the projection. The weight channel enters the same place as the bread curvature, inside the annihilator, so the meat and the projector are entangled. Computing the nested reference law requires the generalised annihilators $\widetilde{M}$ rather than a meat swap, and that is a separate derivation, not a corollary.

6 Verdict

The testing extension partitions cleanly. The goodness-of-fit test needs no correction (Proposition 1); the prevailing scaled statistic is already right. The relative-fit test for distinguishable models is a corollary of the standard-error machinery, with $\omega^2 = c^\top \Gamma c$ and the weight channel (2) computable from quantities the standard error already forms (Proposition 2). The nested difference test, in the regime that does not assume the larger model correct, is its own project, because the weight channel deforms Satorra's projection rather than its meat (Proposition 3). For the manuscript this argues to keep the estimated-weight standard error and the distinguishable relative-fit test together, and to treat the nested weighted- χ^2 law as a sequel [4, 5].

References

- [1] Satorra, A. (1989). Alternative test criteria in covariance structure analysis: a unified approach. *Psychometrika*, 54(1), 131–151. doi:10.1007/BF02294453.
- [2] Satorra, A. (2000). Scaled and adjusted restricted tests in multi-sample analysis of moment structures. In R. D. H. Heijmans, D. S. G. Pollock, & A. Satorra (Eds.), *Innovations in Multivariate Statistical Analysis* (pp. 233–247). Springer. doi:10.1007/978-1-4615-4603-0_17.
- [3] Satorra, A., & Bentler, P. M. (2001). A scaled difference chi-square test statistic for moment structure analysis. *Psychometrika*, 66(4), 507–514. doi:10.1007/BF02296192.
- [4] Vuong, Q. H. (1989). Likelihood ratio tests for model selection and non-nested hypotheses. *Econometrica*, 57(2), 307–333. doi:10.2307/1912557.
- [5] Merkle, E. C., You, D., & Preacher, K. J. (2016). Testing nonnested structural equation models. *Psychological Methods*, 21(2), 151–163. doi:10.1037/met0000038.
- [6] Lai, K., & Simões, F. (2023). Reflecting on the “robust” standard errors for two-stage structural equation modeling. *Structural Equation Modeling*, 30(5).
- [7] Hall, A. R., & Inoue, A. (2003). The large sample behaviour of the generalized method of moments estimator in misspecified models. *Journal of Econometrics*, 114(2), 361–394. doi:10.1016/S0304-4076(03)00075-0.