

Author: Jonas Semprini Næss

## MAT4110: Introduction to Numerical Analysis

### Problem 1.

a.) Let  $(x_k)$  be a sequence obtained by Newton's method for solving the above equation. Explain why  $\lim_{k \rightarrow \infty} x_k = \xi$  if  $x_0$  is sufficiently near  $\xi$  and show that the sequence converges with at least order  $q = 3$ .

#### Solution:

To see that  $(x_k)$  indeed will converge for a sufficient  $x_0$  we need to analyse how  $\arctan(x)$  behaves in regards to contractivity. Indeed,  $f'(x) = \frac{1}{1+x^2} \leq 1$ ,  $\forall x \in \mathbb{R}$  and  $|\arctan(x)| \leq |x|$  which by the Mean Value Theorem tells us that

$$|f(x) - f(y)| \leq |f'(z)||x - y|, \quad \text{for } z \in (x, y).$$

Where WLOG  $y > x$ . Now notice that if  $x \rightarrow \infty$  then  $f'(x) \rightarrow 0$ , implying that  $\arctan(x)$  would violate the contraction criteria given large enough  $x$ . It is also easy seeing that since  $f'(x)$  is rapidly decreasing on  $\mathbb{R}$ , namely  $0 < f'(x) < |\frac{1}{x}|$ , then as  $x \rightarrow 0 \Rightarrow f'(x) \rightarrow 1$ . Thus if  $x \in (0, r)$  where  $r \ll \infty$  then  $|f'(x)| < 1$ . Now let  $z \in (\xi, r)$ , and denote  $f'(z) = L$ , meaning  $0 < L < 1$ , then

$$\begin{aligned} |f(\xi) - f(r)| &\leq |f'(z)||\xi - r| \\ &= L|\xi - r|. \end{aligned}$$

which shows that  $\arctan(x)$  is a contraction for a sufficient  $r$  which we can set as an upper boundary for  $x_0$ .

#### Order of convergence

When showing the order of convergence we start off by denoting  $h(x) = x_n - \frac{f(x_n)}{f'(x_n)}$ . This gives us the functional iteration

$$x_{n+1} = h(x_n)$$

which if  $x_0$  is sufficiently close to  $\xi$  will converge to  $\xi$  (i.e part (1) of this exercise). To express the order of convergence we firstly look at the difference

$$x_{n+1} - \xi = h(x_n) - h(\xi)$$

which if we Taylor expand  $h(x)$  about the point  $x_n$  we get that

$$x_{n+1} - \xi = h(x_n) - h(\xi) = \frac{1}{q!} h^{(q)}(\eta_n)(x_n - \xi)^q, \quad q \in \mathbb{N}$$

where by the Squeeze Theorem will ensure that when  $x_n \rightarrow \eta \Rightarrow \eta_n \rightarrow \xi$ . Moreover, by continuity we then have that

$$\frac{|x_{n+1} - \xi|}{|x_n - \xi|^q} = \frac{1}{q!} |h^{(q)}(\xi)|.$$

Now since  $r = h(r)$  only if  $f(r) = 0$  we always have that  $h'(r) = 0$  since

$$h'(x) = \frac{f(x)f''(x)}{f'(x)^2}.$$

Thus, for the order of convergence to be at least  $k$  the expression

$$\frac{1}{q!} |h^{(q)}(\xi)| \neq 0$$

for at least  $q$ . Now if we use (1.24) from (S.M) we have that

$$\frac{|x_{n+1} - \xi|}{|x_n - \xi|^q} = \frac{1}{2} \left| \frac{f''(\xi)}{f'(\xi)} \right| = 0$$

$$\Downarrow$$

$$h''(r) = 0$$

which means the order of convergence is at least cubic.<sup>1</sup> ■

b.) *Show that*

$$g(x) := \begin{cases} f(x)/xf'(x) & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$

*is a continuous and symmetric function (meaning that  $g(x) = g(-x)$ ), and that Newton's method (for this particular problem) can be written as*

$$x_{k+1} = x_k(1 - g(x_k))$$

**Solution:**

### Continuity

Since we are dealing with a piecewise function we check that the limits as  $x \rightarrow 0$  coincide.

$$\lim_{x \rightarrow 0} \frac{f(x)}{xf'(x)} = \lim_{x \rightarrow 0} \frac{\arctan(x)(1+x^2)}{x} \stackrel{L'H}{=} \frac{1 - \arctan(x)(\frac{2x}{(1+x^2)^2})}{1} = 1 \quad (1)$$

$$\lim_{x \rightarrow 0} 1 = 1 \quad (2)$$

Meaning  $g(x)$  is continuous.

---

<sup>1</sup>In this specific problem it is actually exactly of order 3

## Symmetry

By definition we have that a symmetric function can be expressed by the equality  $g(x) = g(-x)$ . Thus to check that this holds we simply calculate  $g(x) - g(-x)$ . This gives

$$\begin{aligned} g(x) - g(-x) &= \frac{f(x)}{xf'(x)} - \frac{f(-x)}{-xf'(-x)} \\ &= \frac{\arctan(x)(1+x^2)}{x} + \frac{\arctan(-x)(1+(-x)^2)}{x} \\ &= \frac{\arctan(x)(1+x^2) - \arctan(x)(1+x^2)}{x} \\ &= 0 \end{aligned}$$

where we have used that  $\arctan(-x) = -\arctan(x)$  since it is an odd function.

## Newtons Method

To show that Newtons method can be written thaty way we simply write it out as

$$x_{k+1} = x_k(1 - g(x_k)) = x_k(1 - f(x_k)/x_k(f'(x_k))) = x_k - \frac{f(x_k)}{f'(x_k)}$$

and we are done. ■

c.) *Show that there exists a unique point  $x^* \in (0, \infty)$  such that  $g(x^*) = 2$  and that*

$$g(x) > 2, \quad \forall x \in (x^*, \infty).$$

To show the uniqueness of  $x^*$  we will proceed by doing a proof by contradiction. Suppose there exists two points  $x_1, x_2 \in (0, \infty)$  such that  $g(x_1) = g(x_2) = 2$ . As defined we know that

$$g(x) = f(x)/xf'(x) = \frac{\arctan(x)(1+x^2)}{x} > x, \quad x \in (0, \infty)$$

meaning  $g(x)$  is strictly increasing on  $(0, \infty)$ . Since  $g(x)$  is strictly increasing that means

$$g(x_1) < g(x_2) < \dots < g(x_n), \quad \text{for } x_1 < x_2 < \dots < x_n \in (0, \infty)$$

but we defined  $g(x_1) = g(x_2)$  for two arbitrary values, meaning we arrive at a contradiction and there must then exist a unique point  $x^*$  such that  $g(x^*) = 2$ . Thus, must all points  $x_1, x_2, \dots, x_n \in (x^*, \infty)$  yield  $g(x_k) > 2$ , for  $k \in N$ . ■

d.) *Approximate the value of  $x^*$  numerically to 5 correct decimal digits using an iteration method.*

## **Solution:**

We can approximate  $x^*$  by the following script

---

```

import numpy as np
import scipy.linalg as linalg
import sys
import numpy as np

def g(x):
    return ((x**2 + 1) * np.arctan(x)) / (x)

def dg(x):
    return 1 / x + np.arctan(x) - np.arctan(x) / (x**2)

x_0 = 1.0
epsilon = 1
error = 1e-5
x_n = 0

while epsilon > error:
    x_n = x_0 - (g(x_0) - 2) / dg(x_0)
    epsilon = abs(x_n - x_0)
    x_0 = x_n

print(f"x_star: {x_0:.6f}")

```

---

where we have implemented Newtons method on  $k(x) = g(x) - 2$ . This gives ut the following approximated value

```

print(f"x_star: {x_0:.6f}")
x_star: 1.391745

```

e.) Use the results in 1 b) and c) to show that the Newton sequence diverges if  $x_0$  is far from  $\xi$ , and describe the largest possible non-empty interval  $(a,b)$  that contains  $\xi$  and satisfies that

$$\lim_{k \rightarrow \infty} x_k = \xi \quad \text{for all Newton sequences with } x_0 \in (a,b)$$

**Solution:**

Let  $\varphi(x) = x - \frac{f'(x)}{f(x)} = x - (x^2 - 1) \arctan(x)$ , and define the functional iteration  $x_{n+1} = \varphi(x_n)$  (as in (a.)). Remember that  $\varphi$  is odd,  $\varphi(0) = 0$ , and  $\varphi'(x) < 0$  meaning  $\varphi(x)$  is strictly decreasing.

Now let  $x^*$  be a unique positive integer (like in (c)) satisfying  $\varphi(x^*) = -x^*$ . (since  $\varphi$  is odd) If  $0 < |x| < |x^*| \Rightarrow |\varphi(x)| \leq |x|$ . Hence if  $|x_0| < |x^*| \Rightarrow |x_{n+1}| \leq |x_n|$ ,  $\forall n$  and  $|x_n| \rightarrow \xi$  since  $\varphi$  is strictly decreasing and  $\varphi(0) = 0$  which by continuity of  $\varphi$  implies that  $\varphi(\xi) = \xi$ .

If  $|x_0| = |x^*|$  then  $\{x_n\} = \{x_0, -x_0, x_0, \dots\}$  which would neither converge nor diverge.

In the last case suppose now that  $0 < |x^*| < |x|$ , which would imply  $\varphi(x) \geq |x|$ , meaning  $x_n$  is now non-decreasing. Nor is  $x_n$  is bounded (neither from below or above) which then implies that  $|x_n| \rightarrow \infty$ .

Hence to conclude must  $x_0 \in (-x^*, x^*)$  for  $|x_n| \rightarrow \xi$ . ■

e.) Compute the sequence  $(x_k)_{k=0}^6$  with Newton's method

$$x_{k+1} = x_k - \frac{f(x)}{f'(x)}$$

We compute  $(x_k)_{k=0}^6$  by the following script

---

```
import numpy as np
import scipy.linalg as linalg
import sys
import numpy as np
import math

def f(x):
    return np.arctan(x)

def f_prime(x):
    return 1 / (1 + x**2)

def newton_method_arctan(x0, tolerance=1e-13, max_iterations=100):
    x = x0
    for i in range(max_iterations):
        fx = f(x)
        if abs(fx) < tolerance:
            return x, i
        x = x - fx / f_prime(x)
    return None, max_iterations

# Initial guess for the root
x0 = 1.3

# Call Newton's method to find the root
root, iterations = newton_method_arctan(x0)

if root is not None:
    print(f"Approximate root: {root}")
    print(f"Number of iterations: {iterations}")
else:
    print("Newton's method did not converge within the specified tolerance.")
```

---

which gives the following output .

x\_0 = 1.3

-----

Approximate root: 1.2045171040057576e-14

Number of iterations: 6

x\_0 = 1.4

-----

Newton's method did not converge within the specified tolerance.

This aligns well with the theoretical results from a.), c.) and e.) where we can clearly see that  $x^* \in (1.3, 1.4)$  somewhere.<sup>2</sup>

## Problem 2

Compute the QR factorization of the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 0 & -1 \\ 1 & 1 \end{bmatrix}$$

and use the factorization to find the least squares solution of the following equation

$$Ax = \underbrace{\begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}}_{=b}$$

**Solution:**

Let  $a_1 = (1, 0, 1)^T$ ,  $a_2 = (1, -1, 1)^T$ . By applying Gram-Schmidt it follows that

$$\mathbf{u}_1 = a_1 = (1, 0, 1)$$

$$\mathbf{e}_1 = \frac{\mathbf{u}_1}{\|\mathbf{u}_1\|} = (\sqrt{2}^{-1}, 0, \sqrt{2}^{-1})$$

$$\mathbf{u}_2 = a_2 - (a_2 \cdot \mathbf{e}_1)\mathbf{e}_1$$

$$= (1, -1, 1) - ((1, -1, 1) \cdot ((\sqrt{2}^{-1}, 0, \sqrt{2}^{-1}))(\sqrt{2}^{-1}, 0, \sqrt{2}^{-1}))$$

$$= (1, -1, 1) - \left( \frac{2}{\sqrt{2}^2}, 0, \frac{2}{\sqrt{2}^2} \right)$$

$$= (1, -1, 1) - (1, 0, 1)$$

$$= (0, -1, 0)$$

$$\mathbf{e}_2 = \mathbf{u}_2.$$

---

<sup>2</sup>To be completely exact, the value  $x^* \in (-1.391745..., 1.391745...)$

Then by the principles of QR factorization we have that

$$Q = \begin{bmatrix} \mathbf{e}_1 & \mathbf{e}_2 \end{bmatrix}$$

$$R = \begin{bmatrix} a_1\mathbf{e}_1 & a_2\mathbf{e}_1 \\ 0 & a_2\mathbf{e}_2 \end{bmatrix}.$$

To solve the least squares problem we remember that  $Q$  is an orthonormal matrix which means that  $Q^T = Q^{-1}$ , so we can thus re-write the problem as

$$Ax = b \iff QRx = b \iff Rx = Q^T b.$$

Then insert for  $Q, R, b$

$$\begin{bmatrix} \sqrt{2} & \sqrt{2} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} \sqrt{2}/2 & 0 & \sqrt{2}/2 \\ 0 & -1 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$$

and solve the following linear set of equations

$$\sqrt{2}x_1 + \sqrt{2}x_2 = \sqrt{2} + \frac{3}{2}\sqrt{2} \quad (3)$$

$$x_2 = -1 \quad (4)$$

which gives that

$$x = \begin{bmatrix} 3.5 \\ -1 \end{bmatrix}.$$

Can also solve this on the computer by the following script

---

```
import numpy as np
import scipy.linalg as linalg
import sys
import numpy as np

A = np.array([[1.0, 0, 1], [1, -1, 1]])
B = np.array([2, 1, 3])
x = np.linalg.lstsq(A.T, B)
print(x[0])
```

---

which gives the following output .

```
x = np.linalg.lstsq(A.T, B)
[ 3.5 -1.]
```

## Problem 3

Let  $PA = LU$  be an LU factorization of  $A$ , where  $P$  is a permutation matrix,  $L$  is a unit lower triangular matrix and  $U$  is upper triangular. Explain how the matrices  $P, L$  and

*U can be used to construct a more efficient method for computing  $\det(A)$  and compute the computational cost of your approach. You may use that the cost of the LU factorization of  $A$  is  $\frac{2n^3}{3} + \mathcal{O}(n^2)$  arithmetic operations, the cost of  $\det(P)$  is  $\mathcal{O}(n^2)$  arithmetic operations and that  $\det(P) \neq 0$  (it always takes one of the values  $\pm 1$ ).*

**Solution:**

Given that  $A$  has an  $LU$  factorisation we can write the determinant of  $A$  as

$$\det(A) = \det(P)\det(L)\det(U)$$

where  $L$  is a unit lower triangular matrix,  $U$  is an upper triangular matrix and  $P$  is a permutation matrix. The cost of the product  $\det(L)\det(U) = u_{1,1} \cdot u_{2,2} \cdots u_{n,n}$  is only  $\mathcal{O}(n)$  since it is the product of the main diagonal of  $U$ , and by