

Chapter 3

Inference from posterior distribution

Assumption in order:

- (1) Likelihood,
- (2) Parameters,
- (3) Prior.

The assumption results in a Posterior.

Likelihood

- The probability of the data given the parameters.
(Assumption)

Likelihood

- The probability of the data given the parameters.
(Assumption)
- Which parameters? The parameters you need to define a probability distribution.

Likelihood

- The probability of the data given the parameters.
(Assumption)
- Which parameters? The parameters you need to define a probability distribution.
- The likelihood is not a distribution for the parameters.

Example: Placenta praevia

- For the data:

$$X = \begin{cases} 1 & \text{if the child is girl,} \\ 0 & \text{if the child is boy.} \end{cases}$$

Example: Placenta praevia

- For the data:

$$X = \begin{cases} 1 & \text{if the child is girl,} \\ 0 & \text{if the child is boy.} \end{cases}$$

- The parameter is the probability of a child being a girl, p .

Example: Placenta praevia

- If you observe n_F girls of n births, the distribution given the data, is binomial:

$$\mathbb{P}(n_F|p) = \frac{n!}{n_F!(n - n_F)!} p^{n_F} (1 - p)^{n - n_F}$$

Example: Placenta praevia

- If you observe n_F girls of n births, the distribution given the data, is binomial:

$$\mathbb{P}(n_F|p) = \frac{n!}{n_F!(n - n_F)!} p^{n_F} (1 - p)^{n - n_F}$$

Posterior

From the prior, parameters, and likelihood, one gets posterior:

- By Bayes theorem

$$p(p|n_F) = \frac{p(n_F|p)p(p)}{p(n_F)}$$

Posterior

From the prior, parameters, and likelihood, one gets posterior:

- By Bayes theorem

$$p(p|n_F) = \frac{p(n_F|p)p(p)}{p(n_F)}$$

- Or in words:


$$\text{Posterior} = \frac{\text{Likelihood} \cdot \text{Prior}}{\text{Average Likelihood}}$$

Inference from posterior distribution

- The posterior distribution

$$p(p|n_f) \propto p^{n_f}(1-p)^{n-n_f}\mathbb{I}_{[0,1]}(p)$$

.



Inference from posterior distribution

- The posterior distribution


$$p(p|n_f) \propto p^{n_f}(1-p)^{n-n_f}\mathbb{I}_{[0,1]}(p)$$

.

- What is the probability that p is less than 0.5:

$$\mathbb{P}(p < 0.5|n_f) = \int_0^{0.5} p(p|n_f) dp$$

(here we would need the actual density)



Samples of the posterior

```
print(samples)
[1] 0.47474747 0.40404040 0.38383838 0.06060606
[5] 0.21212121 0.37373737 0.31313131 0.24242424 0.60606061
[10] 0.22222222 .....
```

Samples of the posterior

```
print(samples)
[1] 0.47474747 0.40404040 0.38383838 0.06060606
[5] 0.21212121 0.37373737 0.31313131 0.24242424 0.60606061
[10] 0.22222222 .....
```



```
mean(samples < 0.5)
mean(samples)
var(samples)
quantile(samples, c(0.05, 0.95))
```



```
[1] 0.2751
[1] 0.582734
[1] 0.01888366
      5%      95%
0.1808081 0.6166667
```

Posterior distribution

What is the probability that p is less than 0.5?






Figure: Histogram of draws from the posterior. Blue line is the posterior distribution


Posterior intervals




Posterior intervals



Posterior intervals



Posterior intervals



Sampling from the posterior

```
p_grid <- seq(from = 0, to = 1, length.out = 10)  
print(p_grid)
```

- | | | | | |
|-----|-----------|-----------|-----------|-----------|
| [1] | 0.0000000 | 0.1111111 | 0.2222222 | 0.3333333 |
| [5] | 0.4444444 | 0.5555556 | 0.6666667 | 0.7777778 |
| [9] | 0.8888889 | 1.0000000 | | |

Sampling from the posterior

- Grid approximation of the posterior:

```
p_grid <- seq(from = 0, to = 1, length.out = 10)
```

- .

```
prior <- rep(1, 10)
print(prior)
```

```
[1] 1 1 1 1 1 1 1 1 1 1
```

Grid approximation

- Grid approximation of the posterior:

```
p_grid <- seq(from = 0, to = 1, length.out = 10)
prior <- rep(1, 10)
```

- .

```
nf = 6
n = 10
likelihood <- dbinom(x = nf, size = n, prob = p_grid)
print(likelihood)
```

```
[1] 0.0000000000 0.0002466915 0.0092547850
[4] 0.0569018950 0.1541821742 0.2409096472
[7] 0.2276075801 0.1133711163 0.0157882546
[10] 0.0000000000
```

Grid approximation

- Grid approximation of the posterior:

```
p_grid <- seq(from = 0, to = 1, length.out = 10)
prior <- rep(1, 10)
nf = 6
n = 10
likelihood <- dbinom(x = nf, size = n, prob = p_grid)
```

• .

```
posterior <- likelihood * prior
print(posterior) # sum(posterior) = 0.8182621
```

```
[4] 0.0569018950 0.1541821742 0.2409096472
[7] 0.2276075801 0.1133711163 0.0157882546
[10] 0.0000000000
```


Grid approximation

- Grid approximation of the posterior:

```
p_grid <- seq(from = 0, to = 1, length.out = 10)
prior <- rep(1, 10)
nf = 6
n = 10
likelihood <- dbinom(x = nf, size = n, prob = p_grid)
posterior <- likelihood * prior
```

Grid approximation

- Grid approximation of the posterior:


```
p_grid <- seq(from = 0, to = 1, length.out = 10)
prior <- rep(1, 10)
nf = 6
n = 10
likelihood <- dbinom(x = nf, size = n, prob = p_grid)
posterior <- likelihood * prior
```

• .


```
posterior <- posterior / sum(posterior)
print(posterior) # sum(posterior) = 1
```

```
[4] 0.0695399334 0.1884263806 0.2944162197
[7] 0.2781597338 0.1385510953 0.0192948614
[10] 0.0000000000
```


number of grid points = 10




number of grid points = 20



number of grid points = 50



number of grid points = 100



Sampling from the posterior

- Grid approximation of the posterior:

```
p_grid <- seq(from = 0, to = 1, length.out = 1000)
prior <- rep(1, 1000)
nf = 6
n = 10
likelihood <- dbinom(x = nf, size = n, prob = p_grid)
posterior <- likelihood * prior
posterior <- posterior / sum(posterior)
```

Sampling from the posterior

- Grid approximation of the posterior:

```
p_grid <- seq(from = 0, to = 1, length.out = 1000)
prior <- rep(1, 1000)
nf = 6
n = 10
likelihood <- dbinom(x = nf, size = n, prob = p_grid)
posterior <- likelihood * prior
posterior <- posterior / sum(posterior)
```

- Sampling:

```
samples <- sample(p_grid ,
                  prob = posterior ,
                  size = 10000,
                  replace = T)
```


Calculating $\mathbb{P}(p < 0.5)$

- $\mathbb{P}(p|n = 10, n_f = 6) = \int_0^{0.5} p(p|n = 10, n_f = 6)dp = 0.274$

Calculating $\mathbb{P}(p < 0.5)$

- $\mathbb{P}(p|n = 10, n_f = 6) = \int_0^{0.5} p(p|n = 10, n_f = 6)dp = 0.274$
- Approximation by sampling (p_1, p_2, \dots, p_s) :

$$P_{est} = \frac{1}{10^4} \sum_{i=1}^{10^4} \mathbb{I}_{[0,0.5)}(p_i) = 0.281$$

In code:

```
samples <- sample(p_grid ,  
                  prob = posterior ,  
                  size = 10000 ,  
                  replace = T)  
P_est <- mean(samples < 0.5)
```

Approximation

```
samples <- sample(p_grid ,  
  prob = posterior ,  
  size = s ,  
  replace = T)  
P_est <- mean(samples < 0.5)
```

- What is the effect of the code s (the number of samples) on the approximation?

Approximation

```
samples <- sample(p_grid ,  
  prob = posterior ,  
  size = s ,  
  replace = T)  
P_est <- mean(samples < 0.5)
```

- What is the effect of the code s (the number of samples) on the approximation?
- $s = 2$ three runs of the code

$$P_{est}^1 = 0.5, P_{est}^2 = 0, P_{est}^3 = 0.$$

Approximation

```
samples <- sample(p_grid ,  
  prob = posterior ,  
  size = s ,  
  replace = T)  
P_est <- mean(samples < 0.5)
```

- What is the effect of the code s (the number of samples) on the approximation?
- $s = 2$ three runs of the code

$$P_{est}^1 = 0.5, P_{est}^2 = 0, P_{est}^3 = 0.$$

- $s = 10$ three runs of the code

$$P_{est}^1 = 0, P_{est}^2 = 0.3, P_{est}^3 = 0.2.$$

Approximation

```
samples <- sample(p_grid ,  
  prob = posterior ,  
  size = s ,  
  replace = T)  
P_est <- mean(samples < 0.5)
```

- What is the effect of the code s (the number of samples) on the approximation?
- $s = 2$ three runs of the code

$$P_{est}^1 = 0.5, P_{est}^2 = 0, P_{est}^3 = 0.$$

- $s = 10$ three runs of the code

$$P_{est}^1 = 0, P_{est}^2 = 0.3, P_{est}^3 = 0.2.$$

- $s = 10^5$ three runs of the code

$$P_{est}^1 = 0.272, P_{est}^2 = 0.274, P_{est}^3 = 0.274.$$

Predictions

- I will denote future/unobserved data with a star.
- n_f^* number of girls born, n^* number of child births.

Predictions

- I will denote future/unobserved data with a star.
- n_f^* number of girls born, n^* number of child births.
- For a given \hat{p} , the distribution of n_f^* is the likelihood

$$p(n_f^* | n^*, \hat{p}) = \frac{n^*!}{n^*!(n^* - n_f^*)!} \hat{p}^{n_f^*} (1 - \hat{p})^n$$

Predictions

- I will denote future/unobserved data with a star.
- n_f^* number of girls born, n^* number of child births.
- For a given \hat{p} , the distribution of n_f^* is the likelihood


$$p(n_f^* | n^*, \hat{p}) = \frac{n^*!}{n^*!(n^* - n_f^*)!} \hat{p}^{n_f^*} (1 - \hat{p})^n$$

- In R:

```
p_hat <- 0.5  
nf_star <- rbinom(n = 1000, #number of samples  
  size = 16,  
  prob = p_hat)
```


samples n_f^*

Distribution of $p(n_f^* | n = 16, \hat{p} = 0.5)$



which \hat{p}

Which \hat{p} ?



Predictions

- The answer no single \hat{p} !

Predictions

- The answer no single \hat{p} !
- The predictive distribution prior to observing the data (y, n)

$$p(n_f^* | n^*) = \int p(n_f^* | n^*, p) p(p) dp.$$

Predictions

- The answer no single \hat{p} !
- The predictive distribution prior to observing the data (y, n)

$$p(n_f^*|n^*) = \int p(n_f^*|n^*, p)p(p)dp.$$

- The predictive distribution after observing the data (y, n)

$$p(n_f^*|n^*, n, n_f) = \int p(n_f^*|n^*, p)p(p|n, n_f)dp.$$

Sampling from the posterior

- Prior predictive distribution

```
samples <- runif(n = 10000, min = 0, max = 1)
nf_star  <- rbinom(n = 10000,
                  size = 16,
                  prob = samples)
```

Sampling from the posterior

- Prior predictive distribution


```
samples <- runif(n = 10000, min = 0, max = 1)
nf_star <- rbinom(n = 10000,
                  size = 16,
                  prob = samples)
```

- Posterior predictive distribution

```
nf = 6
n = 10
p_grid <- seq(from = 0, to = 1, length.out = 1000)
prior <- rep(1, 1000)
likelihood <- dbinom(x = nf, size = n, prob = p_grid)
posterior <- likelihood * prior
posterior <- posterior / sum(posterior)
samples <- sample(p_grid,
                  prob = posterior,
                  size = 10000,
                  replace = T)
nf_star <- rbinom(n = length(samples),
                  size = 16,
                  prob = samples)
```


samples n_f^*

$$p(n_f^* | n^* = 16) \text{ and } p(n_f^* | n^* = 16, n = 10, n_f = 6)$$



samples n_f^*

$$p(n_f^* | n^* = 16) \text{ and } p(n_f^* | n^* = 16, n = 10, n_f = 6)$$

