ISOCAM observations of the ρ Ophiuchi cloud: Luminosity and mass functions of the pre-main sequence embedded cluster *,**

S. Bontemps^{1,2}, P. André³, A.A. Kaas^{4,2}, L. Nordh², G. Olofsson², M. Huldtgren², A. Abergel⁵, J. Blommaert⁶, F. Boulanger⁵, M. Burgdorf⁶, C.J. Cesarsky³, D. Cesarsky⁵, E. Copet⁷, J. Davies⁸, E. Falgarone⁹, G. Lagache⁵, T. Montmerle³, M. Pérault⁹, P. Persi¹⁰, T. Prusti⁶, J.L. Puget⁵, and F. Sibille¹¹

- Observatoire de Bordeaux, B.P. 89, 33270 Floirac, France email: bontemps@observ.u-bordeaux.fr
- ² Stockholm Observatory, 133 36 Saltsjöbaden, Sweden
- ³ Service d'Astrophysique, CEA Saclay, 91191 Gif-sur-Yvette, France
- ⁴ ESA/ESTEC, Astrophysics Division, Netherlands
- $^{5}\,$ IAS, Université Paris XI, 91405 Orsay, France
- ⁶ ISO/SOC, Astrophysics Division of ESA, Villafranca, Spain
- DESPA, Obs. Paris-Meudon, 5 Pl. J. Janssen, 92195 Meudon, France
- ⁸ JAC, 660 N.A'Ohoku Place, University Park, Hilo, HI 96720, USA
- ⁹ ENS Radioastronomie, 24 Rue Lhomond, 75231 Paris, France
- $^{\rm 10}\,$ IAS, CNR, Area di Ricerca Tor Vergata, 00133 Roma, Italy
- ¹¹ Observatoire de Lyon, 69230 Saint Genis Laval, France

Received June 2000 / Accepted March 2001

Abstract. We present the results of the first extensive mid-infrared (IR) imaging survey of the ρ Ophiuchi embedded cluster, performed with the ISOCAM camera on board the ISO satellite. The main ρ Ophiuchi molecular cloud L1688, as well as the two secondary clouds L1689N and L1689S, have been completely surveyed for point sources at 6.7 μ m and 14.3 μ m. A total of 425 sources are detected in $\sim 0.7 \, \mathrm{deg}^2$, including 16 Class I, 123 Class II, and 77 Class III young stellar objects (YSOs). Essentially all of the mid-IR sources coincide with near-IR sources, but a large proportion of them are recognized for the first time as YSOs. Our dual-wavelength survey allows us to identify essentially all the YSOs with IR excess in the embedded cluster down to $F_{\nu} \sim 10$ –15 mJy. It more than doubles the known population of Class II YSOs and represents the most complete census to date of newly formed stars in the ρ Ophiuchi central region. There are, however, reasons to believe that several tens of Class III YSOs remain to be identified below $L_{\star} \sim 0.2 \text{ L}_{\odot}$. The mid-IR luminosities of most ($\sim 65\%$) Class II objects are consistent with emission from purely passive circumstellar disks. The stellar luminosity function of the complete sample of Class II YSOs is derived with a good accuracy down to $L_{\star} \sim 0.03~{\rm L}_{\odot}$. It is basically flat (in logarithmic units) below $L_{\star} \sim 2 L_{\odot}$, exhibits a possible local maximum at $L_{\star} \sim 1.5 L_{\odot}$, and sharply falls off at higher luminosities. A modeling of the luminosity function, using available pre-main sequence tracks and plausible star formation histories, allows us to derive the mass distribution of the Class II YSOs which arguably reflects the initial mass function (IMF) of the embedded cluster. After correction for the presence of unresolved binary systems, we estimate that the IMF in ρ Ophiuchi is well described by a two-component power law with a low-mass index of -0.35 ± 0.25 , a high-mass index of -1.7 (to be compared with the Salpeter value of -1.35), and a break occurring at $M_{\rm flat}=0.55\pm0.25\,{\rm M}_{\odot}$. This IMF is flat with no evidence for a low-mass cutoff down to at least $\sim 0.06 \, \mathrm{M}_{\odot}$.

Key words. Stars: formation – Stars: low-mass, brown dwarfs – Stars: luminosity function, mass function – Stars: pre-main sequence – ISM: individual objects: ρ Ophiuchi cloud

 $Send \quad offprint \quad requests \quad to {:} \quad S. \quad Bontemps, \quad e{-}mail: \\ bontemps@observ.u{-}bordeaux.fr$

^{*} This work is based on observations with ISO, an ESA project with instruments funded by ESA Member States (especially the PI countries: France, Germany, the Netherlands,

and the United Kingdom) with the participation of ISAS and NASA.

 $^{^{\}star\star}$ Table 1 is only available in electronic form at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via http://cdsweb.u-strasbg.fr/

1. Introduction

Recent observations suggest that most stars in our Galaxy and other galaxies form in compact clusters. In particular, near-IR imaging surveys of nearby molecular cloud complexes have shown that the star formation activity is typically concentrated within a few rich clusters associated with massive dense cores which constitute only a small fraction of the total gas mass available (e.g. Lada, Strom, & Myers 1993, Zinnecker, McCaughrean, & Wilking 1993). These embedded clusters comprise various types of young stellar objects (YSOs) – from still collapsing protostars to young main sequence stars – but are usually dominated in number by low-mass pre-main sequence (PMS) stars, i.e., T Tauri stars. Young clusters provide excellent laboratories for studying the formation and early evolution of stars through the observational analysis of large, genetically homogeneous samples of embedded YSOs. Two key characteristics of these young stellar populations are their luminosity distribution and their mass spectrum, which give important observational constraints on the stellar initial mass function (IMF) in the Galaxy. Observations of young embedded clusters can also help us understand possible links between parent cloud properties and the resulting stellar masses. They are however hampered by: (1) dust extinction from the parent cloud which hides most of the newly formed stars at optical wavelengths, (2) the difficulty to recognize the nature of individual sources (e.g. protostars, T Tauri stars, or background sources), (3) the youth (and thus poorly known intrinsic properties) of most cluster members.

Owing to these difficulties, the census of embedded YSOs provided by IRAS and near-IR studies is far from complete even in the nearest clouds (e.g. Wilking, Lada, & Young 1989 – hereafter WLY89). Thanks to its high sensitivity and good spatial resolution in the mid-IR, the ISOCAM camera on board ISO (Cesarsky et al. 1996; Kessler et al. 1996) was a powerful tool to achieve more complete surveys for YSOs in all major nearby star-forming regions (see Nordh et al. 1996, 1998; Olofsson et al. 1999; Persi et al. 2000).

The nearby ρ Ophiuchi cloud is one of the most actively studied sites of low-mass star formation. Its central region harbors a rich embedded cluster with about 100 members recognized prior to the present work (e.g. WLY89, Casanova et al. 1995). While a dispersed population of optically visible young stars, associated with the Upper-Scorpius OB association, has a typical age of several million years (Myr) (e.g. Preibisch & Zinnecker 1999), the central embedded cluster is recognized as one of the youngest clusters known with an estimated age on the order of $\sim 0.3-1$ Myr (e.g. WLY89, Greene & Meyer 1995, Luhman & Rieke 1999). This young cluster has been extensively studied at wavelengths ranging from the X-ray to the radio band. The satellites Einstein and ROSAT have revealed ~ 70 highly variable X-ray sources associated with magnetically-active young stars, including deeply embedded protostellar sources (Montmerle et al. 1983, Casanova et al. 1995, Grosso et al. 2000). In the near-IR, the cloud has been deeply surveyed from the ground using large-format arrays (Greene & Young 1992, Comerón et al. 1993, Strom et al. 1995, Barsony et al. 1997). Unfortunately, due to difficulties in discriminating between background sources and embedded YSOs without performing time-consuming mid-IR photometry (e.g. Greene et al. 1994) or near-IR spectroscopy (e.g. Greene & Lada 1996; Luhman & Rieke 1999), these recent near-IR surveys have only partially increased the number of classified, recognized members. Finally, while only relatively poor angular resolution IRAS data are available so far in the far-IR (e.g. WLY89), deep imaging surveys at an angular resolution of 10-15'' or better exist at (sub)millimeter wavelengths (e.g., Motte, André, & Neri 1998 – hereafter MAN98 -; Wilson et al. 1999).

It is thanks to the illuminating example of the ρ Ophiuchi embedded cluster that the now widely used empirical classification of YSOs was originally introduced. Three IR classes were initially distinguished based on the shapes of the observed spectral energy distributions (SEDs) between $\sim 2 \ \mu \text{m}$ and $\sim 25\text{--}100 \ \mu \text{m}$ (Lada & Wilking 1984, WLY89). Objects with rising SEDs in this wavelength range were classified as Class I, sources with SEDs broader than blackbodies but decreasing longward of $\sim 2~\mu m$ as Class II, and sources with SEDs consistent with (or only slightly broader than) reddened stellar blackbodies as Class III. These morphological SED classes are interpreted in terms of an evolutionary sequence from (evolved) protostars (Class I), to T Tauri stars with optically thick IR circumstellar disks (Class II), to weak T Tauri stars with at most optically thin disks (Class III) (Lada 1987, Adams, Lada, & Shu 1987, André & Montmerle 1994 – hereafter AM94). A fourth class (Class 0) was subsequently introduced by André, Ward-Thompson, & Barsony (1993) to accommodate the discovery in the radio range of cold sources with large submillimeter to bolometric luminosity ratios and powerful jet-like outflows, such as VLA 1623 in ρ Oph A (e.g. André et al. 1990; Bontemps et al. 1996). Class 0 objects, which have measured circumstellar envelope masses larger than their inferred central stellar masses, are interpreted as young protostars at the beginning of the main accretion phase (e.g. André, Ward-Thompson, & Barsony 2000). The fact that the ρ Ophiuchi central region contains at least two Class 0 protostars (André, Ward-Thompson, & Barsony 1993)), as well as numerous ($\gtrsim 60$) pre-stellar condensations (MAN98), demonstrates that it is still actively forming stars at the present time.

The distance to the ρ Ophiuchi cloud is somewhat uncertain. Usually, a value of 160 pc is adopted (e.g. Chini 1981). However, recent Hipparcos results on the Upper-Scorpius OB association (de Zeeuw et al. 1999) provide a reasonably accurate estimate of 145 ± 2 pc for the average distance to the stars of the OB association. The embedded cluster is located at the inner edge of the molecular complex on the outskirts of the OB association (e.g. de Geus 1989), and not very far, in projection, from the as-

sociation center (less than 4° apart which corresponds to ~ 10 pc; see Fig. 9 of de Zeeuw et al. 1999). In this paper, we therefore adopt a distance of $d=140\pm 10$ pc for the ρ Ophiuchi IR cluster which corresponds to a distance modulus $5\times\log_{10}(d/10\text{pc})=5.73\pm0.15$.

The layout of the paper is as follows. Sect. 2 gives observational details ($\S 2.2$) and describes the way the data have been reduced to obtain mid-IR images and extract point-sources (§ 2.3) along with the photometric uncertainties and the sensitivity levels (§ 2.4). In Sect. 3, the identification of detected sources is discussed (§ 3.1) and the selection of a new population of 123 Class II YSOs, as well as 16 Class I and 77 Class III YSOs, is described (\S 3.2-3.5). We then derive luminosities for these YSOs and build the corresponding luminosity functions in Sect. 4. In Sect. 5, we model the luminosity functions for Class II and Class III YSOs in terms of the underlying mass function and star formation history. In Sect. 6, we discuss the resulting constraints on the IMF of the ρ Ophiuchi cloud down to $\sim 0.06\,\mathrm{M}_\odot$, (§ 6.1-6.2) as well as related implications (\S 6.3-6.5).

2. Observations and data reduction

2.1. Region surveyed by ISOCAM

The survey encompasses the ρ Ophiuchi central region associated with the prominent dark cloud L1688, as well as the two subsidiary sites L1689N and L1689S. The L1688 field is a $\sim 45' \times 45'$ square, while the L1689N and L1689S fields each cover an area of $16.5' \times 16'$ (see Fig. 1). Most previously known members of the ρ Ophiuchi cluster lie within these fields. In particular, this is the case for 94 of a total of 113 recognized members from WLY89, AM94, Greene et al. (1994), and MAN98. The known young stars which lie outside the boundaries of our survey are mostly optically visible, weak-line or post T Tauri stars (belonging to Class III) spread over a large area on the outskirts of the molecular complex (e.g. Martín et al. 1998).

2.2. Observational details

The mapping was performed in the raster mode of ISOCAM in which the mid-IR 32×32 pixel array imaged the sky at consecutive positions along a series of scans parallel to the right-ascension axis. The offset between consecutive array positions along each scan $(\Delta\alpha)$ was 15 pixels, while the offset between two scans $(\Delta\delta)$ was 26 pixels. Each set of scans was then co-added and combined into a single raster image. The final image of the L1688 field (Fig. 1 and Abergel et al. 1996) actually results from the combination of six separate rasters. A pixel field of view of 6" was used for four of these rasters, but smaller 3" pixels were employed for the other two rasters in order to avoid saturating the array on the brightest sources of the cluster. The L1689N and L1689S fields were imaged with one raster each using 6" pixels.

In order to avoid saturation, the individual readout time for the L1688 rasters was set to $t_{\rm int}=0.28\,\rm s$. About 55 of these readouts (i.e. an integration time of $\sim 15\,\rm s$) were performed per sky position. Thanks to the half-frame overlap between subsequent individual images, each sky position was observed twice, yielding an effective total integration time of $\sim 30\,\rm s$. For L1689N and L1689S it was possible to use $t_{\rm int}=2.1\,\rm s$, and about 15 readouts were performed per sky position with the same half-frame overlap, giving an integration time per sky position of $\sim 60\,\rm s$. A total of 1104 individual images were necessary to mosaic the L1688 field, and an additional 60 images each were used to map L1689N and L1689S.

All three fields were mapped in two broad–band filters of ISOCAM: LW2 (5–8.5 $\mu \rm m)$ and LW3 (12–18 $\mu \rm m)$. These filters are approximately centered on two minima of the interstellar extinction curve and are situated apart from the silicate absorption bands (at roughly 10 and 18 $\mu \rm m$). However, they include most of the Unidentified Infrared Bands (UIBs, likely due to PAH-like molecules) which constitute a major source of background emission toward star-forming clouds (e.g. Bernard et al. 1993, Boulanger et al. 1996). The ISOCAM central wavelengths adopted here for LW2 and LW3 are 6.7 $\mu \rm m$ and 14.3 $\mu \rm m$ respectively.

2.3. Image processing, source extraction, and photometry

Each raster consists of a temporal series of individual integration frames (i.e. of 32×32 pixel images) which was reduced using the CAM Interactive Analysis software (CIA)¹. We have subtracted the best dark current from the ISOCAM calibration library, and as a second step we improved it with a second order correction using a FFT thresholding method (Starck et al. 1999). Cosmic-ray hits were detected and masked using the multi-resolution median transform algorithm (Starck et al. 1996). The transients in the time history of each pixel due to detector memory effects were corrected with the inversion method described in Abergel et al. (1996). The images were then flat-fielded with a flat image obtained from the observations themselves. Since these various corrections applied to the images are not perfect, the extraction of faint sources from the images is a difficult task. We have developed an interactive IDL point-source detection and photometry program for raster observations which works in the CIA environment. This program helps to discriminate between astronomical sources and remaining low-level glitches or ghosts due to strong transients (see also Nordh et al. 1996: Kaas et al. 2001). The fluxes of the detected sources were estimated from the series of flux measurements made in the individual images (usually 2 to 4 individual images cover each source) which were obtained from classical

¹ CIA is a joint development by the ESA Astrophysics Division and the ISOCAM consortium led by the ISOCAM PI, C. Cesarsky, Direction des Sciences de la Matière, C.E.A., France.




Fig. 1. Sky map of the ρ Ophiuchi clouds L1688, L1689N, and L1689S showing the spatial distribution of the ISOCAM sources. The filled symbols correspond to "red" YSOs (i.e. Class I and Class II YSOs, see Sect. 3), while the open circles mark the "blue" sources. The surface areas of these symbols are proportional to the 6.7μm flux densities. The open stars mark four young early-type stars (SR3, S1, WL22, WL16 from right to left; see Sect. 3.1). The contours show the L1688 molecular cloud as mapped in CS(2-1) by Liseau et al. (1995). The triangles refer to the peaks of the DCO⁺ dense cores of Loren et al. (1990). The figure emphasizes the presence of four sub-clusters Oph A, B, EF (between Oph E and Oph F), and L1689S (named after the associated dense cores).

aperture photometry. The emission was integrated in a sky aperture, the background emission subtracted, and finally an appropriate aperture correction was applied based on observed point-spread functions available in the ISOCAM calibration library. In practice, the radius of the aperture used was 9'' (i.e., 3 and 1.5 pixels for a pixel size of 3''and 6", respectively). For the weakest sources, however, we reduced the aperture radius to 4.5'' (i.e. 1.5 pixels for a pixel size of 3''), in order to improve the signal-to-noise ratio. Finally, we applied the following conversion factors: 2.33 and 1.97 ADU/gain/s/mJy for LW2 and LW3 respectively (from in-orbit latest calibration - Blommaert 1998). These calibration factors are strictly valid only for sources with a flat SED $(F_{\nu} \sim \nu^{-1})$. Here, a small but significant ($\gtrsim 1$ %) color correction needs to be applied to the bluest sources, recognized as Class III YSOs in Sect. 3 below. For these sources, the conversion factors quoted above were divided by 1.05 for LW2 and 1.02 for LW3 to account for the color effect. The 212 ISOCAM sources recognized as cluster members (see Sect. 3 below) are listed in Table 1 (available only in electronic form at http://cdsweb.u-strasbg.fr/) with their J2000 coordinates, their flux densities and associated rms uncertainties (see Sect. 2.4), as well as the corresponding near-IR identifications.

2.4. Photometric uncertainties and point-source sensitivity

The uncertainties on the final photometric measurements result from systematic errors due to uncertainties in the absolute calibration and the aperture correction factors, and from random errors associated with the flat-fielding noise, the statistical noise in the raw data, the noise due to remaining low-level glitches, and the imperfect correction for the transient behavior of the detectors. The inorbit absolute calibration has been verified to be correct to within 5 % (Blommaert 1998), and we estimate that the maximum systematic error on the aperture correction is $\sim 10\%$ (by comparing theoretical and observed pointspread functions). The maximum systematic error on our photometry is thus $\sim 15 \%$. The magnitudes of the random errors were directly estimated from the data by measuring both a "temporal" noise (noise in the temporal sequence of individual integrations) and a "spatial" noise (due to imperfect flat-fielding and/or spatial structures in the local mid-IR background emission) for each source in the automatic detection procedure. The temporal noise was computed as the standard deviation of the individual aperture measurements divided by the square root of the number of measurements. The spatial noise was estimated as the standard deviation around the mean background (linear combination of the median and the mean of the pixels optimized for the source flux estimates) in the immediate vicinity of each source.




Fig. 2. (a) Distribution of the $F_{\nu}^{6.7}$ flux densities for the 425 sources detected by ISOCAM. The dark histogram shows the flux distribution of the cluster members (i.e., YSOs) discussed in Sect. 3, while the light histogram corresponds to the remaining sources which are likely dominated by background Galactic objects. (b) Same as (a) for the $F_{\nu}^{14.3}$ flux densities. The dark histogram shows the flux distribution for the "red" ISOCAM sources (i.e., Class I and II YSOs), while the light histogram comprises only the "blue" ISOCAM sources not associated with recognized YSOs (see Sect. 3 and Fig. 3). The adopted completeness levels equal to 10 and 15 mJy for $F_{\nu}^{6.7}$ and $F_{\nu}^{14.3}$ respectively are shown as vertical dotted lines. The solid and dashed curves show the expected number of Galactic sources in each bin according to the model of Wainscoat et al. (1992) for $A_V = 0$ and $A_V = 10$ respectively.

The sensitivity limit of the survey was estimated by calculating the average value of the quadratic sum of the temporal and spatial noises measured on the weakest detected sources. The total rms flux uncertainty found in this way, $\sigma_{tot} = (\sigma_{temp}^2 + \sigma_{spat}^2)^{1/2}$, is $\sigma_{6.7} = 2.2$ mJy at 6.7 μ m and $\sigma_{14.3} = 4.1$ mJy at 14.3 μ m, ~ 75 % of which is due to the spatial noise component. The large contribution of the spatial noise originates in the highly structured diffuse mid-IR emission from the ambient molecular cloud itself (see Abergel et al. 1996). Figure 2 displays the distributions of fluxes at 6.7 μ m and 14.3 μ m for all the detected ISOCAM sources. We used the Wainscoat et al. (1992) Galactic model of the mid-IR point source sky to estimate the expected number of foreground and background sources up to a distance of 20 kpc. The model predictions are shown by solid and dashed curves in Fig. 2 for cloud extinctions of $A_V = 0$ and $A_V = 10$, respectively (see Kaas et al. 2001 for more details). It can be

seen that the flux histograms of the ISOCAM sources not associated with YSOs (light shading in Fig. 2) are remarkably similar in shape to the model distributions down to ~ 6 mJy at 6.7 μ m and ~ 10 mJy at 14.3 μ m. These flux densities can be used to estimate the completeness level of our observations which is not uniform over the spatial extent of the survey. The histograms with light (grey) shading in Fig. 2 are dominated by background sources preferentially located in low-noise regions (i.e., outside the crowded central part of L1688), where the total rms flux uncertainty is ~ 2.0 mJy at 6.7 μm and ~ 3.5 mJy at 14.3 μ m. The effective completeness level in these regions is thus $\sim 3 \sigma_{tot}$, where σ_{tot} is the total² flux uncertainty (see above). However, most of the YSOs are located in regions where the noise is somewhat larger. The largest rms noise is reached in the Oph A core area (see Fig. 1), where $\sigma_{tot} \approx 3.4 \, \mathrm{mJy}$ at 6.7 $\mu \mathrm{m}$ and $\sigma_{tot} \approx 5.0 \, \mathrm{mJy}$ at 14.3 $\mu \mathrm{m}$. Therefore, we conservatively estimate the completeness levels of the global ISOCAM survey to be ~ 10 mJy at 6.7 μ m and \sim 15 mJy at 14.3 μ m.

Finally, we note that the $A_V=10$ model curve in Fig. 2b accounts for essentially all the "blue" sources detected at 14.3 μ m and not associated with known YSOs. At 6.7 μ m, the predictions of the Wainscoat et al. model suggest that there might still be a slight excess of ~ 30 unidentified sources belonging to the cloud (Fig. 2a).

3. Identification and nature of the mid-IR sources

3.1. Statistics of detections

Within the ~ 0.7 square degree imaged by ISOCAM, a total of 425 sources has been identified, among which 211 are seen at both 6.7 μ m and 14.3 μ m. The spatial distribution of the sources is shown in Fig. 1, where the "red" sources [those with $\log_{10}(F_{\nu}^{14.3}/F_{\nu}^{6.7}) > -0.2$, see below] are indicated as filled circles. These "red" sources appear to be clustered into four main groupings: three sub-clusters in L1688, i.e., Oph A (West), Oph B (North-East), Oph EF (South) (see also Strom, Kepner, & Strom 1995), as well as a new sub-cluster in L1689S.

Four bright embedded stars (S1, SR3, WL16, WL22) are spatially resolved by ISOCAM in both filters. Their extended mid-IR emission is most likely due to PAH-like molecules excited by relatively strong far-ultraviolet (FUV) radiation fields³. These bright sources are displayed as open star symbols in Fig. 1 (SR3, S1, WL22, WL16 from right to left).

The detection level is more directly related to the temporal noise than to the spatial noise. In terms of σ_{temp} , the effective completeness level is $\sim 12 \, \sigma_{temp}$.

³ S1 and SR3 are optically-visible stars of spectral types B3 and B7, respectively, which clearly emit enough UV photons to excite PAHs. The other two objects, WL16 and WL22, show PAH features in their mid-IR spectra (e.g. Moore et al. 1998), and may be young embedded early-type (i.e., early A or late B) stars.

A total of 89 previously classified YSOs lie within the area of the present survey: 2 Class 0, 69 Class I/II, and 18 Class III YSOs. ISOCAM detected 84 of these 89 YSOs (94 %): 97 % of the Class I/IIs, 94 % of the Class IIIs, and none of the Class 0s. The two undetected Class I/IIs (GY256, GY257), and the undetected Class III (IRS50), are located very close ($\sim 10''-15''$) to bright sources (WL6 and IRS48, respectively), which may account for their non-detection. The two Class 0 objects (VLA 1623 and IRAS 16293–2422) are deeply embedded within massive, cold circumstellar envelopes which are probably opaque at 6.7 μ m and 14.3 μ m (see André, Ward-Thompson, & Barsony 1993)) and too weak to be detected by ISOCAM.

3.2. A population of new YSOs with mid-IR excess

As can be seen in Fig. 3, the mid-IR regime is ideal to detect and characterize the excess emission due to circumstellar disks around young stars. In Fig. 3a, the 207 sources detected at 6.7 μ m and 14.3 μ m (excluding S1, SR3, WL16, WL22) are shown in a diagram which displays the logarithmic flux ratio $\log_{10}(F_{\nu}^{14.3}/F_{\nu}^{6.7})$ against the mid-IR flux $F_{\nu}^{14.3}$. On the right axis of the diagram, the flux ratio $F_{\nu}^{14.3}/F_{\nu}^{6.7}$ has been converted into a classical IR spectral index, $\alpha_{IR} = d\log_{10}(\lambda F_{\lambda})/d\log_{10}(\lambda)$ (e.g. WLY89), calculated between 6.7 μm and 14.3 μm , i.e., α_{IR}^{7-14} . In this diagram, two groups of sources can clearly be distinguished. The lower ratio group has $\log_{10}(F_{\nu}^{14.3}/F_{\nu}^{6.7}) \sim -0.7$, i.e., a spectral index on the order of $\alpha_{\rm IR}^{7-14} \sim -3.0$, which is the value expected for simple photospheric blackbody emission in the Rayleigh-Jeans regime. The dispersion around $\alpha_{\rm IR}^{7-14}\sim-3.0$ is obviously larger for weaker sources. This is mainly due to increasing photometric uncertainty with decreasing flux. The higher ratio group consists of "red" sources defined by $\log_{10}(F_{\nu}^{14.3}/F_{\nu}^{6.7}) > -0.2$ (cf. Nordh et al. 1996) . Most of these actually have $-0.2 \leq \log_{10}(F_{\nu}^{14.3}/F_{\nu}^{6.7}) \leq 0.3$, i.e., $-1.6 \lesssim \alpha_{\rm IR}^{7-14} \lesssim 0$, which is typical of classical T Tauri stars (hereafter CTTS). This range of $\alpha_{\rm IR}^{7-14}$ roughly delineates the domain of Class H VCO (c. a. Adams et al. 1987, AM04) main of Class II YSOs (e.g. Adams et al. 1987, AM94, Greene et al. 1994), and is usually interpreted in terms of optically thick circumstellar disk emission (e.g. Lada & Adams 1992). In Fig. 3, the new ISOCAM sources are distinguished from the previously known cluster members by different symbols. One can see in Fig. 3a that all the previously known Class II sources but one lie above the dividing line for red sources, while all the previously known Class III sources but two lie below it.

The object WL19, classified as a Class I YSO by WLY89 and as a reddened Class II by AM94 (see also Lada & Wilking 1984), is here found to be a "blue" source in the mid-IR range ($\alpha_{\rm IR}^{7-14} = -1.8 \pm 0.2$). This object may correspond to a luminous Class III star located behind the cloud (see also Comerón et al. 1993). Although GY12 formally has a "red" mid-IR spectral index ($\alpha_{\rm IR}^{7-14} = -0.5 \pm 0.5$) here, we still consider it as a Class III object (cf. Greene et al. 1994). (The mid-IR

color is highly uncertain since GY12 is only marginaly resolved at 14.3 $\mu \rm m$ from its bright Class I neighbor GSS30.) Finally the Class III source DoAr21 has a borderline mid-IR spectral index ($\alpha_{\rm IR}^{7-14}=-1.5\pm0.1$) but is kept as a Class III object since its color between 2.2 $\mu \rm m$ and 14.3 $\mu \rm m$ corresponds to $\alpha_{\rm IR}^{2-14}=-2.0\pm0.1$.

A total of 71 sources are identified for the first time as mid-IR excess objects in Fig. 3a. These new "red" sources are most likely all embedded YSOs, i.e., members of the ρ Ophiuchi cluster. Since dust extinction is roughly the same at 6.7 μ m and 14.3 μ m (Rieke & lebofsky 1985; Lutz 1999), any background source should be intrinsically red in order to contaminate the sample of red YSOs. Based on the Galactic model by Wainscoat et al. (1992), the vast majority of background objects should appear "blue" (cf. Fig. 2). Seven background giant stars (GY45, GY65, VSSG6, GY232, GY351, GY411, and GY453 – cf. Luhman & Rieke 1999), and a known foreground dwarf (HD148352 – Garrison 1967) are detected, which all have blue mid-IR colors ($\alpha_{\rm IR}^{7-14} = -2.7, -2.0, -3.1, -3.3, -2.9, -3.2, -3.1,$ and -3.1, respectively).

The newly identified cloud members nicely extend the previously known Class I/II population toward low IR fluxes. While previous studies could only identify sources with $F_{\nu}^{14.3}\gtrsim 100$ mJy, the present census is complete for objects down to $F_{\nu}^{14.3}\sim 15$ mJy. Altogether, a sample of 139 Class I/Class II YSOs is identified, of which 71 are new members. The present survey has thus allowed us to more than double the number of recognized YSOs with circumstellar IR excess in the ρ Ophiuchi cloud.

The 139 red YSOs are listed in Table 2 (Class I YSOs) and Table 3 (Class II YSOs) by decreasing order of $\alpha_{\rm IR}^{2-14}$. Twelve of them (last entries of Table 3) are completely new sources with respect to published IR surveys.

3.3. Class I versus Class II YSOs

In Fig. 3, the YSOs classified as Class I and Class II by AM94 and Greene et al. (1994) are shown as filled stars and filled circles, respectively. In the $\log_{10}(F_{\nu}^{14.3}/F_{\nu}^{6.7})$ versus $F_{\nu}^{14.3}$ diagram, there is no clear color gap between the two classes of objects, even though the Class I YSOs tend to lie in the upper part of the "red" group (see Fig. 3a). Since extinction has a negligible effect on the mid-IR ratio $F_{\nu}^{14.3}/F_{\nu}^{6.7}$, this suggests that Class I and Class II objects have fairly similar intrinsic colors between 6.7 μ m and 14.3 μ m.

The classical IR spectral index calculated from 2 μ m to 10 μ m (or 25 μ m) (e.g. Lada & Wilking 1984 and WLY89) appears to provide a better way of discriminating between envelope-dominated Class I YSOs and disk-dominated Class II sources (see Fig. 3b). In particular, millimeter continuum mapping of optically thin circumstellar dust emission confirms that, apart from a few important exceptions (e.g., WL22, WL16, WL17, IRS37, IRS47), the ρ Ophiuchi objects selected on the basis of $\alpha_{\rm IR}^{2-10} \gtrsim 0$ are indeed Class I protostars surrounded by spheroidal en-




Fig. 3. (a) Logarithmic ratio of the ISOCAM fluxes, $\log_{10}(F_{\nu}^{14.3}/F_{\nu}^{6.7})$, versus $F_{\nu}^{14.3}$ displayed for the 207 ISOCAM sources seen in both filters (excluding the 4 early-type young stars). The purely ISOCAM color $F_{\nu}^{14.3}/F_{\nu}^{6.7}$ is converted into an IR spectral index, $\alpha_{\rm IR}^{7-14}$, on the right axis, and can be used to identify sources with mid-IR excesses (above the dashed line). The completeness limit derived in Sect. 2.4 is displayed as a dotted line. (b) Mid-IR color-color diagram expressed as a spectral index diagram $\alpha_{\rm IR}^{2-7}$ versus $\alpha_{\rm IR}^{2-14}$ for 175 ISOCAM sources also detected in the K band (m_{6.7} and m_{14.3} on the right and upper axes are calculated as in Persi et al. 2000). The inclined heavy dashed line marks the limit to the right of which there is significant mid-IR excess as in (a). The two vertical dashed lines mark the formal boundary between Class II, flat-spectrum, and Class I sources (see text for details). (c) Classical near-IR color-color diagram, J–H versus H–K, for the 152 ISOCAM sources with JHK photometry.

velopes (AM94, MAN98).

As expected, the previously known Class I YSOs are well concentrated in the upper-right part of the $\alpha_{\rm IR}^{2-7}$ versus $\alpha_{\rm IR}^{2-14}$ diagram of Fig. 3b (where $\alpha_{\rm IR}^{2-7}$ and $\alpha_{\rm IR}^{2-14}$ are close to the $\alpha_{\rm IR}^{2-10}$ index used in previous studies – e.g., AM94 and Greene et al. 1994). There is also a hint of two gaps in this diagram at $\alpha_{\rm IR}^{2-14}\sim 0.5$ and $\alpha_{\rm IR}^{2-14}\sim 0.0$, which roughly bracket the regime of flat-spectrum sources as defined by Greene et al. (1994). These may represent a distinct population of transition objects between Class I and Class II (e.g. Calvet et al. 1994).

Here, we thus consider sources with $\alpha_{\rm IR}^{2-14}>0.55$ as Class I YSOs (Table 2), sources with $-0.05<\alpha_{\rm IR}^{2-14}<0.55$ as candidate flat-spectrum objects (see Table 3), and sources with $\alpha_{\rm IR}^{2-14}<0.55$ as Class II YSOs (Table 3). These limiting indices are displayed in Fig. 3b. In the following, the candidate flat-spectrum sources will be treated as Class II YSOs.

3.4. Mid-IR excess versus near-IR excess

Most of the ISOCAM sources (e.g. 90 % of the Class II sources) were also detected in the near-IR (JHK) survey of Barsony et al. (1997). Comparison between Fig. 3b and Fig. 3c illustrates the advantage of mid-IR measurements for selecting sources with intrinsic circumstellar IR excesses. While the red and blue groups of Fig. 3a are well separated in the mid-IR diagram of Fig. 3b, they blend together in the near-IR diagram of Fig. 3c.

Figure 3c also shows that most of the ISOCAM-selected YSOs lie within the reddening band associated with the intrinsic locus of CTTSs as derived by Meyer et al. (1997) in Taurus. The few exceptions, which lie to the right of the reddening band, correspond to Class I and flat-spectrum YSOs.

3.5. The Class III YSO census in ρ Ophiuchi

While Class I/II YSOs are easily recognized in the mid-IR range thanks to their strong IR excesses, Class III objects are difficult to identify without deep X-ray and/or radio centimeter continuum observations. We have used the ROSAT X-ray surveys of Casanova et al. (1995) and Grosso et al. (2000), along with the VLA radio surveys by, e.g., André, Montmerle, & Feigelson (1987) and Stine et al. (1988) to build up a sample of bona-fide Class III YSOs covered by the present survey. With the additional Class III candidate WL19 (see Sect. 3.2 above), there are 38 such YSOs which are listed in order of decreasing $\alpha_{\rm IR}^{2-14}$ in Table 4.

This Class III sample is unfortunately not as complete as the Class I and Class II samples discussed above. According to Grosso et al. (2000), the number of Class IIIs may be roughly as large as the number of Class IIs: above their typical X-ray detection limit of $L_X \sim 3 \times 10^{29} \,\mathrm{erg \, s^{-1}}$ (corresponding to $L_{\star} \sim 0.3 L_{\odot}$ – see Fig. 7 of Grosso et al.), they found a Class III/Class II number ratio of 19/22 in the ROSAT-HRI/ISO-ISOCAM overlapping survey area. If this ratio is representative of the complete population of young stars in ρ Ophiuchi, the total number of Class IIs found here (123 objects) suggests that as many as 106 Class IIIs may be present in the cluster down to $L_{\star} \sim 0.03 \, \rm L_{\odot}$ (our completeness level for Class IIs, see Sect. 4.4). A total of 38 Class IIIs are already known within the ISOCAM survey area, so that ~ 68 unknown Class IIIs may remain to be found. Since it was noted in Sect. 2.4 that ~ 30 sources detected at 6.7 μ m might be unidentified cluster members, about half of the missing Class IIIs may have been actually seen by ISOCAM.

We also note that a large proportion (80/123) of the Class II sources are closely associated with the densest part of L1688 (see Fig. 1). Assuming the same proportion applies to Class IIIs, we would expect \sim 44 unknown Class III sources to be located within the CS contours of Fig. 1. A total of 39 unclassified ISOCAM sources (also detected by Barsony et al. 1997) lie within these CS contours where the number of detected background stars should be small due to high cloud extinction. Most of these 39 sources might thus be yet unidentified Class III YSOs. These candidate Class III sources are listed in order of decreasing $\alpha_{\rm IR}^{2-14}$ in Table 5.

4. Luminosity estimates and luminosity function

Most of the new YSOs identified by ISOCAM are weak IR sources which were not detected by IRAS and were not observed in previous ground-based mid-IR surveys (dedicated to bright near-IR sources). They likely correspond to low-luminosity, low-mass young stars. In Sect. 4.1 below, we derive stellar luminosity estimates for Class II and Class III objects using published near-IR photometry from Barsony et al. (1997). In Sect. 4.2, we provide mid-IR estimates of the disk luminosities, $L_{\rm disk}$, for Class II YSOs. Finally, calorimetric estimates of the bolometric luminosi-

ties, $L_{\rm bol}$, for Class I YSOs are calculated in Sect. 4.3. The luminosity function of the ρ Ophiuchi embedded cluster is then assembled in Sect. 4.4.

4.1. Dereddened J-band stellar luminosities for Class II and Class III sources

The J-band flux provides a good tracer of the stellar luminosity for late-type PMS stars (i.e., T Tauri stars) because the J-band is close to the maximum of the photospheric energy distribution for such cool stars. It is also a good compromise between bands too much affected by interstellar extinction at short wavelengths (very few ρ Ophiuchi YSOs have been detected in the V, R, or I bands), and the H, K and mid-IR bands which are contaminated by intrinsic excesses. Greene et al. (1994) showed that there is a good correlation between the dereddened J-band flux and the stellar luminosity derived by other methods. They pointed out that in ρ Ophiuchi this correlation is roughly consistent with a more theoretically based correlation expected for 1-Myr old PMS stars following the D'Antona & Mazzitelli (1994) evolutionary tracks. More recently, Strom et al. (1995) and Kenyon & Hartmann (1995) used the same model PMS tracks to directly convert dereddened J-band fluxes into stellar masses. We adopt a similar approach here.

4.1.1. Extinction estimates from near-IR colors

The main difficulty and source of uncertainty with this method is due to the foreground extinction affecting the J-band fluxes. One must estimate the interstellar extinction toward each source in order to correct the observed J-band fluxes. We have used the observed near-IR colors to estimate the J-band extinction. The (J-H) color excess is most suitable for this purpose (e.g. Greene et al. 1994) since the dispersion in the intrinsic (J-H) colors of CTTSs is small and observationally well determined (cf. Strom et al. 1989, Meyer et al. 1997).

The reddening law quoted by Cohen et al. (1981), which is determined for the standard CIT system, should be applicable to the JHK photometry of Barsony et al. (1997). We have thus used:

$$A_V = 9.09 \times [(J - H) - (J - H)_0] \tag{1}$$

when the (J-H) color is available, and

$$A_V = 15.4 \times [(H - K) - (H - K)_0] \tag{2}$$

otherwise, where we have adopted $(J-H)_0=0.85$, and $(H-K)_0=0.55$ from the CTTS results of Meyer et al. (1997). The dereddened J_0 magnitude is then obtained as $J_0=J-0.265\times A_V$, or $H_0=H-0.155\times A_V$ for those stars not detected in J (with A_V estimated from H–K). The dereddened J_0 (or H_0) can then be converted into an absolute J-band (H-band) magnitude, M_J (M_H), via the distance modulus of the ρ Ophiuchi cloud equal to 5.73 (for d=140 pc).

The uncertainties on M_J and M_H result from the typical uncertainties on the J, H, K magnitudes and on the intrinsic colors $(J-H)_0$ and $(H-K)_0$. With $\sigma(J)=0.1$, $\sigma(H)\sim\sigma(K)=0.05$ (Barsony et al. 1997), and $\sigma((J-H)_0)=0.1$, $\sigma((H-K)_0)=0.2$ (Meyer et al. 1997), we obtain the following typical uncertainties: $\sigma(M_J)=0.39$ mag, and $\sigma(M_H)=0.60$ mag. In addition, the uncertainty on the cluster distance $(140\pm 10~{\rm pc})$ induces a maximum systematic error of ± 0.15 mag on M_J and M_H .

4.1.2. Relationship between M_J and L_\star for young T Tauri stars

The absolute J-band magnitude M_J can be directly converted into a stellar luminosity L_{\star} if the effective stellar temperature $T_{\rm eff}$ is known: $\log_{10}(L_{\star}) = 1.89 - 0.4 \times$ $(M_J + BC_J(T_{\text{eff}}))$, where BC_J is the bolometric correction for the J band depending only on $T_{\rm eff}$. Pre-main sequence objects in the mass range $0.1 \lesssim M_{\star} \lesssim 2.5 M_{\odot}$ are cool sub-giant stars with typical photospheric temperatures $T_{\rm eff} \sim 2500-5500$ K (e.g. Greene & Meyer 1995, D'Antona & Mazzitelli 1994). In this temperature range (0.34 dex wide), the photospheric blackbody peaks close to the J band $(1.2\mu m)$, so that the J-band bolometric correction spans only a limited range, $1 \lesssim BC_J \lesssim 2$, corresponding to a total shift in luminosity of only 0.4 dex. Therefore, if we use a (geometrical) average value for the effective temperature, $T_{\rm eff}^{~0} \sim 3700$ K, we should not make an error larger than ± 0.2 dex on $\log_{10}(L_{\star})$.

Furthermore, since bright M_{J} sources tend to be of earlier spectral type than faint sources, we can in fact achieve more accurate luminosity estimates. Indeed, PMS stars are predicted to lie within a well-defined strip of the HR diagram. This is illustrated in Fig. 4 which displays model evolutionary tracks and isochrones from D'Antona & Mazzitelli (1998) on a M_J -log₁₀($T_{\rm eff}$) diagram for PMS stars with ages between 0.1 Myr and 3 Myr. We have used the compilations of Hartigan et al. (1994), Kenyon & Hartmann (1995), and Wilking et al. (1999) to derive an approximate linear interpolation for BC_J : $BC_J = 1.65 + 3.0 \times \log_{10}(3700/T_{\text{eff}})$. Figure 4 shows that, for a given observed value of M_J , the possible range of $\log_{10}(T_{\rm eff})$ is reduced to less than ~ 0.15 dex, inducing a maximum error of $\pm 23\%$ (± 0.09 dex) on L_{\star} . Based on Fig. 4, we have adopted a linear relationship between M_J and $T_{\rm eff}$: $\log(T_{\rm eff}/3700) = -0.055 \times (M_J - 4.0)$ (see the heavy dashed line in Fig. 4). This leads to the following $L_{\star} - M_J$ conversion:

$$\log_{10}(L_{\star}) = 1.49 - 0.466 \times M_J. \tag{3}$$

The uncertainty resulting from this conversion should be on the order of 0.045 dex for $0.01 \le L_{\star} \le 10.0 \,\mathrm{L}_{\odot}$. We therefore estimate that the total uncertainty on $\log_{10}(L_{\star})$ is $\sigma(\log_{10}L_{\star}) = (0.045^2 + (0.466 \times \sigma(M_J))^2)^{1/2} = 0.19 \,\mathrm{dex}$.

The $L_{\star}-M_{J}$ conversion described above cannot be applied to sources undetected in the J band. Instead, we




Fig. 4. M_J – $\log_{10}(T_{\rm eff})$ "HR" diagram showing the well-defined strip of the HR diagram occupied by the PMS tracks of D'Antona & Mazzitelli (1998). For $M_J=4.0$ for instance, we see that $\log_{10}(T_{\rm eff})$ can span a range of only ~ 0.15 dex for stellar ages between 0.1 Myr and 3 Myr. We thus adopt the M_J – $\log_{10}(T_{\rm eff})$ linear relationship displayed as a heavy dashed line which corresponds to the conversion: $\log_{10}(L_{\star})=1.49-0.466\times M_J$.

use the H-band magnitude, along with the extinction estimate derived from the (H-K) color, but with the additional complication that the circumstellar (disk) emission cannot be neglected. Meyer et al. (1997) found that the H-band circumstellar excess is on the order of 20 % of the stellar flux, on average, for the Taurus CTTS sample of Strom et al. (1989) (for a ρ Ophiuchi sample, see Greene & Lada 1996). This excess, expressed as a veiling index $r_H = F_{\nu \rm exc}^H/F_{\nu \star}^H$ (e.g. Greene & Meyer 1995), is equal to ~ 0.2 . Accordingly, we have applied a systematic correction $\delta H = -2.5 \log_{10}(1+r_H) \sim 0.2$ mag. The $L_{\star}-M_H$ relationship obtained in a way similar to the J-band relation is then:

$$\log_{10}(L_{\star}) = 1.26 - 0.477 \times (M_H + 0.2) \tag{4}$$

The total uncertainty on $\log_{10}(L_{\star})$ derived from M_H is then $\sigma(\log_{10} L_{\star}) = (0.045^2 + (0.477 \times \sigma(M_H))^2)^{1/2} = 0.29$ dex.

This method can also be applied to Class III YSOs, using different values for $(J-H)_0$ and $(H-K)_0$. We have derived L_{\star} for all Class III YSOs using the following relationships: $A_V = 9.09 \times [(J-H)-0.6]$, or $A_V = 15.4 \times [(H-K)-0.15]$; and $\log(L_{\star}) = 1.49-0.466 \times (J-0.265 \times A_V-5.73)$, or $\log(L_{\star}) = 1.26-0.477 \times (H-0.155 \times A_V-5.73)$

 $A_V - 5.73$) (the H-band IR excess for Class III YSOs is negligible).

The resulting M_J, M_H, A_V , and L_{\star} estimates are listed in Table 3 for Class II YSOs, and in Table 4 and Table 5 for Class III YSOs.

4.2. Disk luminosities for Class II YSOs

Since the SED of an embedded Class II YSO peaks in the mid-IR range, the ISOCAM fluxes should be approximately valid tracers of the total, bolometric luminosities $(L_{\rm bol})$. To estimate $L_{\rm bol}$ for weak Class II sources, an empirical approach thus consists in using this $F_{\nu}^{\text{MIR}}-L_{\text{bol}}$ relationship after proper calibration on a sub-sample of (brighter) objects for which the luminosity can be derived by a more direct method. This approach has been adopted by, e.g., Olofsson et al. (1999). Here, we have used the L_{\star} estimates of Sect. 4.1 to check that a correlation is actually present between L_{\star} and the mid-IR fluxes. Figure 5 displays $F_{\nu}^{14.3}$ (corrected for extinction) as a function of L_{\star} for the 104 Class II sources detected both in the near-IR and in the mid-IR range.




Fig. 5. Correlation between $F_{\nu}^{14.3}$ (corrected for interstellar extinction with $A_{14.3} = 0.03 \times A_V$) and L_{\star} . The observations are compared with a purely reprocessing disk model with $L_{\rm disk} = 0.25 \times L_{\star}$ and a stellar contribution given by Eq. 6. The continuous line corresponds to an average disk inclination ($i = 60^{\circ}$), while the dotted and dashed lines correspond to more extreme inclinations ($i = 0^{\circ}$ and $i = 85^{\circ}$).

A correlation is found, showing that, despite some scatter, the ISOCAM fluxes can be used to give rough estimates of the stellar luminosities of Class II YSOs. This is useful for the few ISOCAM sources of our sample which have not been detected at near-IR wavelengths.

The mid-IR emission of Class II YSOs is usually interpreted as arising from warm dust in an optically thick circumstellar disk. Using a simplified disk model (e.g. Beckwith et al. 1990), it is easy to show that any observed monochromatic flux in the optically thick, powerlaw range of the disk SED is simply proportional to the total disk luminosity L_{disk} divided by the projection factor $\cos(i)$, where i is the disk inclination angle to the line of sight. In the Beckwith et al. (1990) model, the disk is parameterized by a power-law temperature profile with three free parameters, T_0 , r_0 , and q, such that: $T(r) = T_0 \times (r/r_0)^{-q}$. Here, we have adopted $T_0 = 1500$ K, meaning that the disk inner radius is at the dust sublimation temperature, and q = 2/3, corresponding to an IR spectral index $\alpha_{\rm IR} = 2/q - 4 = -1.0$ typical of CTTS spectra.

We must, however, account for the fact that the stellar emission itself is not completely negligible in the mid-IR bands, especially at 6.7 μ m. A simple blackbody emission at $T_{\star} = 3700 \text{ K}$ gives

$$F_{\nu \star}^{6.7} = 0.082 \times (L_{\star}/1 \,\mathrm{L}_{\odot}) \,\mathrm{Jy}$$
 (5)
 $F_{\nu \star}^{14.3} = 0.021 \times (L_{\star}/1 \,\mathrm{L}_{\odot}) \,\mathrm{Jy}.$ (6)

$$F_{\nu \star}^{14.3} = 0.021 \times (L_{\star}/1 \, \text{L}_{\odot}) \, \text{Jy}.$$
 (6)

Since $F_{\nu \star}^{14.3}$ is significantly smaller than $F_{\nu \star}^{6.7}$, only $F_{\nu}^{14.3}$ is used to estimate L_{disk} . Assuming $\cos(i) = 0.5 \ (i = 60^{\circ})$ in all cases, we have computed $L_{\rm disk}$ for each of the Class II YSOs with L_{\star} and A_{V} estimates from Sect. 4.1, based on the following relationship:

$$L_{\text{disk}}/1 \, L_{\odot} = (F_{\nu}^{14.3} \times 10^{+0.4 \times A_{14.3}} - F_{\nu, +}^{14.3})/1.80 \,\text{Jy}, \quad (7)$$

where $A_{14.3} = 0.03 \times A_V$ and $F_{\nu \star}^{14.3}$ from Eq. 6. The results are given in the last column of Table 3.

According to this model, the mid-IR flux is a direct tracer of the disk luminosity, and the $F_{\nu}^{\text{MIR}} - L_{\star}$ correlation of Fig. 5 simply expresses that $L_{\rm disk}$ correlates with L_{\star} . The origin of $L_{\rm disk}$ is either the release of gravitational energy by accretion in the disk, or the absorption/reprocessing of stellar photons by the dusty disk. In the latter case, $L_{\rm disk}$ is naturally proportional to L_{\star} . The fraction $L_{\rm disk}/L_{\star}$ of stellar luminosity reprocessed by the disk depends on the spatial distribution of dust. In the ideal case of an infinite, spatially flat disk, this fraction is 0.25 (Adams & Shu 1986). If the disk is flared, $L_{\rm disk}/L_{\star}$ is larger, while it is smaller if the disk has a inner hole. The theoretical $F_{\nu}^{\text{MIR}} - L_{\star}$ correlations plotted in Fig. 5 correspond to $L_{\rm disk} = 0.25 \times L_{\star}$ and to three representative inclination angles. The fact that this simple model accounts for the observed correlation quite well, suggests that the disks of most ρ Ophiuchi Class II YSOs are passive disks dominated by reprocessing. This is consistent with recent estimates of the disk accretion level in Taurus CTTSs (e.g. Gullbring et al. 1998). The typical disk accretion rate of a CTTS is estimated to be $10^{-8} \,\mathrm{M_{\odot}/yr}$, corresponding to an accretion luminosity $L_{\rm acc} = 0.025 \, \rm L_{\odot}$ for $R_{\star} = 3 \, \mathrm{R}_{\odot}$ and $M_{\star} = 0.25 \, \mathrm{M}_{\odot}$ (i.e., $L_{\star} \sim 0.25 \, \mathrm{L}_{\odot}$ at 1 Myr). In this case, the luminosity due to reprocessing is ~ 5 times larger than the accretion luminosity in the disk.

On the other hand, 37 sources (among a total of 104) are located above the passive disk model lines in Fig. 5. These are good candidates for having an active disk with an accretion rate typically larger than $\sim 5 \times (L_{\star}/0.25 L_{\odot}) \times 10^{-8} M_{\odot}/yr$.

Overall, we find that the median $L_{\rm disk}/L_{\star}$ ratio is 0.41 for the 93 Class II sources detected in the near-IR and with $F_{\nu}^{14.3} > 15 \,\mathrm{mJy}$ (i.e. the completeness level derived in Sect. 2.4). Using this ratio, we have derived rough estimates of the stellar luminosities of the 15 Class II YSOs which have no near-IR photometry (see Table 3) as follows:

$$L_{\star}(L_{\odot}) \approx L_{\rm disk}/0.41 \approx (0.97 \times 1.6/0.41) \times (F_{\nu}^{14.3}/1.8 \, \rm Jy).(8)$$

[The factor 0.97 corresponds to a typical stellar contribution of 3% at 14.3 μ m, while the factor 1.6 is an average extinction correction at 14.3 μ m (corresponding to $< A_V >= 17$ mag).]

4.3. Calorimetric luminosities for Class I sources

The most direct method of estimating the total luminosities $L_{\rm bol}$ of embedded YSOs consists in integrating the observed SEDs (cf. WLY89). However, since most of the ρ Ophiuchi Class II and Class III YSOs are deeply embedded within the cloud $(A_V \gtrsim 10)$, only a negligible fraction of their bolometric luminosity can be recovered by finitebeam IR observations (e.g. Comerón et al. 1993). We thus do not attempt to derive calorimetric estimates of $L_{\rm bol}$ for these sources. In contrast, the calorimetric method is believed to be appropriate for Class I YSOs since these are self-embedded in substantial amounts of circumstellar material which re-radiate locally the absorbed luminosity (cf. WLY89 and AM94). Using our new mid-IR measurements, we have evaluated the calorimetric luminosities (L_{cal}) of the 16 Class I YSOs observed in our survey. Only 7 of them have reliable IRAS fluxes up to 60 or 100 μ m (IRS54, IRS44, GSS30, IRS43, EL29, IRS48, IRS51). For these, the median of the ratio of $L_{\rm cal}(6.7-14.3\mu{\rm m})$ to $L_{\rm bol}$ is found to be 9.8, suggesting that the typical fraction of a Class I source's luminosity radiated between 6.7 and 14.3 μm is $\sim 10 \%$. Assuming that this ratio is representative of all Class I YSOs, we have derived estimates of L_{bol} for the remaining 9 weaker Class I sources (i.e., CRBR85, LFAM26, LFAM1, WL12, IRS46, CRBR12, IRS67, CRBR42, WL6). These luminosities are listed in Table 2.

4.4. Luminosity function of the ρ Ophiuchi embedded cluster

Combining the L_{\star} luminosities determined in Sect. 4.1 for the sources detected in the near-IR with the L_{\star} estimates from $L_{\rm disk}$ for the sources without near-IR measurements (Sect. 4.2), we have built a luminosity function for Class II YSOs which represents a major improvement over previous studies (see Fig. 6a). In terms of L_{\star} , the completeness level for this population can be estimated from the $F_{\nu}^{14.3}$ completeness limit derived in Sect. 2.4 ($F_{\nu}^{14.3} \sim 15\,{\rm mJy}$) using Eq. 8: $L_{\star}^{comp}({\rm ClassII}) = 0.032\,{\rm L}_{\odot}$. While the luminosity function previously published by Greene et al.

(1994) included only 33 (bright) Class II sources and suffered from severe incompleteness below $L_{\star} \sim 1-2\,\mathrm{L}_{\odot}$, our present completeness level is a factor $\sim 30\text{--}50$ lower. The new luminosity function shows a marked flattening in logarithmic units at $L_{\star} \sim 2\,\mathrm{L}_{\odot}$, well above our completeness limit. This important new feature is discussed in Sect. 5 below.

Based on the $L_{\rm cal}$ estimates of Sect. 4.3, a new bolometric luminosity function for the 16 Class I YSOs of ρ Ophiuchi is displayed in Fig. 6b. The associated completeness level is derived from $F_{\nu}^{6.7}=10\,\mathrm{mJy}$ and $F_{\nu}^{14.3}=15\,\mathrm{mJy}$ (Sect. 2.4) using $L_{\rm bol}/L_{\rm cal}(6.7$ –14.3 μ m) \approx 9.8 (Sect. 4.3): $L_{\star}^{\rm comp}({\rm Class I})=0.017\,\mathrm{L}_{\odot}$. The median $L_{\rm bol}$ for Class I YSOs is 1.6 L $_{\odot}$, which is \sim 8 times larger than the median L_{\star} of Class II YSOs (0.20 L $_{\odot}$). The luminosities of Class I YSOs span a range of two orders of magnitude between $\sim 0.1\,\mathrm{L}_{\odot}$ and $\sim 10\,\mathrm{L}_{\odot}$, which is roughly as wide as the luminosity range spanned by Class II YSOs. The comparatively large value of $L_{\rm bol}$ for Class I YSOs is probably due to a dominant contribution of accretion luminosity as expected in the case of protostars.

In Fig. 6c, we plot the luminosity function of the 55 Class III sources that are located within the CS contours of Fig. 1 and for which we have enough near-IR data to derive L_{\star} according to the procedure described in Sect. 4.1. This sample comprises 19 confirmed Class IIIs from Table 4 together with 36 candidate Class III sources from Table 5. It might be contaminated by a few background/foreground sources but is characterized by a relatively well defined completeness luminosity. From the completeness level $F_{\nu}^{6.7} = 10 \,\mathrm{mJy}$ and using Eq. 5 with an average extinction correction corresponding to $A_V =$ 17 mag, we get $L_{\star}^{comp}(\text{ClassIII}) = 0.20 \, \text{L}_{\odot}$, which is ~ 6 times higher than L_{\star}^{comp} (ClassII). Deep X-ray observations with XMM should improve the completeness luminosity for Class III YSOs by an order of magnitude in the near future (cf. discussion by Grosso et al. 2000 in a companion paper).

5. Modeling of the luminosity function

If the mass-luminosity relationship of PMS stars were a simple, fixed power-law function, the shape of the luminosity function would directly reflect the underlying mass function. Unfortunately, the mass-luminosity relation is a complex function with inflexion points and is strongly age-dependent. Deriving the mass function from the luminosity function thus requires knowledge of the stellar age distribution. The ρ Ophiuchi embedded cluster is believed to be younger than most known star clusters (e.g. Luhman & Rieke 1999), with a typical age on the order of, or less than, 1 Myr (e.g. WLY89, Greene & Meyer 1995).

The 123 Class II sources identified with ISOCAM are presently the most complete sample of young stars available in the cluster. The corresponding mass function is estimated down to $0.055\,\rm M_\odot$ in Sect. 5.1. We investigate the effect of including Class III YSOs by modeling the lu-




Fig. 6. Luminosity functions (LF) (a) for the 123 Class II YSOs (continuous histogram with statistical error bars). The function corresponding to a similar histogram shifted by half the 0.2 dex bin size is shown as a thin curve to illustrate the level of statistical fluctuations due to binning. The LF of 33 Class II sources from Greene et al. (1994) is displayed as a darker histogram (rebinned to 0.2 dex bins, and rescaled to d = 140 pc for better comparison with the new LF). The typical $\pm 1\sigma$ uncertainty on the position of each bin resulting from the individual uncertainties on L_{\star} is indicated, along with the effect of the distance uncertainty; (b) for the 16 Class I YSOs; and (c) for the sample of 55 (19 confirmed + 36 candidate) Class III sources from Table 4 and 5. The LF of (a) is shown in the background for comparison. The dashed vertical lines show the respective completeness levels.

minosity function of the 135 Class II and Class III objects located inside the CS contours of Fig. 1 (Sect. 5.3).

5.1. Constraints on the mass function of Class II YSOs

We have modeled the observed luminosity function using a two-segment power-law mass function with two free parameters, $M_{\rm flat}$ and α_1 : ${\rm d}N/{\rm d}\log M_{\star} \propto {M_{\star}}^{\alpha_1}$ for $M_{\star} < M_{\rm flat}$, and $dN/d \log M_{\star} \propto {M_{\star}}^{-1.7}$ for $M_{\star} > M_{\rm flat}$. The high-mass range (for $M_{\star} > M_{\rm flat}$) is simply taken to be a power-law with index -1.7 (whereas the Salpeter index would be -1.35), in agreement with the IMFs favored by Kroupa et al. (1993) and Scalo (1998). We assume that the cluster forms stars with a fixed IMF, independent of time. The evolutionary tracks of D'Antona & Mazzitelli (1998)⁴ are used to derive a set of mass-luminosity relations adapted to the ρ Ophiuchi PMS population, under the following three hypotheses about the star formation history: Constant star formation rate over a) 0.5 Myr (Fig. 7a), b) 1 Myr (Fig. 7b), and c) 2 Myr (Fig. 7c). Cases b) and c) correspond to the simplest scenarios consistent with current observational constraints (e.g. WLY89). Case a) mimics a recent "burst" of star formation suggested by

recent studies (e.g. Greene & Meyer 1995 and Luhman & Rieke 1999). These three scenarios are roughly representative of our current (imperfect) knowledge of the stellar age distribution in the cluster.

The fitting analysis is performed on the 12 (logarithmic) luminosity bins above the completeness level of $0.032\,\mathrm{L}_\odot$ and below $10\,\mathrm{L}_\odot$ by varying α_1 and M_flat . Due to the age ambiguity for a particular (M_{\star}, L_{\star}) in the PMS tracks for $M_{\star} \gtrsim 2 \,\mathrm{M}_{\odot}$ and $age < 3 \,\mathrm{Myr}$ as a result of the transition from convective to radiative interiors (e.g. Fig. 4), our simplified modeling is not valid for $L_{\star} \gtrsim 10 \, \mathrm{L}_{\odot}$. The best fit is shown as a heavy curve for each of the three assumed age distributions (Fig. 7a, 7b, and 7c). The flattening of the luminosity function below $2 L_{\odot}$ is well reproduced in all three cases. The best-fit model in the 1 Myr case (Fig. 7b) is however not quite as good as the two others since it predicts a peak at $L_{\star} \sim 0.4 - 0.5 \, \mathrm{L}_{\odot}$ where the data show a dip. This is reflected by the largest value 1.41 for the reduced $\chi^2_{\nu} \equiv \frac{\chi^2}{N-n}$ (with N=12 fitted data points and n=2 free parameters). The model of Fig. 7c reproduces the data somewhat better ($\chi^2_{\nu} = 1.10$) as the predicted peak luminosity moves down to $L_{\star} \sim 0.2 - 0.3 \, \mathrm{L}_{\odot}$. A marginally better fit is obtained with the 0.5 Myr model (Fig. 7a, $\chi^2_{\nu} = 0.66$). In this case, the model luminosity function has a local maximum at $L_{\star} \gtrsim 1 \, \mathrm{L}_{\odot}$ as observed and the flattening for $L_{\star} \sim 2 \, \mathrm{L}_{\odot}$ is particulary well reproduced.

⁴ We here adopt the tracks of D'Antona & Mazzitelli (1998) as they provide the most extensive set of PMS models available to date, including a tight grid of masses and ages. In the future, we plan to check the details of our results using other tracks such as those of Baraffe et al. (1998) or Palla & Stahler (1999).




Fig. 7. Best model fits (heavy curves) to the luminosity function of Class II sources obtained for a stellar age distribution corresponding to a constant formation rate over (a) 0.5 Myr, (b) 1 Myr, and (c) 2 Myr (the first 0.2 Myr have been avoided in the PMS tracks to account for a typical protostellar timescale). In (d), similar best fits are shown for the combined luminosity function of 135 Class II and Class III sources inside the CS contours of Fig. 1. The vertical dashed lines show the completeness levels of the respective luminosity functions (see Fig. 6).

The best-fit values of $M_{\rm flat}$ along with the formal $1\,\sigma$ errors resulting from the fitting procedure are as follows: $0.33\pm0.04\,{\rm M}_{\odot},\ 0.55\pm0.09\,{\rm M}_{\odot},\ {\rm and}\ 0.74\pm0.13\,{\rm M}_{\odot}$ for Fig. 7a, 7b, and 7c, respectively. These $M_{\rm flat}$ values follow the expected trend that the older the stars, the higher the derived masses for the same luminosities. The final error on the determination of $M_{\rm flat}$ is clearly dominated by the uncertainties on the stellar age distribution. We adopt the following average value: $M_{\rm flat}=0.55\pm0.25\,{\rm M}_{\odot}.$ The best-fit values for α_1 are $-0.02\pm0.15,\ -0.27\pm0.13,\ {\rm and}\ -0.19\pm0.13,\ {\rm respectively}.$ There is no correlation with age in this case. We conservatively adopt $\alpha_1=-0.15\pm0.2.$ These constraints on the mass function are valid for $0.032\,{\rm L}_{\odot} < L_{\star} < 10\,{\rm L}_{\odot},\ {\rm which}\ {\rm approximately}\ {\rm corresponds}\ {\rm to}\ 0.055\,{\rm M}_{\odot} \lesssim M_{\star} \lesssim 2\,{\rm M}_{\odot}.$

Under the same three assumptions about the stellar age distribution as above, we have also derived a range of stellar masses directly from the M_J magnitude for each of the 123 Class II YSOs. We display the results as a mass function in Fig. 8, where the vertical error bars reflect the uncertainties induced by the various age assumptions rather than the statistical errors. The best two-component power-law mass function derived above is superposed as a heavy curve. The error bars are particularly large for the two mass bins close to $1 \, \mathrm{M}_{\odot}$, i.e., just above M_{flat} . This is due to a shift of a large number of stars beyond M_{flat} when the stellar ages are varied upward. The global shape of the mass function is however fairly well determined: It

is basically flat (in logarithmic units) at low masses down to $0.055\,\rm M_\odot$ and shows a steep decline beyond $\sim 0.5\,\rm M_\odot$.




Fig. 8. Mass function of the 123 Class II YSOs. The range of possible age distributions (see Sect. 5.1) induces a range of masses for each star and thus an uncertainty on the derived histogram displayed as vertical error bars. The best two-segment power-law mass function of Sect. 5.1 (heavy solid curve), and the effect of binarity as a function of the binary fraction from 50 % to 100 % (light curves – see Sect. 6.2) are shown. The vertical dashed line marks the completeness level of $0.055\,\mathrm{M}_{\odot}$.

5.2. Possible deuterium feature in the luminosity function of Class II YSOs

Some of the features (peaks and dips) apparent in the observed luminosity function might be real as they are also predicted by the models (see Fig. 7). The feature expected in the luminosity function of young stars as a result of deuterium burning during PMS contraction is particularly interesting since its location is age-dependent (e.g. Zinnecker et al. 1993). As already noted, the model fit obtained under the 0.5 Myr scenario is particularly good as it predicts a peak at $L_{\star} \sim 1.5 \, \mathrm{L}_{\odot}$, consistent with the observations (see Fig. 7a). In the model, this $1.5 \, \mathrm{L}_{\odot}$ peak corresponds to an inflexion point in the mass-luminosity relation due to deuterium burning in ~ 0.4 -Myr old stars with $M_{\star} \sim 0.5 \, \mathrm{M}_{\odot}$. If this peak in the luminosity function is confirmed, it will provide strong evidence that the population of Class II YSOs is particularly young in ρ Ophiuchi.

5.3. Effect of Class III YSOs

There is most probably a significant population of Class III objects embedded in the cloud (e.g. Sect. 3.5). To evaluate how the presence of this population may affect our conclusions on the IMF of the cluster (see Sect. 6 below), we here consider the luminosity function of a combined sample of Class II and Class III sources. The sample comprises the 80 Class II and 55 Class III YSOs located inside the CS contours of Fig. 1. It should be complete down to $\sim 0.2\,L_\odot$. (Note, however, that only 19 of the 55 Class III sources of this sample are confirmed YSOs at this stage – see Sect. 4.4.) The associated luminosity function, displayed in Fig. 7d, is very similar to the luminosity function of Class II YSOs. In particular, it exhibits a flattening below $\sim 1-2\,L_\odot$.

The results of a modeling similar to that of Sect. 5.1, but performed on only 8 luminosity bins between 0.2 and $10\,\mathrm{L}_\odot$, are shown in Fig. 7d. The best fits found under the same three star formation scenarios as in Sect. 5.1 are good (with $\chi^2_\nu = 0.64$, 0.94, and 1.15, respectively). The best-fit values for the parameters M_{flat} and α_1 of the mass function are as follows: $M_{\mathrm{flat}} = 0.32 \pm 0.04\,\mathrm{M}_\odot$, $0.48 \pm 0.08\,\mathrm{M}_\odot$, $0.66 \pm 0.10\,\mathrm{M}_\odot$, and $\alpha_1 = -0.08 \pm 0.45$, 0.00 ± 0.35 , $+0.05 \pm 0.19$ for star formation durations of 0.5, 1, and 2 Myr, respectively. Combining these results, we obtain $M_{\mathrm{flat}} \sim 0.5 \pm 0.2\,\mathrm{M}_\odot$ and $\alpha_1 \sim 0.0 \pm 0.4$. Within the uncertainties these values are essentially identical to those found in Sect. 5.1 for the sample of Class II YSOs, although they apply to a more limited mass range, approximately $0.17\,\mathrm{M}_\odot \lesssim M_\star \lesssim 2\,\mathrm{M}_\odot$.

In contrast to the Class II case, the luminosity function of the Class II—Class III combined sample does not show a peak at $L_{\star} = 1.5 \, \mathrm{L}_{\odot}$ (cf. Fig. 7d). This may be understood if the average stellar age of the combined population is somewhat larger than that of the population of Class II sources (as is expected).

6. Discussion

6.1. The initial mass function in ρ Ophiuchi

Since the ρ Ophiuchi molecular cloud is still actively forming stars, the final IMF of the cluster cannot be directly measured. The masses of the stars already formed in the cluster provide only a "snapshot" of the local IMF (cf. Meyer et al. 2000). However, assuming that the mass distribution of formed stars does not change significantly with time during the cluster's history (an assumption made in the models of Sect. 5.1), any snapshot of the mass distribution taken on a large, complete population of PMS objects should accurately reflect the end-product IMF

Compared to previous investigations of the IMF in the ρ Ophiuchi cloud based exclusively on near-IR data (e.g. Comerón et al. 1993, Strom et al. 1995, Williams et al. 1995, Luhman & Rieke 1999), the use of mid-IR photometry has allowed us to consider a much larger sample of young stars (see, e.g., Fig. 6). Since our sub-sample of Class II YSOs is fairly large (123 objects) and complete down to low luminosities, it provides an excellent opportunity to derive improved constraints on the ρ Ophiuchi IMF down to low masses.

In a statistical sense at least, Class II YSOs are believed to represent a specific phase of PMS evolution which follows the (Class 0 and Class I) protostellar phases, and precedes the Class III phase (see Sect. 1). Due to their short lifetime ($\lesssim 10^5\,\mathrm{yr}$), protostars make up only a small fraction of a young cluster's population (cf. Fletcher & Stahler 1994), and can be neglected in the global mass function. Furthermore, in contrast to PMS stars, protostars have not yet reached their final stellar masses.

Class III objects are more numerous and should thus contribute significantly to the global mass distribution. Furthermore, it has been suggested that some YSOs evolve quickly to the Class III stage, perhaps as early as the "birthline" for PMS stars (e.g. Stahler & Walter 1993), and spend virtually no time in the Class II phase (cf. André et al. 1992, Greene & Meyer 1995). Since such objects cannot be identified through IR observations, their exact number and mass distribution will not be known until the results of deep X-ray (and follow-up) surveys are available (see Sect. 3.5). However, providing the (range of) evolutionary timescale(s) from Class II to Class III is independent of mass, both classes of PMS objects should have identical mass functions. The results of Sect. 5.3 do seem to support this view, as they suggest that the mass functions of the Class II and Class III samples do not differ down to $M_{\star} \sim 0.17 \,\mathrm{M}_{\odot}$.

We therefore conclude (and will assume in the following) that the mass distribution of Class II YSOs determined in Sect. 5.1 and shown in Fig. 8 currently represents our best estimate of the IMF in the ρ Ophiuchi embedded cluster down to $M_{\star}\sim 0.055\,{\rm M}_{\odot}.$ As discussed in Sect. 6.2 below, this mass function applies to stellar systems rather than single stars.

6.2. Influence of binary stars

A large proportion ($\gtrsim 50\%$) of field stars are in fact multiple (e.g. binary) systems (e.g. Duquennoy & Mayor 1991). This is also true for young PMS stars, and there is a growing body of evidence that the binary fraction is even larger in a young PMS cluster like ρ Ophiuchi than for main sequence stars in the field (e.g. Leinert et al. 1993; Simon et al. 1995). Since the present study is based on ISOCAM/near-IR observations which do not have enough angular resolution to separate most of the expected binaries, a significant population of low-mass stellar companions are presumably missing from the mass function derived above. These low-mass companions are hidden by the corresponding primaries.

To estimate the magnitude of this binary effect, we show three simple models in Fig. 8 which assume a population of hidden secondaries corresponding to a binary fraction f of 50%, 75%, and 100 %, respectively. In each case, we start from a primary mass function with the two-segment power law form derived in Sect. 5.1. We then add a population of secondaries with masses distributed uniformly (in logarithmic units) between a minimum mass of $0.02\,\mathrm{M}_\odot$ and the mass of the primary. We thus assume that the component masses are uncorrelated and drawn from essentially the same mass function (cf. Kroupa et al. 1993).

The global mass functions resulting from addition of companions to the primary mass function are shown in Fig. 8. It can be seen that the global mass functions are similar in form to the original primary mass function. Neither the position of the break point $(M_{\rm flat})$ nor the slope in the high-mass range are affected by the addition of companions. However, the slope in the low-mass range (α_1) steepens as the binary fraction increases. Indeed, the power-law index between $0.055\,{\rm M}_{\odot}$ and $M_{\rm flat}=0.55\,{\rm M}_{\odot}$ changes from $\alpha_1=-0.15$ to $\alpha_1=-0.31$ for f=50%, $\alpha_1=-0.37$ for f=75%, and $\alpha_1=-0.42$ for f=100%.

In summary, if we account for uncertainties in the binary fraction (50% $\lesssim f \lesssim 100\%$), our best estimate of the single-star mass function in ρ Ophiuchi is well described by a two-segment power-law with a low-mass index $\alpha = -0.35 \pm 0.25$ down to $0.055\,\mathrm{M}_\odot$, a high-mass index $\alpha_2 = -1.7$, and a break (flattening) occurring at $M_\mathrm{flat} \sim 0.55 \pm 0.25\,\mathrm{M}_\odot$.

6.3. Luminosity/mass functions in individual cloud cores

Our sample of Class II YSOs is large enough that we can study the properties of the four sub-clusters associated with the dense cores Oph A, Oph B, Oph EF, and L1689S (see Fig. 1). All 123 Class II sources but 5 belong to these 4 sub-clusters. The luminosity functions of Class II objects in the individual sub-clusters are displayed in Fig. 9 along with the 'best-fit' model of Sect. 5.1 (solid curve). They all agree reasonably well in shape with both the total luminosity function and the model: all four luminosity




Fig. 9. Individual luminosity functions for the four subclusters associated with the dense cores Oph A, Oph B, Oph EF, and L1689S (see Fig. 1). These are displayed with 0.4 dex bins and scaled by a factor 1/2 for direct comparison with the 0.2 dex bin total luminosity function (light grey histogram in the background). The light solid curves correspond to the model of Fig. 7a scaled to the number of stars in each sub-cluster.

functions are essentially flat over two orders of magnitude in luminosity and appear to have a peak at $L_{\star} \sim 1.5 \, \rm L_{\odot}$. The agreement is particularly good for the sub-cluster with the largest number of stars, Oph A (see Fig. 9a), but even the smallest sub-cluster, L1689S, tends to reproduce the shape of the global luminosity function on a smaller scale (Fig. 9d). In contrast, for instance, the luminosity function derived for the Chamaeleon I cloud based on ISOCAM data (Persi et al. 2000) differs markedly from the ρ Ophiuchi luminosity functions. It does not exhibit any peak at $1.5 \, \rm L_{\odot}$ and is consistent with an older ($\sim 3 \, \rm Myr)$ PMS population having a similar underlying mass function (see Kaas & Bontemps 2001).

The similarity of the individual luminosity functions suggests similar distributions of stellar ages and stellar masses in each of the four sub-clusters.

6.4. Global aspects of star formation in ρ Ophiuchi

After correction for unresolved binaries (assuming a binary fraction f=75%), the total number of Class II sources (including companions) down to $0.055\,\mathrm{M}_\odot$ is ~ 145 . Assuming a Class III to Class II number ratio of 19/22 as found by X-ray surveys (Grosso et al. 2000 – see Sect. 3.5), we infer the presence of ~ 125 Class III stars (including associated companions) in the same mass range. The typical number ratio of Class Is (plus Class 0s) to Class IIs is 18/123 suggesting an additional ~ 21 embedded YSOs. Altogether, we therefore estimate that there are currently $N_{\star} \sim 291$ YSOs down to $\sim 0.055\,\mathrm{M}_\odot$ including $\sim 19\,\%$ of brown dwarfs. The average and median masses of these objects are $\sim 0.35\,\mathrm{M}_\odot$ and $\sim 0.20\,\mathrm{M}_\odot$ respectively. The total mass of condensed objects (includ-

ing brown dwarfs) in the cluster is thus estimated to be $M_{\star}^{\rm clust} = 291 \times 0.35 \sim 102 \, {\rm M}_{\odot}$. (The brown dwarfs with $0.055 \, {\rm M}_{\odot} < M_{\star} < 0.08 \, {\rm M}_{\odot}$ contribute only $\sim 4 \, \%$ of this mass.) Restricting ourselves to L1688 (thus subtracting the $\sim 10 \, \%$ contribution from L1689) whose average radius is approximately 0.4 pc (cf. CS contours in Fig. 1), we find $N_{\star}^{\rm L1688} \sim 262, \, M_{\star}^{\rm L1688} \sim 92 \, {\rm M}_{\odot}, \, n_{\star}^{\rm L1688} \sim 980 \, {\rm stars/pc^3},$ and $\rho_{\star}^{\rm L1688} \sim 340 \, {\rm M}_{\odot}/{\rm pc^3},$ where n_{\star} and ρ_{\star} are the stellar number density and stellar mass (volume) density of the cluster, respectively.

Adopting a conservative value of 2 Myr for the cluster age, the total mass of $M_{\star}^{\rm clust} = 102 \, {\rm M}_{\odot}$ translates into an average star formation rate of $5.1 \times 10^{-5} \, {\rm M}_{\odot}/{\rm yr}$, corresponding to one new YSO (of typical mass $0.20 \, {\rm M}_{\odot}$) every $\sim 4000 \, {\rm yr}$.

Lastly, we can derive the star formation efficiency (SFE) in L1688, defined as SFE= $M_{\rm star}/(M_{\rm star}+M_{\rm gas})$. The total molecular gas mass, $M_{\rm gas}$, of L1688 has been estimated to range between 550 M $_{\odot}$ (from C¹8O measurements – Wilking & Lada 1983) and 1500 M $_{\odot}$ (from CS(2-1) data – Liseau et al. 1995). Using $M_{\star}^{\rm L1688}=92\,{\rm M}_{\odot}$, we thus get SFE^{L1688} $\sim 6-14$ %, which is somewhat lower than previous estimates ($\geq 22\%$ – WLY89). Note, however, that active star formation in L1688 appears to be limited to the three sub-clusters/dense cores Oph A, Oph EF, and Oph B (see Fig. 1 and Loren et al. 1990), where the local star formation efficiency is significantly higher: $SFE \sim 31\%$, using a total core mass of $200\,{\rm M}_{\odot}$ (Loren et al. 1990).




Fig. 10. Comparison of the pre-stellar mass spectrum measured by MAN98 for 58 protocluster condensations (dark histogram with statistical error bars) with the YSO mass function derived in Sect. 5.1 for 123 Class II systems (light histogram with error bars accounting for both statistical and age uncertainties).

6.5. Comparison with the ρ Ophiuchi protocluster condensations

In an extensive 1.3 mm dust continuum imaging survey of L1688 with the IRAM 30 m telescope (11" resolution), MAN98 could identify 58 compact starless condensations. Molecular line observations (e.g. Belloche, André, & Motte 2001) indicate that the condensations are gravitationally bound and thus likely pre-stellar in nature. MAN98 noted a remarkable similarity between the mass spectrum of these pre-stellar condensations and the IMF of Miller & Scalo (1979).

In Fig. 10, we compare the pre-stellar mass spectrum determined by MAN98 with the mass distribution of Class II YSOs derived in Sect. 5.1. (As such, both distributions are uncorrected for the presence of close binary systems.) It can be seen that there is a good agreement in shape between the two mass spectra. A Kolmogorov-Smirnov test performed on the corresponding cumulative distributions confirms that they are statistically indistinguishable at the 95% confidence level. This supports the suggestion of MAN98 that the IMF of embedded clusters is primarily determined by cloud fragmentation at the prestellar stage of star formation. The fact that both the prestellar and the YSO spectrum of Fig. 10 present a break at roughly the same mass $\sim 0.5\,\mathrm{M}_\odot$ is quite remarkable. A small, global shift of the masses by only $\sim 30\%$ upward or downward in one of the spectra would make them differ at the 2σ statistical level. Although in absolute terms, both sets of masses are probably uncertain by a factor of ~ 2 (due to uncertainties in the 1.3 mm dust opacity and in the cluster age, respectively), this suggests that the protocluster condensations identified at 1.3 mm may form stars/systems with an efficiency larger than $\sim 50-70\%$.

7. Conclusions

We have used ISOCAM to survey the ρ Ophiuchi main cloud for embedded YSOs down to a completeness level of $\sim 10-15\,\mathrm{mJy}$ at $6.7\,\mu\mathrm{m}$ and $14.3\,\mu\mathrm{m}$. Our main findings are as follows:

- 1. A total of 425 point sources are detected in the \sim 0.7 deg² field covered by the survey, of which 211 are seen at both 6.7 μ m and 14.3 μ m. The observed distribution of the flux density ratio $F_{\nu}^{14.3}/F_{\nu}^{6.7}$ is clearly bimodal with a gap separating two distinct groups of sources: a "red" group corresponding to Class I and Class II YSOs with optically thick mid-IR excesses, and a "blue" group consisting of Class III YSOs and background stars (cf. Fig. 3).
- 2. The red group consists of 139 cluster members, 71 of them being newly identified YSOs. Based on their mid-IR colors from 2 μ m to 6.7 μ m and 14.3 μ m, essentially all of these new YSOs are low-luminosity Class II objects. This brings the total number of Class II members to 123, a factor of 2 larger than previously known. Only 50 % of these Class II sources have a near-IR excess

- large enough to be recognizable in a near-IR color-color diagram.
- 3. Combining near-IR data from Barsony et al. (1997) with our mid-IR photometry, we derive stellar luminosities for 123 Class II and 74 Class III YSOs. We also estimate bolometric luminosities for 16 Class I objects. The corresponding luminosity functions (Fig. 6) are complete down to $\sim 0.02\,L_{\odot}, \sim 0.03\,L_{\odot},$ and $\sim 0.2\,L_{\odot}$ for Class I, Class II, and Class III objects, respectively. The luminosity function of Class II YSOs is essentially flat (in logarithmic units) below $\sim 2\,L_{\odot}$, with a possible peak at $1.5\,L_{\odot}$ and dip at $0.5\,L_{\odot}$ seen in the four sub-regions Oph A, Oph B, Oph EF, and L1689S (Fig. 9).
- 4. The large proportion of Class II objects observed above the completeness level for Class III sources $(M_{\star} \sim 0.2\,\mathrm{M}_{\odot})$ suggests that more than 50% of the ρ Ophiuchi embedded YSO population have optically thick circumstellar disks at mid-IR wavelengths. In a majority of cases, the luminosities of these disks are consistent with pure reprocessing of stellar light. Only $\sim 35\%$ of the Class II objects have excess mid-IR luminosities suggestive of substantial disk accretion rates.
- 5. The luminosity function of Class II objects is well modeled by a population of PMS stars with ages in the range $\sim 0.3-2$ Myr and a roughly flat mass distribution below $M_{\rm flat} \sim 0.5\,{\rm M}_{\odot}$, i.e., ${\rm d}N/{\rm d}\log M_{\star} \propto M_{\star}^{-0.15}$ down to $\sim 0.06\,{\rm M}_{\odot}$.
- 6. Pending the results of a deep X-ray census of Class III objects, we argue that the mass distribution of Class II YSOs is representative of the emergent IMF of the embedded cluster. If we account for the presence of unresolved binaries, this emergent mass function is well described by a two-segment power law with a low-mass index $\alpha_1 = -0.35 \pm 0.25$, a high-mass index $\alpha_2 = -1.7$, and a transition mass $M_{\rm flat} = 0.55 \pm 0.25 \,{\rm M}_{\odot}$ (where ${\rm d}N/{\rm d}\log M_{\star} \propto {M_{\star}}^{\alpha}$). We find no evidence for a sharp turnover at low masses down to at least $\sim 0.06 \,{\rm M}_{\odot}$.
- 7. The shape of the mass function for Class II systems is statistically indistinguishable from the mass spectrum determined at 1.3 mm by Motte, André, & Neri (1998) for the pre-stellar condensations of the protocluster. This supports the conclusion of these authors that the IMF may be primarily determined by fragmentation at the pre-stellar stage of star formation. It also suggests that the 1.3 mm protocluster condensations should form stars/systems with an efficiency larger than ~ 50–70%.

Acknowledgements. S.B. was supported by an ESA Research Fellowship during his stay at the Stockholm Observatory. The authors thank the referee, Andrea Moneti, for constructive criticisms.

References

Abergel, A., Bernard, J.P., Boulanger, F., et al., 1996, A&A 315, L329

Adams, F.C., Shu, F.H., 1986, ApJ 308, 836

Adams, F.C., Lada, C.J., Shu, F.H., 1987, ApJ 312, 788 André, P., Montmerle, T., 1994, ApJ 420, 837 (AM94)

André, P., Montmerle, T., Feigelson, E.D., 1987, AJ 93, 1182
 André, P., Montmerle, T., Feigelson, E.D., Stine, P.C., Klein, K.L., 1988, ApJ 335, 940

André, P., Martín-Pintado, J., Despois, D., Montmerle, T., 1990, A&A 236, 180

André, P., Deeney, B.D., Phillips, R.B., Lestrade, J.-F., 1992, ApJ 401, 667

André, P., Ward-Thompson, D., Barsony, M., 1993, ApJ 406, 122

André, P., Ward-Thompson, D., Barsony, M., 2000. In: Protostars and Planets IV, Mannings V., Boss A.P., Russell S.S. (eds.), University of Arizona Press, Tucson, p. 59

Baraffe, I., Chabrier, G., Allard, F., Hauschildt, P.H., 1998, A&A 337, 403

Barsony, M., Kenyon, S.J., Lada, E.A., Teuben, P.J., 1997, ApJS 112, 109

Beckwith, S.V., Sargent, A.I., Chini, R.S., Guesten, R., 1990, AJ 99, 924

Belloche, A., André, P., Motte, F., 2001. In: From Darkness to Light - Origin and Early Evolution of Young Stellar Clusters, Montmerle T. & André P. (eds.), ASP Conf. Ser., in press

Bernard, J.P., Boulanger, F., Puget, J.L., 1993, A&A 277, 609 Blommaert, J., 1998, ISOCAM Photometric Report, Experimental Astronomy, in press

Bontemps, S., André, P., Terebey, S., Cabrit, S., 1996, A&A 311, 858

Boulanger, F., Reach, W.T., Abergel, A., et al., 1996, A&A 315, L325

Calvet, N., Hartmann, L., Kenyon, S.J., Whitney, B.A., 1994, ApJ 434, 330

Casanova, S., Montmerle, T., Feigelson, E.D., André, P., 1995, ApJ 439, 752

Censori, C., D'Antona, F., 1998. In: Brown Dwarfs and Extrasolar Planets, Rebolo R., Martín E.L., Zapatero Osorio M.R. (eds.), ASP Conf. Ser., vol. 134, 518

Cesarsky, C., Abergel, A., Agnèse, P., et al., 1996, A&A, 315, L32

Chini, R., 1981, A&A 99, 346

Cohen, J.G., Frogel, J.A., Persson, S.E., Elias, J.H., 1981, ApJ 249, 481

Comerón, F., Rieke, G.H., Burrows, A., Rieke, M.J., 1993, ApJ 416, 185

D'Antona, F., Mazzitelli, I., 1994, ApJS 90, 467

D'Antona, F., Mazzitelli, I., 1998. In: Brown Dwarfs and Extrasolar Planets, Rebolo R., Martín E.L., Zapatero Osorio M.R. (eds.), ASP Conf. Ser., vol. 134, 442

de Geus, E.J., de Zeeuw, P.T., Lub, J., 1989, A&A 216, 44

de Zeeuw, P.T., Hoogerwerf, R., de Bruijne, J.H.J, Brown, A.G.A., Blaauw, A., 1999, AJ 117, 354

Duquennoy, A., Mayor, M., 1991, A&A 248, 485

Elias, J.H., 1978, ApJ 224, 453

Fletcher, A.B., Stahler, S.W., 1994, ApJ 435, 329

Garrison, R.F., 1967, ApJ 147, 1003

Greene, T.P., Young, E.T., 1992, ApJ 395, 516

Greene, T.P., Meyer, M.R., 1995, ApJ 450, 233

Greene, T.P., Lada, C.J., 1996, AJ 112, 2184

Greene, T.P., Wilking, B.A., André, P., Young, E.T., Lada, C.J., 1994, ApJ 434, 614

Grosso, N., Montmerle, T., Bontemps, S., André, P., Feigelson, E.D., 2000, A&A 359, 113

preparation

Gullbring, E., Hartmann, L., Briceño, C., Calvet, N. 1998, ApJ 492, 323

Hartigan, P., Strom, K.M., Strom, S.E., 1994, ApJ 427, 961 Kaas, A.A., Olofsson, G., Bontemps, S., et al., 2001, A&A in

Kaas, A.A., Bontemps, S., 2001. In: From Darkness to Light
 Origin and Early Evolution of Young Stellar Clusters,
 Montmerle T. & André P. (eds.), ASP Conf. Ser., in press

Kenyon, S.J., Hartmann, L., 1995, ApJS, 101, 117

Kenyon, S.J., Lada, E.A., Barsony, M., 1998, AJ 115, 252

Kessler, M.F., Steinz, J.A., Anderegg, M.E., et al., 1996, A&A 315, L27

Kroupa, P., Tout, C., Gilmore, G.F., 1993, MNRAS 262, 545Lada, C.J., 1987. In: Star Forming Regions, IAU Symposia, vol. 115, 1

Lada, C.J., Wilking, B.A., 1984, ApJ 287, 610

Lada, C.J., Adams, F.C., 1992, ApJ 393, 278

Lada, E.A., Strom, K.M., Myers, P.C., 1993. In: Protostars and Planets III, Mannings V., Boss A.P., Russel S.S. (eds.), University of Arizona Press, Tucson, p. 245

Leinert, C., Zinnecker, H., Weitzel, N., et al., 1993, A&A 278, 129

Liseau, R., Lorenzetti, D., Molinari, S., et al., 1995, A&A 300, 493

Loren, R.B., 1989, ApJ 338, 902

Loren, R.B., Wootten, A., Wilking, B.A., 1990, ApJ 365, 269Luhman, G.H., Rieke, G.H., 1999, ApJ 525, 440

Lutz, D., et al., 1999. In: The Universe as seen by ISO, Cox P., Kessler M.F. (eds.), ESA Publications Division, vol. SP-427, 623

Martín, E.L., Montmerle, T., Gregorio-Hetem, J., Casanova, S., 1998, MNRAS 300, 733

Meyer, M.R., Calvet, N., Hillenbrand, L.A., 1997, AJ 114, 288
Meyer, M.R., Adams, F.C., Hillenbrand, L.A., Carpenter,
J.M., Larson, R.B., 2000. In: Protostars and Planets IV,
Mannings V., Boss A.P., Russell S.S. (eds.), University of
Arizona Press, Tucson, p. 121

Miller, G.E., Scalo, J.M., 1979, ApJS 41, 513

Montmerle, T., Koch-Miramond, L., Falgarone, E., Grindlay, J.E., 1983, ApJ 269, 182

Moore, T.J.T., et al., 1998, MNRAS 299, 1209

Motte, F., André, P., Neri, R., 1998, A&A 336, 150 (MAN98)

Nordh, L., Olofsson, G., Abergel, A., et al., 1996, A&A 315, L185

Nordh, L., Olofsson, G., Bontemps, S., et al., 1998. In: Star Formation with ISO, Yun J.L., Liseau R. (eds.), ASP Conf. Ser., vol. 132, 127

Olofsson, G., Huldtgren, M., Kaas, A.A., et al., 1999, A&A 350, 883

Palla, F., Stahler, S.W., 1999, ApJ 525, 772

Persi, P., Marenzi, A.R., Olofsson, G., et al., 2000, A&A 357, 219

Piskunov, A., Belikov, A., 1996, Astron. Lett. 22-4, 466

Preibisch, T., Zinnecker, H., 1999, AJ 117, 2381

Rieke, G.H., Lebofsky, M.J., 1985, ApJ 288, 618

Salpeter, E., 1955, ApJ 121, 161

Scalo, J., 1998. In: The Stellar Initial Mass Function, Gilmore G., Howell D. (eds.), ASP Conf. Ser., vol. 142, 201

Shu, F.H., Adams, F.C., Lizano, S., 1987, ARA&A 25, 23

Simon, M., Ghez, A.M., Leinert, C., et al., 1995, ApJ 443, 625 Stahler, S.W, & Walter, F.M. 1993. In: Protostars and Planets

III, Mannings V., Boss A.P., Russel S.S. (eds.), University of Arizona Press, Tucson, p. 405

Starck, J.L., Murtagh, F., Pirenne, B., Albrecht, M. 1996, PASP 108, 446

Starck, J.L., Abergel, A., Aussel, H., et al. 1999, A&AS 134, 135

Stine, P.C., Feigelson, E.D., André, P., Montmerle, T., 1988, AJ 96, 1394

Strom, K.M., Strom, S.E., Edwards, S., Cabrit, S., Skrutskie, M.F., 1989, AJ 97, 1451

Strom, K.M., Kepner, J., Strom, S.E., 1995, ApJ 438, 813

Wainscoat, R.J., Cohen, M., Volk, K., Walker, H.J., Schwartz, D.E., 1992, ApJS 83, 111

Wilking, B.A., Lada, C.J., 1983, ApJ 274, 698

Wilking, B.A., Schwartz, R.D., Blackwell, J.H., 1987, AJ 94, 106

Wilking, B.A., Lada, C.J., Young, E.T., 1989, ApJ 340, 823 (WLY89)

Wilking, B.A., Greene, T.P., Meyer, M.R., 1999, AJ 117, 469Wilking, B.A., Bontemps, S., Schuler, R.E., Greene, T.P., André, P., 2001, ApJ, in press

Williams, D.M., Comerón, F., Rieke, G.H., Rieke, M.J., 1995, ApJ 454, 144

Wilson, T.L., Mauersberger, R., Gensheimer, P.D., Muders, D., Bieging, J.H., 1999, ApJ 525, 343

Zinnecker, H., McCaughrean, M.J., Wilking, B.A., 1993. In: Protostars and Planets III, Mannings V., Boss A.P., Russel S.S. (eds.), University of Arizona Press, Tucson, p. 429

Table 1. List of the 212 ISOCAM sources identified as members of the ρ Ophiuchi cluster

700	<i>C</i>	(10000)2	F 6.7	Ь	F14 3	h	T1 (10 (1) 0	TD d
ISO		$es(J2000)^{a}$	$F_{\nu}^{6.7}$	$\sigma_{6.7}^{600000000000000000000000000000000000$	$F_{\nu}^{14.3}$	$\sigma_{14.3}^{b}$	Identification ^c	IR^d
#	α	δ	[Jy]	[Jy]	[Jy]	[Jy]		Class
1	$16^{\rm h}25^{\rm m}36.8$	$-24^{\circ}15'51''$	0.157	0.031	0.249	0.023	IRS2	II
2	$16^{\rm h}25^{\rm m}38^{\rm s}.2$	$-24^{\circ}22'40''$	0.075	0.014	0.088	0.007	B162538 - 242238	nII
3	$16^{\rm h}25^{\rm m}39^{\rm s}.7$	$-24^{\circ}26'37''$	0.197	0.020	0.449	0.013	IRS3	nII
4	16 ^h 25 ^m 41.1	$-24^{\circ}21'38''$	0.008	0.008	-	-	B162541-242138	III
5	16 ^h 25 ^m 50 ^s .6	$-24^{\circ}39'18''$	0.045	0.012	-	-	IRS10	III
6	16 ^h 25 ^m 56 ^s 3	$-24^{\circ}20'50''$	0.400	0.042	0.662	0.024	SR4/IRS12	II
7	16 ^h 25 ^m 57 ^s .5 16 ^h 26 ^m 01 ^s .5	$-24^{\circ}30'34''$	0.049	0.007	0.009	0.006	GSS20	III
8 9	16 ^h 26 ^m 01.5 16 ^h 26 ^m 01.5	$-24^{\circ}29'47''$ $-24^{\circ}25'20''$	$0.034 \\ 0.025$	0.010 0.009	0.032	0.007	B162601-242945 SKS1-4	III nII
10	16 26 01.5 16 ^h 26 ^m 03.2	$-24^{\circ}23'40''$	0.025 0.507	0.009 0.017	0.032 0.341	0.007 0.024	DoAr21/GSS23	III
11	16 26 03.2 16 ^h 26 ^m 03.2	$-24^{\circ}17'49''$	0.025	0.017 0.002	0.041	0.024 0.003	VSSG19	III
12	16 26 03.2 16 ^h 26 ^m 04.6	$-24^{\circ}17'52''$	0.025	0.002	0.000	0.003	B162604-241753	nII
13	16 ^h 26 ^m 07 ^s 0	$-24^{\circ}17'32'$ $-24^{\circ}27'28''$	0.003 0.112	0.002 0.022	0.012 0.128	0.003	B162607-242725	nII
14	16 ^h 26 ^m 07 ^s .5	$-24^{\circ}27'43''$	0.032	0.007	-	-	B162607-242742	III
15	16 ^h 26 ^m 08.7	$-24^{\circ}18'53''$	0.010	0.003	_	_	CRBR4	III?
16	16 ^h 26 ^m 09 ^s .7	$-24^{\circ}34'12''$	_e	-	$_{-}^{e}$	_	SR3/GSS25	III
17	$16^{\rm h}26^{\rm m}10^{\rm s}.4$	$-24^{\circ}20'58''$	0.386	0.011	0.43	0.017	GSS26	II
18	$16^{\rm h}26^{\rm m}15.8$	$-24^{\circ}19'23''$	0.026	0.002	0.005	0.003	SKS1-7	III?
19	$16^{\rm h}26^{\rm m}17^{\rm s}.0$	$-24^{\circ}22'26''$	0.262	0.005	0.414	0.012	GSS29	II
20	$16^{\rm h}26^{\rm m}17\stackrel{\rm s}{.}2$	$-24^{\circ}20'23''$	0.146	0.003	0.293	0.004	DoAr24/GSS28	II
21	$16^{\rm h}26^{\rm m}17\stackrel{\rm s}{.}3$	$-24^{\circ}23'49''$	0.246	0.004	0.363	0.004	CRBR12	nI
22	$16^{\rm h}26^{\rm m}18\stackrel{\rm s}{.}6$	$-24^{\circ}17'11''$	0.010	0.001	-	-	B162618 - 241712	III?
23	$16^{\rm h}26^{\rm m}18.8$	$-24^{\circ}26'13''$	0.033	0.008	0.027	0.007	SKS1-10	nII
24	16 ^h 26 ^m 18 ^s .9	$-24^{\circ}28'22''$	0.673	0.025	0.575	0.024	VSSG1	II
25	16 ^h 26 ^m 19 ^s .0	$-24^{\circ}23'07''$	0.051	0.008	-	-	CRBR17	III?
26	16 ^h 26 ^m 19 ^s 2	$-24^{\circ}24'16''$	0.056	0.006	0.074	0.004	CRBR15	nII
27	16 ^h 26 ^m 20 ^s .7	$-24^{\circ}08'48''$	0.030	0.010	0.014	0.006	WSB28	III
28	16 ^h 26 ^m 21.50	$-24^{\circ}15'42''$	0.037	0.004	0.013	0.005	B162621-241544	III?
29	16 ^h 26 ^m 21 ^s 5	$-24^{\circ}23'07''$	4.09	0.12	19.1	0.26	GSS30/GY6	I
30	$16^{\rm h}26^{\rm m}21.7$ $16^{\rm h}26^{\rm m}21.8$	$-24^{\circ}26'02''$ $-24^{\circ}22'49''$	0.028	0.007	0.033	0.014	GY5	nII I
$\frac{31}{32}$	16 26 21.8 16 ^h 26 ^m 22.2	$-24^{\circ}22^{\circ}49$ $-24^{\circ}44'36''$	$0.066 \\ 0.027$	0.013 0.009	0.119	0.025	LFAM1 GY3	nII
32 33	16 26 22.2 16 ^h 26 ^m 22 ^s .4	$-24^{\circ}44^{\circ}30$ $-24^{\circ}24'08''$	0.027 0.009	0.009	$0.028 \\ 0.017$	$0.012 \\ 0.003$	GY11	nII
34	16 26 22.4 16 ^h 26 ^m 22.6	$-24^{\circ}22'56''$	0.009 0.052	0.003	0.017	0.003	GY12	III
$\frac{34}{35}$	16 26 22.0 16 ^h 26 ^m 23.1	$-24^{\circ}28'49''$	0.032 0.018	0.008	0.034 0.026	0.010 0.014	GY15	nII
36	16 ^h 26 ^m 23 ^s .4	$-24^{\circ}21'02''$	2.59	0.068	2.56	0.014	GSS31/GY20	II
37	16 ^h 26 ^m 23 ^s .7	$-24^{\circ}24'40''$	0.395	0.005	0.405	0.007	LFAM3/GY21	II
38	$16^{\rm h}26^{\rm m}24^{\rm s}.0$	$-24^{\circ}43'09''$	0.174	0.012	0.206	0.017	DoAr25/GY17	II
39	$16^{\rm h}26^{\rm m}24.1$	$-24^{\circ}24'50''$	2.65	0.10	2.16	0.067	S2/GY23	II
40	$16^{\rm h}26^{\rm m}24.2$	$-24^{\circ}16'14''$	2.73	0.027	3.02	0.030	EL24	II
41	$16^{\rm h}26^{\rm m}25^{\rm s}.3$	$-24^{\circ}24'47''$	0.064	0.008	0.049	0.011	GY29	$_{ m nII}$
42	$16^{\rm h}26^{\rm m}27^{\rm s}.7$	$-24^{\circ}26'43''$	0.003	0.001	_	_	VSSG29/GY37	III?
43	$16^{\rm h}26^{\rm m}28^{\rm s}.0$	$-24^{\circ}41'51''$	0.029	0.002	0.115	0.006	GY33	$_{ m nII}$
44	$16^{\rm h}26^{\rm m}28\rlap.{}^{\rm s}5$	$-24^{\circ}15'45''$	0.023	0.004	0.010	0.004	B162628 - 241543	III?
45	$16^{\rm h}26^{\rm m}29^{\rm s}.7$	$-24^{\circ}19'08''$	0.015	0.002	0.005	0.003	LFAM8/SKS1-19	III
46	$16^{\rm h}26^{\rm m}30^{\rm s}.5$	$-24^{\circ}22'59''$	0.347	0.094	0.303	0.065	VSSG27/GY51	II
47	$16^{\rm h}26^{\rm m}30^{\rm s}.9$	$-24^{\circ}31'07''$	0.012	0.008	-	-	IRS14/GY54	III?
48	$16^{\rm h}26^{\rm m}34^{\rm s}.2$	$-24^{\circ}23'30''$	_ ^e	-	$\underline{}^e$	-	$\mathrm{S1/GY70}^f$	III
49	$16^{\rm h}26^{\rm m}36^{\rm s}.7$	$-24^{\circ}18'10''$	0.007	0.002	-	-	B162636 - 241811	III?
50	16 ^h 26 ^m 36 ^s .8	$-24^{\circ}19'01''$	0.007	0.003	-	-	B162636-241902	III?
51	16 ^h 26 ^m 36 ^s 9	$-24^{\circ}15'54''$	0.307	0.004	0.636	0.009	B162636-241554	nII
52	16 ^h 26 ^m 38 ^s .0	$-24^{\circ}23'01''$	0.057	0.049	0.066	0.059	VSSG4/GY81	nII
53	16 ^h 26 ^m 38 ^s .6	$-24^{\circ}23'10''$	0.035	0.045	0.046	0.043	$GY84^g$	nII
54	16 ^h 26 ^m 40 ^s .6	$-24^{\circ}27'16''$	0.089	0.003	0.158	0.004	GY91/CRBR42	I
55	$16^{\rm h}26^{\rm m}40\stackrel{\rm s}{.}7$	$-24^{\circ}30'53''$	0.004	0.002	-	-	IRS16/GY92	III?

					0 0		p	
56	$16^{\rm h}26^{\rm m}41\stackrel{\rm s}{.}6$	$-24^{\circ}40'15''$	0.060	0.005	0.045	0.006	WSB37/GY93	II
57	$16^{\rm h}26^{\rm m}41\stackrel{\rm s}{.}8$	$-24^{\circ}18'01''$	0.011	0.002	-	-	B162641 - 241801	III?
58	$16^{\rm h}26^{\rm m}42^{\rm s}.0$	$-24^{\circ}33'24''$	0.172	0.026	0.089	0.014	WL8/GY96	III
59	$16^{\rm h}26^{\rm m}42^{\rm s}.1$	$-24^{\circ}31'02''$	0.047	0.002	0.059	0.003	WL7/GY98	II
60	$16^{\rm h}26^{\rm m}42^{\rm s}.3$	$-24^{\circ}26'26''$	0.019	0.002	-	-	GY101	III
61	$16^{\rm h}26^{\rm m}42^{\rm s}.3$	$-24^{\circ}26'34''$	0.012	0.001	-	-	GY103	III
62	$16^{\rm h}26^{\rm m}42^{\rm s}.9$	$-24^{\circ}20'32''$	0.284	0.007	0.290	0.008	GSS37/GY110	II
63	$16^{\rm h}26^{\rm m}43^{\rm s}.1$	$-24^{\circ}23'01''$	0.024	0.008	0.028	0.007	GY109	nII
64	$16^{\rm h}26^{\rm m}43^{\rm s}.8$	$-24^{\circ}16'35''$	0.033	0.002	0.009	0.003	VSSG11	III
65	$16^{\rm h}26^{\rm m}44^{\rm s}.0$	$-24^{\circ}34'48''$	1.27	0.043	2.57	0.029	WL12/GY111	I
66	$16^{\rm h}26^{\rm m}44^{\rm s}.3$	$-24^{\circ}43'18''$	0.020	0.004	-	-	GY112	III
67	$16^{\rm h}26^{\rm m}45^{\rm s}.1$	$-24^{\circ}23'10''$	0.291	0.013	0.307	0.012	GSS39/GY116	II
68	$16^{\rm h}26^{\rm m}46^{\rm s}.5$	$-24^{\circ}12'03''$	0.352	0.006	0.240	0.004	VSS27	II
69	$16^{\rm h}26^{\rm m}47^{\rm s}.0$	$-24^{\circ}44'30''$	0.011	0.002	-	-	GY122	III
70	$16^{\rm h}26^{\rm m}48{\stackrel{\rm s}{.}}6$	$-24^{\circ}28'39''$	0.137	0.004	0.188	0.005	WL2/GY128	II
71	$16^{\rm h}26^{\rm m}48^{\rm s}.7$	$-24^{\circ}26'28''$	0.006	0.002	-	-	GY130	III?
72	$16^{\rm h}26^{\rm m}49\stackrel{\rm s}{.}0$	$-24^{\circ}38'27''$	0.101	0.002	0.086	0.002	WL18/GY129	II
73	$16^{\rm h}26^{\rm m}49^{\rm s}.2$	$-24^{\circ}20'05''$	0.074	0.004	0.022	0.004	VSSG3/GY135	III
74	$16^{\rm h}26^{\rm m}50^{\rm s}_{\cdot}9$	$-24^{\circ}20'52''$	0.019	0.003	-	-	IRS20/GY143	III?
75	$16^{\rm h}26^{\rm m}52^{\rm s}.1$	$-24^{\circ}30'39''$	0.026	0.002	0.026	0.003	GY144	$_{ m nII}$
76	$16^{\rm h}26^{\rm m}53^{\rm s}.6$	$-24^{\circ}32'36''$	0.022	0.003	0.017	0.004	GY146	$_{ m nII}$
77	$16^{\rm h}26^{\rm m}54^{\rm s}.4$	$-24^{\circ}24'38''$	0.020	0.003	0.017	0.004	GY152	$_{ m nII}$
78	$16^{\rm h}26^{\rm m}54^{\rm s}.5$	$-24^{\circ}26'22''$	0.091	0.003	0.070	0.004	VSSG5/GY153	II
79	$16^{\rm h}26^{\rm m}54^{\rm s}.9$	$-24^{\circ}27'05''$	0.013	0.003	0.013	0.004	GY154	$_{ m nII}$
80	$16^{\rm h}26^{\rm m}55^{\rm s}.0$	$-24^{\circ}22'31''$	0.026	0.004	0.008	0.004	GY156	III
81	$16^{\rm h}26^{\rm m}55^{\rm s}.3$	$-24^{\circ}20'29''$	0.055	0.002	0.010	0.003	VSSG7/GY157	III?
82	$16^{\rm h}26^{\rm m}56^{\rm s}.7$	$-24^{\circ}28'38''$	0.003	0.001	-	-	GY163	III?
83	$16^{\rm h}26^{\rm m}57^{\rm s}.3$	$-24^{\circ}14'00''$	0.114	0.005	0.084	0.004	B162656 - 241353	$_{ m nII}$
84	$16^{\rm h}26^{\rm m}57^{\rm s}.4$	$-24^{\circ}35'39''$	0.009	0.003	0.010	0.005	WL21/GY164	$_{ m nII}$
85	$16^{\rm h}26^{\rm m}58\stackrel{\rm s}{.}3$	$-24^{\circ}37'40''$	0.015	0.004	0.011	0.007	CRBR51	$_{ m nII}$
86	$16^{\rm h}26^{\rm m}58^{\rm s}.4$	$-24^{\circ}21'29''$	0.022	0.003	0.018	0.004	IRS26/GY171	$_{ m nII}$
87	$16^{\rm h}26^{\rm m}58^{\rm s}.6$	$-24^{\circ}18'34''$	0.020	0.003	0.014	0.003	B162658-241836	nII
88	$16^{\rm h}26^{\rm m}58^{\rm s}.8$	$-24^{\circ}45'37''$	1.41	0.090	2.29	0.17	$\mathrm{SR}24^h$	II
89	$16^{\rm h}26^{\rm m}59^{\rm s}.3$	$-24^{\circ}35'57''$	0.004	0.002	0.008	0.005	WL14/GY172	nII
90	$16^{\rm h}26^{\rm m}59^{\rm s}.3$	$-24^{\circ}35'01''$	_e	-	_e	-	WL22/GY174	II
91	$16^{\rm h}27^{\rm m}01^{\rm s}.8$	$-24^{\circ}21'37''$	0.060	0.003	0.013	0.003	VSSG8/GY181	III?
92	$16^{\rm h}27^{\rm m}02^{\rm s}.5$	$-24^{\circ}37'30''$	_e	-	\mathbf{e}^{e}	-	WL16/GY182	II
93	$16^{\rm h}27^{\rm m}03^{\rm s}.0$	$-24^{\circ}26'16''$	0.026	0.003	0.021	0.004	GY188	nII
94	$16^{\rm h}27^{\rm m}03^{\rm s}.7$	$-24^{\circ}20'07''$	0.008	0.002	0.007	0.003	B162703-242007	nII
95	16 ^h 27 ^m 04 ^s .1	$-24^{\circ}28'33''$	0.103	0.006	0.164	0.004	WL1/GY192	II
96	16 ^h 27 ^m 04 ^s .4	$-24^{\circ}43'00''$	0.020	0.003	-	-	GY193	III
97	16 ^h 27 ^m 04 ^s .6	$-24^{\circ}42'15''$	0.017	0.003	-	-	GY194	III
98	16 ^h 27 ^m 04 ^s .6	$-24^{\circ}27'16''$	0.046	0.006	0.066	0.008	GY195	nII
99	16 ^h 27 ^m 05 ^s .4	$-24^{\circ}36'31''$	0.032	0.005	0.064	0.009	LFAM26/GY197	I
100	16 ^h 27 ^m 05 ^s .7	$-24^{\circ}40'12''$	0.007	0.002	-	-	B162705-244013	III?
101	16 ^h 27 ^m 06 ^s .0	$-24^{\circ}26'19''$	0.025	0.002	-	-	IRS30/GY203	III
102	16 ^h 27 ^m 06 ^s .6	$-24^{\circ}41'48''$	0.024	0.002	0.020	0.004	GY204	nII
103	16 ^h 27 ^m 07 ^s .0	-24°38′16″	0.449	0.006	0.732	0.015	WL17/GY205	II
104	16 ^h 27 ^m 07 ^s .9	$-24^{\circ}40'27''$	0.004	0.002	- 0.150	-	GY207	III?
105	16 ^h 27 ^m 09 ^s .2	$-24^{\circ}34'10''$	0.174	0.005	0.172	0.008	WL10/GY211	II
106	16 ^h 27 ^m 09 ^s .3	$-24^{\circ}12'07''$	0.058	0.003	0.044	0.003	B162708-241204	nII
107	16 ^h 27 ^m 09 ^s .6	$-24^{\circ}40'25''$	0.079	0.002	0.096	0.004	GY213	nII
108	16 ^h 27 ^m 09 ^s .6	$-24^{\circ}37'21''$	24.0	0.66	26.2	0.50	EL29/GY214	III.5
109	$16^{\rm h}27^{\rm m}09^{\rm s}.6$ $16^{\rm h}27^{\rm m}10^{\rm s}.2$	$-24^{\circ}29'55''$	0.014	0.006	2 94	- 0.026	GY215	III?
110	16 ^h 27 ^m 10.2 16 ^h 27 ^m 10.3	$-24^{\circ}19'16''$ $-24^{\circ}33'22''$	1.57	0.026	2.84	0.026	SR21/VSSG23	II 1112
111	16 ^h 27 ^m 10.3 16 ^h 27 ^m 11.4	$-24^{\circ}33^{\circ}22^{\circ}$ $-24^{\circ}40'46''$	0.009	0.002	- 0.301	- 0.010	WL9/GY220 GY224	III?
112 113	16 27 11.4 16 ^h 27 ^m 11.7	$-24^{\circ}40^{\circ}40'$ $-24^{\circ}23'49''$	0.221 0.034	$0.006 \\ 0.010$	0.391	0.010	IRS32/GY228	II III?
113	16 27 11.7 16 ^h 27 ^m 11.9	$-24^{\circ}38'31''$	0.034 0.230	0.010	0.126	0.012	WL19/GY227	III! II/III?
114	10 21 11.8	27 JO J1	0.200	0.000	0.120	0.012	W 110/ G 1 221	11/111;

115	$16^{\rm h}27^{\rm m}12^{\rm s}.1$	$-24^{\circ}34'48''$	0.024	0.003	0.019	0.006	WL11/GY229	nII
116	$16^{\rm h}27^{\rm m}13\stackrel{\rm s}{.}9$	$-24^{\circ}18'25''$	0.074	0.005	0.073	0.005	B162713-241818	$_{ m nII}$
117	$16^{\rm h}27^{\rm m}14\stackrel{\rm s}{.}0$	$-24^{\circ}43'31''$	0.093	0.003	0.168	0.006	GY235	$_{ m nII}$
118	$16^{\rm h}27^{\rm m}14^{\rm s}.6$	$-24^{\circ}26'55''$	0.119	0.009	0.101	0.009	IRS33/GY236	nII
119	$16^{\rm h}27^{\rm m}15.4$	$-24^{\circ}30'54''$	0.054	0.005	0.061	0.007	IRS35/GY238	nII
120	16 ^h 27 ^m 15.7	$-24^{\circ}36'34''$ $-24^{\circ}26'46''$	0.054 0.167	0.003	0.001 0.152	0.014	IRS34/GY239	II
		$-24^{\circ}38'46''$					•	
121	16 ^h 27 ^m 15 ^s .9		0.151	0.004	0.782	0.010	WL20/GY240	II
122	16 ^h 27 ^m 16 ^s .1	$-24^{\circ}25'22''$	0.018	0.010	0.019	0.005	IRS36/GY241	nII
123^{*}	$16^{\rm h}27^{\rm m}17^{\rm s}.6$	$-24^{\circ}05'19''$	0.021	0.002	0.035	0.004	ISO1627176-240519	nII
124	16 ^h 27 ^m 17 ^s .6	$-24^{\circ}28'58''$	0.286	0.011	0.347	0.009	IRS37/GY244	II
125	$16^{\rm h}27^{\rm m}18\stackrel{\rm s}{.}0$	$-24^{\circ}28'55''$	0.186	0.014	0.049	0.017	WL5/GY246	III
126	$16^{\rm h}27^{\rm m}18\stackrel{\rm s}{.}0$	$-24^{\circ}24'31''$	0.028	0.004	-	-	GY248	III
127	$16^{\rm h}27^{\rm m}18\stackrel{\rm s}{.}5$	$-24^{\circ}39'15''$	0.075	0.004	0.096	0.009	GY245	II
128	$16^{\rm h}27^{\rm m}18\stackrel{\rm s}{.}6$	$-24^{\circ}29'07''$	0.284	0.019	0.272	0.012	WL4/GY247	II
129	$16^{\rm h}27^{\rm m}19\stackrel{\rm s}{.}3$	$-24^{\circ}28'45''$	0.285	0.015	0.218	0.006	WL3/GY249	II
130	$16^{\rm h}27^{\rm m}19^{\rm s}.5$	$-24^{\circ}41'43''$	0.053	0.004	_	_	SR12/GY250	III
131	$16^{\rm h}27^{\rm m}21.6$	$-24^{\circ}21'52''$	0.014	0.004	_	_	GY255	III?
132	$16^{\rm h}27^{\rm m}21.7$	$-24^{\circ}41'42''$	2.68	0.096	2.13	0.057	IRS42/GY252	II.
133	16 ^h 27 ^m 21 ^s .8	$-24^{\circ}43'34''$	0.035	0.002	0.008	0.004	GY253	III
134	16 27 21.8 16 ^h 27 ^m 21.8	$-24^{\circ}43^{\circ}54^{\circ}$ $-24^{\circ}29'55''$	1.09		1.26		WL6/GY254	I
	16 27 21.8 16 ^h 27 ^m 22 ^s .9	$-24^{\circ}29^{\circ}35$ $-24^{\circ}18'00''$		0.029		0.033	•	
135			0.049	0.003	0.011	0.003	VSSG22	III
136	16 ^h 27 ^m 24 ^s .3	$-24^{\circ}41'46''$	0.008	0.002	-	-	GY258	III?
137	$16^{\rm h}27^{\rm m}24.8$	$-24^{\circ}41'03''$	0.177	0.006	0.362	0.015	CRBR85	Ι
138	$16^{\rm h}27^{\rm m}26\stackrel{\rm s}{.}0$	$-24^{\circ}19'28''$	0.006	0.002	0.006	0.002	B162726 - 241925	nII
139	$16^{\rm h}27^{\rm m}26^{\rm s}.2$	$-24^{\circ}42'45''$	0.032	0.003	0.039	0.006	GY260	nII
140	$16^{\rm h}27^{\rm m}26\stackrel{\rm s}{.}8$	$-24^{\circ}39'22''$	0.160	0.005	0.197	0.019	GY262	II
141	$16^{\rm h}27^{\rm m}27^{\rm s}.1$	$-24^{\circ}40'51''$	2.21	0.019	4.43	0.091	IRS43/GY265	I
142	$16^{\rm h}27^{\rm m}27^{\rm s}.5$	$-24^{\circ}31'18''$	0.095	0.004	0.098	0.011	VSSG25/GY267	II
143	$16^{\rm h}27^{\rm m}28^{\rm s}.3$	$-24^{\circ}39'33''$	4.19	0.14	11.4	0.12	IRS44/GY269	I
144	$16^{\rm h}27^{\rm m}28^{\rm s}\!.4$	$-24^{\circ}27'21''$	0.407	0.016	0.481	0.018	IRS45/GY273	II
145	$16^{\rm h}27^{\rm m}29^{\rm s}.7$	$-24^{\circ}39'16''$	0.294	0.004	0.649	0.011	IRS46/GY274	I
146	$16^{\rm h}27^{\rm m}29.7$	$-24^{\circ}33'36''$	0.013	0.002	-	-	GY278	III?
147	$16^{\rm h}27^{\rm m}30^{\rm s}.1$	$-24^{\circ}27'43''$	1.78	0.002	1.54	0.017	IRS47/GY279	II
148	16 ^h 27 ^m 30.8	$-24^{\circ}34'04''$	0.014	0.002	-	0.011	GY283	III?
149	16 27 30.8 16 ^h 27 ^m 30.9	$-24^{\circ}47'25''$		0.002	0.012	0.006		III
	16 27 30.9 16 ^h 27 ^m 30.9		0.029			0.006	B162730-244726	
150*		$-24^{\circ}27'34''$	0.078	0.008	0.094	0.007	ISO1627309-242734	nII
151	16 ^h 27 ^m 31 ^s .0	$-24^{\circ}25'00''$	0.059	0.007	0.098	0.008	GY284	nII
152	$16^{\rm h}27^{\rm m}32^{\rm s}.5$	$-24^{\circ}33'22''$	0.030	0.002	0.018	0.004	GY289	III
153	$16^{\rm h}27^{\rm m}32\stackrel{\rm s}{.}6$	$-24^{\circ}32'42''$	0.020	0.003	-	-	GY290	III?
154	$16^{\rm h}27^{\rm m}32^{\rm s}.8$	$-24^{\circ}32'34''$	0.058	0.002	0.061	0.006	GY291	$_{ m nII}$
155	$16^{\rm h}27^{\rm m}33^{\rm s}2$	$-24^{\circ}41'15''$	0.391	0.006	0.534	0.013	GY292	II
156	$16^{\rm h}27^{\rm m}35^{\rm s}.2$	$-24^{\circ}38'31''$	0.017	0.002	-	-	GY295	III?
157	$16^{\rm h}27^{\rm m}35^{\rm s}.6$	$-24^{\circ}45'31''$	0.010	0.002	-	-	GY296	III
158	$16^{\rm h}27^{\rm m}36^{\rm s}.3$	$-24^{\circ}28'34''$	0.006	0.003	_	_	GY297	III?
159	$16^{\rm h}27^{\rm m}37^{\rm s}2$	$-24^{\circ}30'34''$	5.87	0.11	6.49	0.099	IRS48/GY304	Ι
160	16 ^h 27 ^m 37 ^s .4	$-24^{\circ}17'58''$	0.011	0.002	0.010	0.003	B162737-241756	nII
161	16 ^h 27 ^m 37 ^s .6	$-24^{\circ}42'40''$	0.071	0.003	0.161	0.008	GY301	nII
162	16 ^h 27 ^m 38.0	$-24^{\circ}25'25''$	0.005	0.003	-	-	GY309	III?
163	16 27 38.0 16 ^h 27 ^m 38.4	$-24^{\circ}36'58''$	0.303	0.003	0.332			III:
						0.010	IRS49/GY308	
164	16 ^h 27 ^m 38 ^s 8	$-24^{\circ}38'39''$	0.020	0.002	0.039	0.006	GY310	nII
165	16 ^h 27 ^m 39 ^s .1	$-24^{\circ}40'21''$	0.049	0.002	0.077	0.008	GY312	nII
166	$16^{\rm h}27^{\rm m}39.5$	$-24^{\circ}39'16''$	0.266	0.005	0.411	0.008	GY314	II
167	$16^{\rm h}27^{\rm m}40^{\rm s}.0$	$-24^{\circ}43'13''$	0.924	0.016	1.06	0.014	IRS51/GY315	Ι
168	$16^{\rm h}27^{\rm m}40^{\rm s}.5$	$-24^{\circ}22'07''$	0.464	0.009	0.449	0.007	SR9/IRS52	II
169	$16^{\rm h}27^{\rm m}41^{\rm s}.5$	$-24^{\circ}35'37''$	0.024	0.002	-	-	GY322	III?
170	$16^{\rm h}27^{\rm m}41\stackrel{\rm s}{.}8$	$-24^{\circ}46'45''$	0.026	0.002	0.043	0.006	B162741 - 244645	nII
171	$16^{\rm h}27^{\rm m}41\stackrel{\rm s}{.}9$	$-24^{\circ}43'37''$	0.050	0.002	0.058	0.005	GY323	nII
172	$16^{\rm h}27^{\rm m}42^{\rm s}.7$	$-24^{\circ}38'49''$	0.021	0.004	0.013	0.007	GY326	$_{ m nII}$
173	$16^{\rm h}27^{\rm m}43^{\rm s}.7$	$-24^{\circ}43'07''$	0.027	0.003	-	-	IRS53/GY334	III?
							,	

	1							
174	$16^{\rm h}27^{\rm m}45^{\rm s}.9$	$-24^{\circ}38'00''$	0.003	0.002	-	-	GY346	III?
175	$16^{\rm h}27^{\rm m}46^{\rm s}.0$	$-24^{\circ}44'52''$	0.061	0.002	0.082	0.004	GY344	nII
176	$16^{\rm h}27^{\rm m}46^{\rm s}.2$	$-24^{\circ}31'40''$	0.024	0.003	0.022	0.005	GY350	nII
177	$16^{\rm h}27^{\rm m}47^{\rm s}2$	$-24^{\circ}45'33''$	0.047	0.003	0.037	0.007	GY352	nII
178	$16^{\rm h}27^{\rm m}49^{\rm s}.9$	$-24^{\circ}25'25''$	0.055	0.005	0.041	0.005	GY371	nII
179	$16^{\rm h}27^{\rm m}50^{\rm s}.0$	$-24^{\circ}44'15''$	0.009	0.002	-	-	GY370	III?
180	$16^{\rm h}27^{\rm m}50^{\rm s}.2$	$-24^{\circ}25'43''$	0.182	0.005	0.045	0.006	VSSG14/GY372	III
181	$16^{\rm h}27^{\rm m}50^{\rm s}.3$	$-24^{\circ}39'01''$	0.004	0.002	-	-	GY373	III?
182	$16^{\rm h}27^{\rm m}51\stackrel{\rm s}{.}7$	$-24^{\circ}31'46''$	4.65	0.094	5.3	0.11	IRS54/GY378	I
183	$16^{\rm h}27^{\rm m}51\stackrel{\rm s}{.}9$	$-24^{\circ}46'28''$	0.015	0.002	-	-	GY377	III
184	$16^{\rm h}27^{\rm m}52\stackrel{\rm s}{.}0$	$-24^{\circ}40'48''$	0.067	0.002	0.017	0.004	IRS55/GY380	III
185	$16^{\rm h}27^{\rm m}55\stackrel{ m s}{.}0$	$-24^{\circ}28'39''$	0.025	0.002	0.026	0.002	GY397	nII
186	$16^{\rm h}27^{\rm m}55.5$	$-24^{\circ}44'49''$	0.011	0.002	-	-	GY398	III?
187	$16^{\rm h}27^{\rm m}55\stackrel{ m s}{.}9$	$-24^{\circ}26'22''$	0.102	0.003	0.101	0.004	SR10/GY400	II
188	$16^{\rm h}27^{\rm m}57^{\rm s}.7$	$-24^{\circ}40'01''$	0.017	0.001	-	-	GY410	III
189	$16^{\rm h}27^{\rm m}57^{\rm s}.8$	$-24^{\circ}36'01''$	0.004	0.003	-	-	GY412	III?
190	$16^{\rm h}28^{\rm m}03.8$	$-24^{\circ}34'43''$	0.009	0.001	0.012	0.003	GY450	nII
191	$16^{\rm h}28^{\rm m}04\stackrel{\rm s}{.}9$	$-24^{\circ}34'52''$	0.036	0.002	0.012	0.003	GY463	III?
192	$16^{\rm h}28^{\rm m}05.5$	$-24^{\circ}33'55''$	0.002	0.001	-	-	GY472	III?
193	$16^{\rm h}28^{\rm m}12\stackrel{\rm s}{.}2$	$-24^{\circ}11'37''$	0.018	0.003	0.025	0.004	B162812 - 241138	nII
194	$16^{\rm h}28^{\rm m}13\stackrel{\rm s}{.}7$	$-24^{\circ}32'48''$	0.052	0.001	0.040	0.002	B162813 - 243249	nII
195*	$16^{\rm h}28^{\rm m}16\stackrel{\rm s}{.}8$	$-24^{\circ}05'19''$	0.102	0.004	0.065	0.004	ISO1628168 - 240519	nII
196	$16^{\rm h}28^{\rm m}16\stackrel{\rm s}{.}8$	$-24^{\circ}37'04''$	0.108	0.002	0.128	0.002	WSB60	II
197	$16^{\rm h}28^{\rm m}22^{\rm s}.1$	$-24^{\circ}42'49''$	0.017	0.001	0.010	0.002	B162821 - 244246	nII
198	$16^{\rm h}28^{\rm m}33\stackrel{\rm s}{.}2$	$-24^{\circ}22'46''$	0.701	0.020	0.265	0.004	SR20	III
199	$16^{\rm h}28^{\rm m}45\stackrel{\rm s}{.}6$	$-24^{\circ}28'22''$	0.315	0.004	0.230	0.006	SR13	II
200*	$16^{\rm h}31^{\rm m}43\stackrel{\rm s}{.}4$	$-24^{\circ}55'24''$	0.170	0.003	0.430	0.006	ISO1631434-245524	$_{ m nII}$
201*	$16^{\rm h}31^{\rm m}48\stackrel{\rm s}{.}7$	$-24^{\circ}54'32''$	0.016	0.002	0.011	0.004	ISO1631487 - 245432	nII
202^{*}	$16^{\rm h}31^{\rm m}51\stackrel{\rm s}{.}8$	$-24^{\circ}57'26''$	0.003	0.001	0.011	0.005	ISO1631518 - 245726	nII
203*	$16^{\rm h}31^{\rm m}52^{\rm s}.0$	$-24^{\circ}55'36''$	0.080	0.002	0.322	0.004	ISO1631520-245536	nII
204	$16^{\rm h}31^{\rm m}52^{\rm s}.0$	$-24^{\circ}56'18''$	1.19	0.011	1.84	0.016	L1689-IRS5	II
205^{*}	$16^{\rm h}31^{\rm m}53\stackrel{\rm s}{.}1$	$-24^{\circ}55'04''$	0.073	0.001	0.076	0.002	ISO1631531-245504	nII
206^{*}	$16^{\rm h}31^{\rm m}54\stackrel{\rm s}{.}8$	$-25^{\circ}03'50''$	0.055	0.002	0.045	0.004	ISO1631548-250350	nII
207	$16^{\rm h}31^{\rm m}55\stackrel{\rm s}{.}2$	$-25^{\circ}03'24''$	0.355	0.005	0.582	0.006	IRAS16289 - 2457	$_{ m nII}$
208*	$16^{\rm h}31^{\rm m}59\stackrel{\rm s}{.}1$	$-24^{\circ}54'42''$	0.023	0.002	0.026	0.002	ISO1631591-245442	nII
209	$16^{\rm h}32^{\rm m}01\stackrel{\rm s}{.}0$	$-24^{\circ}56'44''$	0.982	0.019	1.17	0.011	IRS67/L1689-IRS6	I
210*	$16^{\rm h}32^{\rm m}02^{\rm s}.1$	$-24^{\circ}56'16''$	0.012	0.003	0.031	0.003	ISO1632021-245616	nII
211*	$16^{\rm h}32^{\rm m}05.6$	$-25^{\circ}02'34''$	0.019	0.002	0.031	0.003	ISO1632056-250234	nII
212	$16^{\rm h}32^{\rm m}21^{\rm s}.2$	$-24^{\circ}30'35''$	0.489	0.005	0.807	0.004	L1689-IRS7	II

^a The absolute ISOCAM coordinates have been determined using the Barsony et al. (1997) database as a reference frame. The resulting positional uncertainty is estimated to be $\sim 3''$ (rms).

The quoted rms uncertainties on $F_{\nu}^{6.7}$ and $F_{\nu}^{14.3}$ comprise the random errors estimated from the temporal and spatial noises measured on the data. They do not include the systematic errors due to absolute calibration and aperture correction uncertainties. The maximum systematic error is estimated to be less than $\sim 15\%$.

^c For a more complete cross-identification of the ρ Ophiuchi IR sources, see, e.g., Table 1 of AM94 and Table 3 of Barsony et al. (1997). "B" refers to Barsony et al. (1997). "ISO" refers to the present survey. Twelve ISO sources are completely new sources with respect to published IR surveys; they are named after their ISOCAM coordinates and marked by a * .

^d Adopted SED classes for the ISOCAM sources (see text in Sect. 3). "nI" and "nII" indicate newly identified Class I and Class II YSOs; a question mark indicates an uncertain classification.

^e ISO16=SR3, ISO48=S1, ISO90=WL22, and ISO92=WL16 are extended sources in the mid-IR range, most probably due to PAH-like emission. No attempt has been made to estimate their integrated $F_{\nu}^{6.7}$ and $F_{\nu}^{14.3}$ fluxes.

No precise mid-IR position could be derived for ISO48=S1/GY70. The quoted coordinates correspond to the near-IR position of Barsony et al. (1997).

The position of ISO53 is offset by $\sim 16''$ to the south of the near-IR source GY84. This large mid-IR offset is most likely due to the presence of extended, structured emission from the bright neighboring source S1.

^h ISO88 = SR24 is a binary system (SR24S/GY167 and SR24N/GY168; e.g. Greene et al. 1994) which is not resolved in the ISOCAM images.

Table 2. Class I YSOs

ISO #a	Identification	$\alpha_{ m IR}^{2-14}$	${L_{ m bol}}^b \ [{ m L}_{\odot}]$
182	IRS54/GY378	1.76	6.6
143	IRS44/GY269	1.57	8.7
137	CRBR85	1.48	0.36
99	LFAM26/GY197	1.25	0.064
29	GSS30/GY6	1.20	21.
31	LFAM1	1.08	0.13
65	WL12/GY111	1.04	2.6
108	EL29/GY214	0.98	26.
141	IRS43/GY265	0.98	6.7
145	IRS46/GY274	0.94	0.62
21	CRBR12	0.91	0.42
209	IRS67/L1689-IRS6	0.74	1.5
54	GY91/CRBR42	0.70	0.17
134	WL6/GY254	0.59	1.7
159	$IRS48/GY304^c$	0.18	7.4
167	$IRS51/GY315^c$	-0.04	1.1

^a The ISO number refers to the numbering of Table 1(available only in electronic form at http://cdsweb.u-strasbg.fr/).

Table 3. Class II YSOs

ISO #a	Identification	$\alpha_{ m IR}^{2-14}$	M_J [mag]	M_H [mag]	A_V [mag]	L_{\star} [L $_{\odot}$]	$L_{ m disk} \ [{ m L}_{\odot}]$
170	B162741-244645	0.51	-	6.1	24.9	0.018	0.047
103	WL17/GY205	0.42	5.2	_	22.5	0.12	0.76
124	IRS37/GY244	0.35	-	2.4	36.9	0.99	0.50
112	GY224	0.34	-	2.4	36.2	1.1	0.56
118	IRS33/GY236	0.32	-	3.6	38.3	0.28	0.15
33	GY11	0.31	10.1	-	2.7	0.001	0.010
119	IRS35/GY238	0.30	-	3.4	45.7	0.34	0.11
129	WL3/GY249	0.23	-	2.2	42.2	1.3	0.34
75	GY144	0.20	-	5.9	26.8	0.023	0.030
147	IRS47/GY279	0.17	2.6	-	26.8	1.9	1.8
46	VSSG27/GY51	0.17	5.2	-	21.6	0.11	0.30
127	GY245	0.17	6.7^{b}	-	24.7	0.023	0.11
161	GY301	0.12	-	2.1	44.9	1.5	0.25
132	IRS42/GY252	0.08	2.1	-	27.7	3.1	2.5
77	GY152	0.05	-	-	-	0.037^{c}	0.015
70	WL2/GY128	0.05	-	2.3	38.6	1.1	0.27
165	GY312	0.03	6.6	-	14.8	0.027	0.064
85	CRBR51	0.03	-	-	-	0.025^{c}	0.010
175	GY344	0.02	6.5	-	17.4	0.030	0.074
26	CRBR15	0.01	6.7	-	14.6	0.022	0.061
139	GY260	-0.03	-	3.7	40.3	0.24	0.058
37	LFAM3/GY21	-0.06	4.5	-	14.5	0.25	0.33
121	WL20/GY240	-0.07	3.3	-	16.5	0.86	0.67
51	B162636 - 241554	-0.09	4.4	-	7.6	0.28	0.43
95	WL1/GY192	-0.11	4.9	-	20.9	0.15	0.16
122	IRS36/GY241	-0.11	-	4.9	34.3	0.066	0.025

^b L_{bol} is estimated by integrating under the observed SED from 1.2 μ m to 60 μ m or 100 μ m and extrapolating from 60 μ m or 100 μ m to 200 μ m with a spectral index of $\alpha_{\text{IR}} = -1.0$. For the 9 weak sources (ISO # 137, 99, 31, 65, 146, 21, 209, 54 and 133) without reliable IRAS fluxes, L_{bol} is estimated using a typical $L_{\text{bol}}/L_{\text{cal}}$ (6.7 - 14.3 μ m) ratio of 9.8 (see Sect. 4.3).

¹³³⁾ without reliable IRAS fluxes, $L_{\rm bol}$ is estimated using a typical $L_{\rm bol}/L_{\rm cal}(6.7-14.3\mu{\rm m})$ ratio of 9.8 (see Sect. 4.3).

^c Although ISO159 = IRS48 and ISO167 = IRS51 formally lie below our practical Class I-Class II limit ($\alpha_{\rm IR}^{2-14}=0.55$), they are still considered as Class I YSOs here (cf. WLY89 and AM94).

171	GY323	-0.12		4.0	30.9	0.19	0.070
107	GY213		- 5.6			0.19 0.075	
76	GY146	-0.15		- 4.1	$19.4 \\ 43.5$	0.075 0.17	0.089
	IRS34/GY239	-0.16	-				0.025
120	IRS45/GY273	-0.23		2.4	34.8	1.1	0.19
144		-0.24	3.6	-	19.5	0.66	0.45
204	$L1689-IRS5^d$	-0.25	2.6	-	12.8	1.9	1.4
17	GSS26	-0.30	3.2	-	22.6	0.95	0.43
93	GY188	-0.36	6.3	-	20.5	0.034	0.020
98	GY195	-0.36	5.5	-	20.6	0.087	0.063
53	GY84	-0.39	5.8	-	14.1	0.060	0.037
190	GY450	-0.39	9.2	-	8.6	0.002	0.008
23	SKS1-10	-0.41	7.2	-	8.2	0.013	0.018
13	B162607-242725	-0.41	4.6	-	19.5	0.22	0.12
117	GY235	-0.43	4.9	-	9.9	0.16	0.12
212	$L1689-IRS7^d$	-0.44	2.6	-	14.7	1.8	0.64
79	GY154	-0.44	-	5.5	24.3	0.033	0.013
24	VSSG1	-0.49	3.2	-	17.1	0.97	0.49
3	IRS3	-0.50	4.3	-	4.6	0.31	0.28
39	S2/GY23	-0.51	2.0	-	11.8	3.7	1.6
140	GY262	-0.52	3.5	-	23.7	0.71	0.20
40	EL24	-0.54	1.8	-	10.0	4.5	2.1
52	VSSG4/GY81	-0.54	5.2	-	17.6	0.11	0.058
154	GY291	-0.60	4.3	-	23.0	0.29	0.058
94	B162703 - 242007	-0.61	-	6.6	16.6	0.010	0.006
128	WL4/GY247	-0.67	3.2	-	19.5	1.1	0.24
59	WL7/GY98	-0.69	4.1	-	27.2	0.37	0.061
84	WL21/GY164	-0.70	7.8	-	14.2	0.007	0.008
67	GSS39/GY116	-0.72	3.1	-	16.3	1.1	0.25
41	GY29	-0.73	5.1	-	19.1	0.13	0.043
88	SR24N/GY168	-0.74	2.5	_	7.7	2.1	0.76
35	GY15	-0.75	6.2	_	11.5	0.040	0.019
164	GY310	-0.75	6.4	_	4.0	0.032	0.024
63	GY109	-0.77	5.8	_	14.6	0.064	0.022
151	GY284	-0.77	4.9	_	7.6	0.15	0.065
43	GY33	-0.78	4.2	_	15.5	0.36	0.092
138	B162726-241925	-0.78	7.9	_	11.2	0.006	0.004
36	GSS31/GY20	-0.79	1.5	_	6.1	5.9	1.6
177	GY352	-0.79	5.2	_	17.7	0.11	0.031
197	B162821-244246	-0.81	6.0	_	20.3	0.050	0.009
110	SR21/VSSG23	-0.81	1.9	_	3.5	4.0	1.7
166	GY314	-0.83	3.4	_	6.4	0.80	0.26
9	SKS1-4	-0.85	6.0	_	9.5	0.051	0.022
12	B162604-241753	-0.86	6.6	_	13.4	0.027	0.009
155	GY292	-0.90	2.7	_	10.8	1.6	0.37
88	SR24S/GY167	-0.90 -0.91	$\frac{2.7}{2.5}$	_	5.9	$\frac{1.0}{2.2}$	0.72
19	GSS29	-0.91 -0.91	$\frac{2.5}{2.9}$	-	9.4	1.4	0.72
115	WL11/GY229	-0.92	6.3	-	13.8	0.037	0.015
196	$WSB60^e$	-0.92	4.9	-	2.8	0.16	0.075
30	GY5	-0.92	6.3	-	$\frac{2.5}{6.4}$	0.036	0.019
176	GY350	-0.94	6.4	-	6.4	0.033	0.014
72	WL18/GY129	-0.94	4.7	-	10.4	0.19	0.061
163	IRS49/GY309	-0.96	3.1	-	10.1	1.1	0.23
193	B162812-241138	-0.99	6.2	-	6.2	0.039	0.016
2	B162538-242238	-1.00	4.6	-	10.9	0.22	0.063
78	VSSG5/GY153	-1.02	3.8	-	19.7	0.55	0.056
86	IRS26/GY171	-1.04	5.2	-	19.9	0.12	0.015
105	WL10/GY211	-1.05	3.4	-	12.5	0.78	0.12
87	B162658-241836	-1.06	6.0	-	14.2	0.050	0.011
160	B162737-241756	-1.08	7.2	-	4.5	0.014	0.006
83	B162656-241353	-1.08	4.2	-	11.1	0.33	0.058

32	GY3	-1.09	6.5	-	0.4	0.029	0.015
1	IRS2	-1.09	3.4	-	4.9	0.79	0.15
185	GY397	-1.10	6.1	-	4.1	0.045	0.016
6	SR4/IRS12	-1.12	2.8	-	1.9	1.5	0.37
142	VSSG25/GY267	-1.13	4.0	-	9.8	0.42	0.065
20	DoAr24/GSS28	-1.14	3.5	-	1.8	0.72	0.16
178	GY371	-1.16	5.2	-	7.1	0.11	0.026
62	GSS37/GY110	-1.19	2.7	-	8.5	1.6	0.18
89	WL14/GY172	-1.24	5.8	-	17.2	0.059	0.006
194	B162813 - 243249	-1.24	5.2	-	5.9	0.12	0.024
116	B162713 - 241818	-1.27	3.7	-	10.7	0.57	0.046
106	B162708 - 241204	-1.33	4.5	-	8.0	0.24	0.027
102	GY204	-1.33	6.2	-	1.5	0.039	0.011
168	SR9/IRS52	-1.35	2.7	-	-0.7	1.6	0.23
199	SR13	-1.37	3.4	-	-0.9	0.78	0.12
187	SR10/GY400	-1.39	4.1	-	-3.2	0.38	0.052
68	VSS27	-1.53	2.6	-	5.6	1.9	0.13
172	GY326	-1.56	5.3	-	8.5	0.11	0.008
56	WSB37/GY93	-1.58	4.7	-	1.2	0.20	0.024
38	DoAr25/GY17	-1.58	3.4	-	0.7	0.83	0.11
-	GY256	-	-	4.5	34.5	0.11	-
-	GY257	-	-	4.1	33.2	0.17	-
90	WL22/GY174	-	-	-	-	$29.^{f}$	-
92	WL16/GY182	-	-	-	-	$44.^{f}$	-
207	IRAS16289 - 2457	-	-	-	-	1.3^{c}	0.51
123	New^g	-	-	-	-	0.077^{c}	0.031
150	New^g	-	-	-	-	0.21^{c}	0.083
195	New^g	-	-	-	-	0.14^{c}	0.058
200	New^g	-	-	-	-	0.95^{c}	0.38
201	New^g	-	-	-	-	0.025^{c}	0.010
202	New^g	-	-	-	-	0.024^{c}	0.010
203	New^g	-	-	-	-	0.71^{c}	0.28
205	New^g	-	-	-	-	0.17^{c}	0.067
206	New^g	-	-	-	-	0.099^{c}	0.040
208	New^g	-	-	-	-	0.058^{c}	0.023
210	New^g	-	-	-	-	0.067^{c}	0.027
211	New^g	-	-	-	-	0.068^{c}	0.027

The sample of 21 Class II sources with $-0.05 \le \alpha_{\rm IR}^{2-14} \le 0.55$ might contain a significant population of transition objects (flat-spectrum objects) between Class I protostars and Class II T Tauri stars.

^a The ISO number refers to the numbering in Table 1 (available only in electronic form at http://cdsweb.u-strasbg.fr/).

^b The J-band flux of ISO127 = GY245 is taken from Greene et al. (1994).

ISO77 = GY152, ISO85 = CRBR51, and the last 13 sources are not detected in all near-IR bands, and no reliable M_J or M_H can be estimated. L_{\star} has been obtained using Eq. 8 (Sect. 4.2).

^d ISO204 = L1689-IRS5 and ISO212 = L1689-IRS7 refer to the IR sources listed by Greene et al. (1994) in L1689.

 $[^]e$ ISO196 = WSB60 corresponds to the source B162816 – 243657 in Barsony et al. (1997).

Since ISO90 = WL22 and ISO92 = WL16 are young early-type stars (see Sect. 3.1), L_{\star} cannot be derived accurately using the method described in Sect. 4.1. The quoted luminosities for WL22 and WL16 are taken from WLY89 and Comerón et al. (1993), respectively. They have been scaled to $d = 140 \,\mathrm{pc}$.

 $[^]g$ The adopted names and the J2000 coordinates of the completely new IR sources are given in Table 1.

Table 4. Class III YSOs

ISO #a	Identification	$\begin{array}{c} \alpha_{\rm IR}^{2-14} \\ (\alpha_{\rm IR}^{2-7}) \end{array}$	M_J	M_H	A_V	L_{\star}
#		$(\alpha_{ m IR}^{2-7})$	[mag]	[mag]	[mag]	$[\mathrm{L}_{\odot}]$
114	$\mathrm{WL}19/\mathrm{GY}227^{b}$	-0.05	-	-1.0	73.4	52.
125	$WL5/GY246^b$	-1.02	-	-0.7	59.8	39.
34	$GY12^b$	-1.06	3.3	-	19.5	0.89
58	$WL8/GY96^b$	-1.11	1.9	-	35.5	4.2
152	$GY289^b$	-1.33	3.3	-	27.5	0.90
198	SR20	-1.65	1.9	-	4.0	4.0
133	$\text{GY}253^b$	-1.73	3.3	-	31.2	0.85
45	$LFAM8/SKS1-19^b$	-1.82	4.2	-	25.2	0.34
80	$GY156^b$	-2.00	3.4	-	22.5	0.84
10	$DoAr21/GSS23^b$	-2.00	0.7	-	6.0	15.
27	WSB28	-2.06	4.1	-	4.3	0.36
149	B162730 - 244726	-2.20	3.7	-	10.3	0.61
135	VSSG22	-2.23	3.0	-	17.1	1.2
64	$VSSG11^b$	-2.24	3.4	-	15.2	0.83
73	VSSG3/GY135	-2.26	2.3	-	15.7	2.8
180	VSSG14/GY372	-2.51	2.2	-	5.5	3.0
184	IRS55/GY380	-2.65	2.6	-	6.0	1.8
11	$\mathrm{VSSG}19^b$	-2.75	3.8	-	3.9	0.50
7	GSS20	-2.84	3.1	-	4.7	1.1
60	$GY101^b$	(0.00)	-	2.0	55.5	2.0
61	$GY103^b$	(-0.45)	-	2.6	48.5	1.1
126	$GY248^b$	(-0.82)	4.2	-	25.5	0.36
101	$IRS30/GY203^b$	(-0.99)	-	2.2	36.5	1.6
14	B162607 - 242742	(-1.43)	3.5	-	20.6	0.70
8	B162601 - 242945	(-1.96)	3.9	-	8.1	0.47
183	$GY377^b$	(-1.98)	4.0	-	16.0	0.40
157	$GY296^b$	(-2.02)	5.5	-	5.1	0.081
4	B162541 - 242138	(-2.05)	5.8	-	6.7	0.060
69	GY122	(-2.17)	5.5	-	2.7	0.080
96	GY193	(-2.24)	4.2	-	7.4	0.34
97	GY194	(-2.35)	4.1	-	9.1	0.40
188	GY410	(-2.36)	4.0	-	10.2	0.43
66	GY112	(-2.42)	4.2	-	3.6	0.35
130	$SR12/GY250^b$	(-2.48)	3.4	-	1.2	0.83
5	IRS10	(-2.78)	2.7	-	5.4	1.7
-	IRS50/GY306	_	3.7	-	11.5	0.60
16	SR3/GSS25	_	-	-	-	$100.^{c}$
48	$\mathrm{S1/GY70}^b$	_	-	-	-	1100. ^c

^a The ISO number refers to the numbering in Table 1 (available only in electronic form at http://cdsweb.u-strasbg.fr/).

^b Class III YSOs located inside the CS contours of Fig. 1 (see Sect. 3.5 and 4.4).

^c For the two B stars ISO16 = SR3 and ISO48 = S1 (e.g. Elias 1978), L_{\star} cannot be derived using the method described in Sect. 4.1. The quoted values are taken from Lada & Wilking (1984), and have been scaled to $d = 140 \,\mathrm{pc}$.

Table 5. Class III candidates located within the CS contours of Fig. 1

$\#^a$			r 1	r i	A_V	L_{\star}
		$\begin{array}{c} \alpha_{\rm IR}^{2-14} \\ (\alpha_{\rm IR}^{2-7}) \end{array}$	[mag]	[mag]	[mag]	$[L_{\odot}]$
191	GY463	-1.46	4.4	-	25.2	0.28
44	B162628-241543	-1.63	3.9	-	21.7	0.45
81	VSSG7/GY157	-2.13	2.1	-	29.8	3.4
91	VSSG8/GY181	-2.20	2.4	-	22.9	2.4
28	B162621-241544	-2.23	2.9	-	15.5	1.4
	SKS1-7	-2.38	3.6	-	17.4	0.64
25	CRBR17	(0.68)	8.3	-	10.1	0.004
153	GY290	(0.41)	-	3.7	41.5	0.31
100	$B162705 - 244013^b$	(0.12)	-	-	-	-
109	$GY215^b$	(0.05)	-	-	-	-
104	GY207	(-0.31)	-	4.6	39.5	0.11
49	$B162636 - 241811^b$	(-0.36)	-	-	-	-
15	CRBR4	(-0.41)	-	1.9	58.9	2.2
146	GY278	(-0.58)	-	2.4	47.8	1.3
136	GY258	(-0.72)	8.2	-	10.8	0.004
55	IRS16/GY92	(-0.80)	-	5.0	29.1	0.079
173	IRS53/GY334	(-0.81)	-	2.0	40.2	2.1
71	GY130	(-0.87)	-	3.7	38.8	0.31
162	GY309	(-1.01)	-	3.3	42.5	0.47
50	B162636-241902	(-1.06)	6.6	-	14.0	0.025
57	B162641-241801	(-1.22)	-	3.1	32.3	0.58
111	WL9/GY220	(-1.27)	5.5	-	21.6	0.086
22	B162618-241712	(-1.50)	4.2	-	26.8	0.34
113	IRS32/GY228	(-1.52)	3.6	-	18.5	0.62
	GY163	(-1.53)	-	4.1	32.8	0.19
131	GY255	(-1.57)	4.2	-	21.6	0.35
169	GY322	(-1.59)	4.0	-	16.3	0.43
47	IRS14/GY54	(-1.85)	4.7	-	16.6	0.21
	GY412	(-1.90)	5.2	-	17.6	0.12
181	GY373	(-1.91)	6.4	-	7.7	0.032
174	GY346	(-1.93)	5.6	-	19.5	0.075
74	IRS20/GY143	(-1.97)	3.8	-	16.0	0.50
179	GY370	(-1.99)	5.4	-	10.0	0.094
148	GY283	(-2.09)	4.3	-	13.1	0.32
42	VSSG29/GY37	(-2.14)	6.8	-	6.5	0.021
192	GY472	(-2.14)	5.9	-	20.1	0.054
	GY398	(-2.20)	5.2	-	5.2	0.11
	GY297	(-2.22)	6.0	-	0.7	0.047
156	GY295	(-2.52)	4.3	-	4.8	0.32

 $[^]a$ The ISO number refers to the numbering in Table 1 (available only in electronic form at http://cdsweb.u-strasbg.fr/).

^b A_V and L_{\star} were not derived for these sources as they were detected only in the K band by Barsony et al. (1997).