



Adult Tachycardia

NARROW (≤ 0.11 sec)



History


- Age
- Past medical history (MI, Angina, Diabetes, post menopausal)
- Recent physical exertion
- Palpitations, irregular heart beat
- Time (onset /duration / repetition)

Signs and Symptoms

- Chest pain, heart failure, dyspnea
- AMS
- Shock, poor perfusion, hypotension
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness

Differential

- Trauma vs. Medical
- Sinus Tachycardia vs. dysrhythmia
- Fever, sepsis, infection
- Pericarditis, pulmonary embolism
- Aortic dissection or aneurysm
- Overdose: Stimulants





Adult Tachycardia

NARROW (≤ 0.11 sec)



ECG and rhythm information should be interpreted in context of the entire patient assessment:

For example, if you have a patient which is ill from a likely infection with fever and is tachycardic to 140 – 160 there overall symptoms are unlikely related to tachycardia and more likely related to overwhelming sepsis and potentially hypoxia.

- Tachycardia is defined as heart rate > 100 but rarely causes symptoms unless > 120 in the adult.
- The most important decision point in care is whether the patient is stable or unstable.

Rate controlled:

- Heart rate is considered controlled when rate is ≤ 120 beats per minute.

DRUG SHORTAGE STRATEGIES:

If DILTIZEM IS NOT AVIALBLE:

- Amiodarone 150 mg in 100 mL of D5W IV / IO and infuse over 10 minutes then start infusion of Amiodarone 450 mg in 250 mL of D5W at 1 mg/min (33 mL/hr) IV / IO.
- Or Metoprolol 2.5 mg IV / IO Slow IV Push and repeat every 5 minutes as needed to a Maximum 15 mg.
- Preference Amiodarone $>$ referred Metoprolol
- DO NOT give Metoprolol to a patient with a CHF worsening.

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Most important goal is to differentiate the type of tachycardia and if STABLE or UNSTABLE and SYMPTOMATIC.**
- **12-Lead ECG:**
 - 12 Lead ECG not necessary to diagnose and treat
 - Obtain when patient is stable and/or following rhythm conversion.
- **Unstable condition**
 - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
 - If at any point patient becomes unstable move to unstable arm in algorithm.
- Search for underlying cause of tachycardia such as fever, sepsis, dyspnea, etc.
- Typical sinus tachycardia is in the range of 100 to (200 - patient's age) beats per minute.
- **Symptomatic condition**
 - Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
 - Symptomatic tachycardia usually occurs at rates ≥ 150 beats per minute.
 - Patients symptomatic with heart rates < 150 likely have impaired cardiac function such as CHF.
- **Serious Signs / Symptoms:**
 - Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute CHF.
- **If patient has history or 12 Lead ECG reveals Wolfe Parkinson White (WPW):**
 - DO NOT administer a Calcium Channel Blocker (e.g. Diltiazem) or Beta Blockers.
 - Use caution with Adenosine and give only with defibrillator available.
- **Regular Narrow-Complex Tachycardia:**
 - Vagal maneuvers and adenosine are preferred. Vagal maneuvers may convert 19% to 54 % of SVT.
 - Using passive leg raise with Valsalva is more effective.
 - Adenosine should be pushed rapidly via proximal IV site followed by 20 mL Normal Saline rapid flush.
 - Adenosine should not be used in the post-cardiac transplant patient without **Contact of Medical Control**.
 - Agencies using both calcium channel blockers and beta blockers should choose one primarily. Giving the agents sequentially requires **Contact of Medical Control**. This may lead to profound bradycardia / hypotension.
- **Irregular Narrow-Complex Tachycardia:**
 - Rate control is more important in pre-hospital setting rather than focus on rhythm conversion.
- **Synchronized Cardioversion:**
 - Recommended to treat UNSTABLE Atrial Fibrillation, Atrial Flutter and SVT.
- Monitor for hypotension after administration of Calcium Channel Blockers or Beta Blockers.
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.