

Where You Can Park — Instructor Notes

Class 13 · replacement, convenience sampling, bias vs. precision · 20 min · handout + skeleton notebook

The idea. How a sample is drawn matters more than how big it is. Part 1 shows it with one keyword: `replace=True` almost halves the chance of winning the marble contest. Part 2 shows it with geography. A crew that samples only where it can park collects three times the data and gets estimates twice as tight, and it is wrong nearly every time, because in this stream access and chloride are linked through the road bridges.

What students produce. Two simulated win probabilities, a prediction and a comparison of two sampling designs, and one discussion answer. **What this does not claim to teach:** stratified or systematic designs, the finite-population correction, or anything about real chloride in a real creek (the stream is simulated, and the handout says so).

Materials. Handout, skeleton notebook, `data/stream_sim.csv`. The notebook reads from `data/`, so upload it that way on the hub.

Timing (20 min)

Segment	Min	What happens
Part 1	5	Predict, simulate both ways, 1.3.
Part 2	11	Truth, predict, convenience loop, comparison table, 2.4.
Discussion	4	<i>Literary Digest</i> , whole class.

Watch out. Part 2 is the core. If a team hasn't reached the convenience loop by minute 9, give them the sampling line from the Appendix so they get to the table and 2.4. Part 1 can shrink to a show of hands on 1.3 if the room is slow to start.

Note. Cut for time (October 2026). The full 50-minute version also had the tank callback (1.4), three hand-drawn random samples, the careless-repeat count with `creek.sample(12)`, Part 3 (the two-humped histogram and the box plot), the streetlight question and the field-dataset question. It's in git history (commit `f22088e`) if you want to reuse any of it.

Handout and notebook question numbers now match.

The numbers

All computed in the local `ds313` environment from `stream_sim.csv` (240 reaches).

Marble contest, exact	with replacement $6/27 = 0.222$; without $48/120 = 0.400$
True mean chloride	90.2 mg/L (population SD 36.9, range 18.4–189.4)
Access	87 easy, 153 hard (36% easy); easy mean 111.6, hard mean 78.0

	centre	SD	within 10 of truth
random, n = 12	90.2	10.3	0.68
convenience, n = 40	111.6	5.1	0.013
convenience, n = 40, with replacement	111.5	7.0	0.05

One run of 1000 each; another run gave 90.1 / 10.3 / 0.667 and 111.6 / 4.9 / 0.006. Theory agrees: the SE for random $n = 12$ is 10.4; for convenience $n = 40$ drawn without replacement from 87 reaches, 5.0. The bias is $111.6 - 90.2 = 21.4$ mg/L, about twice the random design's SD. The third row is what a team gets if it forgets `with_replacement=False` in the convenience loop.

Part 1. Two Words That Change the Answer

(≈ 5 min)

1.1 Anything, committed. Guesses tend to cluster around 1/3 or “about 1/4.”

1.2 Replaced about 0.22; kept out about 0.40 (one run: 0.2225 / 0.3956). The notebook blank is `three_different_colors(False)`.

1.3 KEPT OUT: you reach into a bag and keep what you draw. `np.random.choice` samples *with* replacement by default (`replace=True`), and has all semester. The default cut the answer from 0.40 to 0.22, almost in half. Why: without replacement, once a red is drawn only one red is left among five marbles, so a second red is less likely. With replacement the bag never “remembers.” Exact: $(6 \cdot 4 \cdot 2) / (6 \cdot 5 \cdot 4) = 2/5$ vs. $3!/3^3 = 6/27$.

Note. Dice and coins are *supposed* to be sampled with replacement, since a die can't run out of sixes. That's why the default never mattered before. It bites the moment the population is a set of physical things.

Part 2. A Stream With No Name

(≈ 11 min)

2.1 90.2 mg/L (90.15).

2.2 TOO HIGH: access means roads, roads cross at bridges, and bridges are where road salt enters. Accept any answer with a mechanism. TOO LOW (“parks are cleaner”) is a reasonable prediction and worth hearing; the data says otherwise.

2.3 See *The numbers*. Random: centre about 90, SD about 10, within 10 of truth about 0.68. Convenience: centre about 112, SD about 5, within 10 of truth about 0.01.

Watch out. Convenience-loop bugs, by symptom.

- Centre about 90 instead of 112: the sample was drawn from creek, not easy.
- SD about 7 instead of 5: `easy.sample(40)` without `with_replacement=False`. That's Part 1 again; point it out rather than just fixing it.
- 2000 estimates or a column-length mismatch: the estimates were appended to `random_estimates`. The notebook's check cell names this.

