

Math 243 Analysis 2 Assignment 6

Jonathan Pearce, 260672004

February 21, 2017

Problem 2a. Derivative at 0:

$$\lim_{x \rightarrow 0} \frac{F(x) - F(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{x^2 \cos(\frac{1}{x^2}) - 0}{x} = \lim_{x \rightarrow 0} x \cos(\frac{1}{x^2})$$

Since $|\cos(\frac{1}{x^2})| \leq 1$. It follows,

$$|x \cos(\frac{1}{x^2})| \leq |x|$$

hence,

$$\lim_{x \rightarrow 0} -x \leq \lim_{x \rightarrow 0} x \cos(\frac{1}{x^2}) \leq \lim_{x \rightarrow 0} x$$

By the Squeeze Theorem,

$$\begin{aligned} \lim_{x \rightarrow 0} -x = 0 = \lim_{x \rightarrow 0} x &\implies \lim_{x \rightarrow 0} x \cos(\frac{1}{x^2}) = 0 \\ &\implies F'(0) = 0 \end{aligned}$$

Let $c \in \mathbb{R} \setminus \{0\}$. Derivative at c : Differentiate F at $\mathbb{R} \setminus \{0\}$. There exists $g : \mathbb{R} \rightarrow \mathbb{R}$ and $h : \mathbb{R} \rightarrow \mathbb{R}$ such that $(g \cdot h)(x) = F(x)$. Suppose,

$$g(x) = x^2 \text{ and } h(x) = \begin{cases} \cos(\frac{1}{x^2}), & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

Therefore by the product rule, $F'(x) = (g \cdot h)'(x) = g(x)h'(x) + h(x)g'(x)$
Differentiate g at $c \in \mathbb{R}$:

$$\lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c} = \lim_{x \rightarrow c} \frac{(x - c)(x + c)}{x - c} = \lim_{x \rightarrow c} (x + c) = 2c$$

Therefore $g'(x) = 2x, \forall x \in \mathbb{R}$

h can be expressed as the composition of two functions $\varphi \circ \psi$ where $\varphi = \cos(x)$ and $\psi = \frac{1}{x^2}$
Therefore $h(x) = (\varphi \circ \psi)(x) = \varphi(\psi(x))$. From the chain rule,

$$h'(x) = (\varphi \circ \psi)'(x) = \varphi'(\psi(x)) \cdot \psi'(x).$$

By definition $\varphi'(x) = -\sin(x)$.

Differentiate ψ at $c \in \mathbb{R} \setminus \{0\}$:

$$\lim_{x \rightarrow c} \frac{\frac{1}{x^2} - \frac{1}{c^2}}{x - c} = \lim_{x \rightarrow c} \frac{\frac{c^2 - x^2}{x^2 c^2}}{x - c} = \lim_{x \rightarrow c} \frac{\frac{(c-x)(c+x)}{x^2 c^2}}{x - c} = \lim_{x \rightarrow c} \frac{-(c+x)}{x^2 c^2} = \frac{-2c}{c^4} = \frac{-2}{c^3}$$

Therefore $\psi'(x) = -2x^{-3}$, $\forall x \in \mathbb{R} \setminus \{0\}$.

Therefore $h'(x) = (\varphi \circ \psi)'(x) = \varphi'(\psi(x)) \cdot \psi'(x) = -\sin(\frac{1}{x^2})(-2x^{-3}) \forall x \in \mathbb{R} \setminus \{0\}$

From the Product Rule,

$$F'(x) = (g \cdot h)'(x) = g(x)h'(x) + h(x)g'(x) = 2x \cos(\frac{1}{x^2}) + x^2(-\sin(\frac{1}{x^2}))(-2x^{-3}), \forall x \in \mathbb{R} \setminus \{0\}$$

$$\implies F'(x) = \begin{cases} 2x \cos(\frac{1}{x^2}) + \frac{2}{x} \sin(\frac{1}{x^2}), & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

Therefore F is differentiable.

Consider the sequence $x_n := \sqrt{\frac{1}{2\pi n + \frac{\pi}{2}}}$. We have,

$$\lim(x_n) = 0$$

$$\begin{aligned} F'(x_n) &= 2x_n \cos(\frac{1}{x_n^2}) + \frac{2}{x_n} \sin(\frac{1}{x_n^2}) = 2\sqrt{\frac{1}{2\pi n + \frac{\pi}{2}}} \cos(2\pi n + \frac{\pi}{2}) + \frac{2}{\sqrt{\frac{1}{2\pi n + \frac{\pi}{2}}}} \sin(2\pi n + \frac{\pi}{2}) \\ &= 2\sqrt{2\pi n + \frac{\pi}{2}} \\ \lim(F'(x_n)) &= \infty \end{aligned}$$

Therefore $F'(x)$ is unbounded on $[-1, 1]$. It follows that $F'(x)$ is not Riemann Integrable on $[-1, 1]$.

Problem 2.

$$\int_0^x f = \int_x^1 f \implies \int_0^x f = \int_0^1 f - \int_0^x f$$

Consider $x = 0$,

$$\int_0^0 f = \int_0^1 f \implies \int_0^1 f = 0$$

It follows,

$$\int_0^x f = - \int_0^x f$$

Since f is continuous on $[0, 1]$ it is Riemann Integrable on $[0, 1]$. Therefore define the function F as follows

$$\begin{aligned} F(x) &:= \int_0^x f \\ F(x) &= \int_0^x f = - \int_0^x f = -F(x) \end{aligned}$$

From the Fundamental Theorem (Second Form) we have,

$$F'(x) = f$$

$$-F'(x) = -f$$

Therefore,

$$f = -f$$

We conclude $f(x) = 0 \forall x \in [0, 1]$.