

**Due Friday, February 10**

1. Define  $T \in \mathcal{L}(\mathbb{R}^3, \mathbb{R}^2)$  in the standard bases by

$$T(x, y, z) = (x + y + z, 3x - 2z).$$

- (a) Find  $M(T)$  with respect to the standard bases for  $\mathbb{R}^3$  and  $\mathbb{R}^2$ .  
 (b) Find  $M(T)$  with respect to the standard basis for  $\mathbb{R}^3$  and the basis  $((1, 1), (1, 2))$  for  $\mathbb{R}^2$ .
2. Define  $f : \mathbb{F}^4 \rightarrow \mathbb{F}^2$  by

$$f(x_1, x_2, x_3, x_4) = (x_1 - 2x_3 + x_4, 3x_3 - 2x_4).$$

Find a basis for the kernel and prove that it is a basis. Then, use the fundamental theorem of linear maps to find a basis of the image of  $f$ .

3. Suppose that  $V$  is finite-dimensional and that the dimension of  $V$  is odd. Show that there does not exist a linear map  $T : V \rightarrow V$  such that

$$\ker(T) = \text{range}(T).$$

4. Suppose  $D \in \mathcal{L}(\mathcal{P}_3(\mathbb{R}), \mathcal{P}_2(\mathbb{R}))$  is the differentiation map defined by  $Dp = p'$ . Find a basis of  $\mathcal{P}_3(\mathbb{R})$  and a basis of  $\mathcal{P}_2(\mathbb{R})$  such that the matrix of  $D$  with respect to these bases is

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

5. (Ax 3.D.7) Suppose  $V$  and  $W$  are finite-dimensional. Let  $v \in V$ . Let

$$E = \{T \in \mathcal{L}(V, W) \mid Tv = 0\}.$$

- (a) Show that  $E$  is a subspace of  $\mathcal{L}(V, W)$ .  
 (b) Suppose  $v \neq 0$ . What is  $\dim E$ ? (Extended hint: If  $v \neq 0$ , we can extend it to a basis for  $V$ ,  $v, v_1, \dots, v_n$ . We can also fix a basis for  $W$ :  $w_1, \dots, w_m$ . It may help to think of a linear map in terms of its matrix with respect to the bases we just listed. What condition on the matrix forces  $v$  to go to 0? What is a basis for this set of matrices?)

The following exercises are not required to be handed in but might be helpful.

- (Ax 3.B.13) Suppose  $T$  is a linear map from  $\mathbb{F}^4 \rightarrow \mathbb{F}^2$  such that

$$\ker(T) = \{(x_1, x_2, x_3, x_4) \in \mathbb{F}^4 \mid x_1 = 5x_2 \text{ and } x_3 = 7x_4\}.$$

Prove that  $T$  is surjective.

- Suppose  $V$  is finite-dimensional and  $\dim V > 1$ . Prove that the set of noninvertible operators on  $V$  is not a necessarily a subspace of  $\mathcal{L}(V)$ .