

## ORIGINAL ARTICLE

# Combined Mesenchymal Stromal Cell Therapy and Extracorporeal Membrane Oxygenation in Acute Respiratory Distress Syndrome

## A Randomized Controlled Trial in Sheep

Jonathan E. Millar<sup>1,2,3</sup>, Nicole Bartnikowski<sup>1,4</sup>, Margaret R. Passmore<sup>1,2</sup>, Nchafatso G. Obonyo<sup>1,5</sup>, Maximillian V. Malfertheiner<sup>1,6</sup>, Viktor von Bahr<sup>1,7</sup>, Meredith A. Redd<sup>8</sup>, Louise See Hoe<sup>1,2</sup>, Katrina K. Ki<sup>1,2</sup>, Sanne Pedersen<sup>1,4</sup>, Andrew J. Boyle<sup>3</sup>, J. Kenneth Baillie<sup>9</sup>, Kiran Shekar<sup>1,2</sup>, Nathan Palpant<sup>8</sup>, Jacky Y. Suen<sup>1,2</sup>, Michael A. Matthay<sup>10,11</sup>, Daniel F. McAuley<sup>3</sup>, and John F. Fraser<sup>1,2</sup>; on behalf of the CELTIC Investigators

<sup>1</sup>Critical Care Research Group, the Prince Charles Hospital, Brisbane, Queensland, Australia; <sup>2</sup>Faculty of Medicine and <sup>8</sup>Institute for Molecular Bioscience, University of Queensland, Brisbane, Queensland, Australia; <sup>3</sup>Wellcome-Wolfson Institute for Experimental Medicine, Queen's University Belfast, Belfast, United Kingdom; <sup>4</sup>Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, Queensland, Australia; <sup>5</sup>Wellcome Trust Centre for Global Health Research, Imperial College London, London, United Kingdom; <sup>6</sup>Department of Internal Medicine II, Cardiology and Pneumology, University Medical Center Regensburg, Regensburg, Germany; <sup>7</sup>Section for Anesthesiology and Intensive Care Medicine, Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden; <sup>9</sup>Roslin Institute, University of Edinburgh, Edinburgh, United Kingdom; and <sup>10</sup>Department of Medicine and <sup>11</sup>Department of Anesthesia, Cardiovascular Research Institute, University of California, San Francisco, San Francisco, California

ORCID IDs: 0000-0002-4853-9377 (J.E.M.); 0000-0002-0859-8882 (M.R.P.).

### Abstract

**Rationale:** Mesenchymal stromal cell (MSC) therapy is a promising intervention for acute respiratory distress syndrome (ARDS), although trials to date have not investigated its use alongside extracorporeal membrane oxygenation (ECMO). Recent preclinical studies have suggested that combining these interventions may attenuate the efficacy of ECMO.

**Objectives:** To determine the safety and efficacy of MSC therapy in a model of ARDS and ECMO.

**Methods:** ARDS was induced in 14 sheep, after which they were established on venovenous ECMO. Subsequently, they received either endobronchial induced pluripotent stem cell–derived human MSCs (hMSCs) ( $n = 7$ ) or cell-free carrier vehicle (vehicle control;  $n = 7$ ). During ECMO, a low  $V_T$  ventilation strategy was employed in addition to protocolized hemodynamic support. Animals were monitored and supported for 24 hours. Lung tissue,

bronchoalveolar fluid, and plasma were analyzed, in addition to continuous respiratory and hemodynamic monitoring.

**Measurements and Main Results:** The administration of hMSCs did not improve oxygenation ( $\text{PaO}_2/\text{FiO}_2$ , mean difference =  $-146$  mm Hg;  $P = 0.076$ ) or pulmonary function. However, histological evidence of lung injury (lung injury score mean difference =  $-0.07$ ;  $P = 0.04$ ) and BAL IL-8 were reduced. In addition, hMSC-treated animals had a significantly lower cumulative requirement for vasopressor. Despite endobronchial administration, animals treated with hMSCs had a significant elevation in transmembrane oxygenator pressure gradients. This was accompanied by more pulmonary artery thromboses and adherent hMSCs found on explanted oxygenator fibers.

**Conclusions:** Endobronchial hMSC therapy in an ovine model of ARDS and ECMO can impair membrane oxygenator function and does not improve oxygenation. These data do not recommend the safe use of hMSCs during venovenous ECMO.

**Keywords:** ARDS; ECMO; MSC; models; animal

(Received in original form November 7, 2019; accepted in final form April 14, 2020)

Supported by the Intensive Care Society (UK), the Prince Charles Hospital Foundation, the Queensland Government, and the National Health and Medical Research Council (Australia). M.A.M. was supported by NHLBI grants HL123004, HL140026, and HL126456.

Author Contributions: J.E.M.—study conception, model development, study design and conduct, animal surgery, data collection and analyses, and manuscript preparation; N.B. and J.Y.S.—model development, study design and conduct, data collection and analyses, and manuscript review; N.G.O., M.V.M., and V.v.B.—model development, study design and conduct, animal surgery, and manuscript review; M.R.P., M.A.R., L.S.H., and K.K.K.—sample analyses, data analyses, and manuscript review; S.P.—model development, study design and conduct, animal surgery, and manuscript review; J.K.B.—data analysis and manuscript review; A.J.B., K.S., N.P., and M.A.M.—study design, data interpretation, and manuscript review; D.F.M. and J.F.F.—study conception, model development, study design, data interpretation, and manuscript review.

Correspondence and requests for reprints should be addressed to Jonathan E. Millar, M.D., Critical Care Research Group, the Prince Charles Hospital, Chermiside, Brisbane, Queensland 4032, Australia. E-mail: j.millar@doctors.org.uk.

This article has a related editorial.

This article has an online supplement, which is accessible from this issue's table of contents at [www.atsjournals.org](http://www.atsjournals.org).

Am J Respir Crit Care Med Vol 202, Iss 3, pp 383–392, Aug 1, 2020

Copyright © 2020 by the American Thoracic Society

Originally Published in Press as DOI: 10.1164/rccm.201911-2143OC on April 15, 2020

Internet address: [www.atsjournals.org](http://www.atsjournals.org)

## At a Glance Commentary

### Scientific Knowledge on the

**Subject:** Mesenchymal stromal cell (MSC) therapy is a promising novel intervention for acute respiratory distress syndrome (ARDS). Trials to date have failed to explore the safety of cell therapy during extracorporeal membrane oxygenation (ECMO). Preclinical investigations have found that MSC therapy during ECMO attenuate the efficacy of ECMO.

### What This Study Adds to the Field:

This study, employing a 24-hour large animal model of ARDS and ECMO, examined the safety and efficacy of MSC therapy. We found that endobronchially administered MSCs adhere to, and impair, membrane oxygenators *in vivo*. MSCs did not improve oxygenation or other ventilatory parameters but did reduce the severity of histological evidence of lung injury. Our data also suggest that MSCs may reduce circulatory shock associated with ARDS. There were no adverse effects of MSC administration on renal or liver function.

The quest for an effective pharmacological treatment for the acute respiratory distress syndrome (ARDS) has been unsuccessful. Recently, mesenchymal stromal cells (MSCs) have attracted attention as a candidate therapy for ARDS (1).

MSCs are multipotent adult stem cells found in tissues of mesodermal origin, such as bone marrow (2). Therapeutic interest in these cells has arisen because of their pleiotropic immunomodulatory abilities. During acute inflammation, MSCs appear to be immunosuppressive, influencing both innate and adaptive immune responses (3). In ARDS, their beneficial effects are believed to be mediated in a variety of ways, including secretion of antiinflammatory paracrine factors (4), restoration of epithelial and endothelial integrity (5), enhancement of alveolar fluid clearance (6), direct antimicrobial activity (7), and mitochondrial transfer (8). In preclinical models of acute lung injury, MSCs have been shown to reduce mortality (9). A phase 2 study has been conducted in patients with ARDS with no reported infusion-related adverse events (10).

To date, trials of MSCs in ARDS have excluded patients supported with extracorporeal membrane oxygenation (ECMO). The use of ECMO in acute severe respiratory failure has increased substantially in the last decade, and is now an established tool for supporting those with refractory illness (11). The use of MSCs during ECMO, while potentially attractive, raises some unique considerations. First, MSCs are large cells, with an average diameter between 10 and 30  $\mu\text{m}$  (12), which, when administered therapeutically, may pose a risk to the patency of a membrane oxygenator. Second, a defining characteristic of MSCs is avid plastic adherence (13); this too may threaten membrane oxygenators, which are constructed largely from plastics. Recent *ex vivo* and small animal experimentation has confirmed these concerns (14, 15). Conversely, immunomodulation by MSCs may provide additional benefits for patients on ECMO, where the institution of extracorporeal support results in an additional inflammatory insult (16).

Given the paucity of evidence to support the safe use of MSC therapy during ECMO, we conducted a controlled trial of clinical-grade induced pluripotent stem cell (iPSC)-derived human MSCs (hMSCs) in an ovine model of ARDS, supported with venovenous ECMO (VV-ECMO). The primary objective was to assess the safety of MSC therapy and to investigate its effect on physiologic and biologic markers of pulmonary and systemic injury.

## Methods

### Study Design

Ethical approvals were obtained from University Animal Ethics Committees of Queensland University of Technology and the University of Queensland (QUT1600001108 and UQPC/483/17) and authorization for *in vivo* use of hMSCs was granted by the Australian Department of Agriculture (2017/075). The study was conducted in accordance with the *Australian Code for the Care and Use of Animals for Scientific Purposes* (17) and is reported in compliance with Animal Research: Reporting of *In Vivo* Experiments guidelines (18). Detailed methods and a comprehensive description of the experiments and analyses are provided in the online supplement. A schematic of the study protocol is provided in Figure 1.

### Animal Model

A total of 14 healthy Border Leicester Cross ewes, aged between 1 and 3 years and weighing between 46 and 55 kg (mean =  $52.6 \pm 3$  kg), was randomly assigned to one of two groups: endobronchial iPSC-derived hMSC treatment ( $n = 7$ ) or endobronchial carrier vehicle only ( $n = 7$ ).

In brief, animals were anesthetized with a combination of ketamine, midazolam, and fentanyl. Continuous neuromuscular blockade was maintained by infusion of vecuronium. In the supine position, animals were tracheostomized and ventilated using a low- $V_T$  strategy (6 ml/kg actual body weight [ABW]). After instrumentation, acute lung injury was induced by combining an intravenous infusion of oleic acid (0.06 ml/kg, O1008; Sigma-Aldrich) with endobronchial *Escherichia coli* LPS (LPS, 100  $\mu\text{g}$ , O55:B5; Sigma-Aldrich). Once a  $\text{PaO}_2/\text{FiO}_2$  ratio  $< 100$  mm Hg (positive end-expiratory pressure  $\geq 10$  cm  $\text{H}_2\text{O}$ ) was obtained (Time 0 [ $T_0$ ]), animals were established on VV-ECMO via a right-sided jugular–jugular configuration ( $T_1$ ) and positioned in sternal recumbency. VV-ECMO was combined with a lower- $V_T$  strategy (4 ml/kg ABW) for 22 hours, at which time ( $T_{23}$ ) ECMO was stopped and a standardized recruitment maneuver was performed. Animals were returned to pre-ECMO ventilatory settings for 1 hour before being killed ( $T_{24}$ ).

### iPSC-derived hMSCs


After 1 hour of VV-ECMO ( $T_2$ ), animals received a fixed dose of  $3 \times 10^8$  iPSC-derived hMSCs suspended in a carrier vehicle (hMSC) or carrier vehicle alone (vehicle control). Cells were provided by Cynata Therapeutics Ltd. (CYP-001; Cynata Therapeutics Ltd.). These cells were at least 99% positive for CD-73, CD-90, and CD-105, but negative for CD-31 and CD-45. The total volume of vehicle was 60 ml (57.5% plasmalyte-A, 40% flexbumin 25%, and 2.5% DMSO). Cells were at least 97% viable before administration. The distribution of delivery is described in Figure E1 in the online supplement.

### Statistical Analysis

An *a priori* sample size calculation, based on the primary outcome of  $\text{PaO}_2/\text{FiO}_2$  ratio at 24 hours, is detailed in the online supplement. Data are expressed as mean

## Mechanical ventilation

Mode	Volume controlled
Tidal volume	6 mL/kg
Respiratory rate	≤35 breaths per minute*
FiO <sub>2</sub>	Lowest required for PaO <sub>2</sub> ≥80 mmHg <sup>†</sup>
PEEP	Start 5 cmH <sub>2</sub> O <sup>‡</sup>
Plateau pressure	≤30 cmH <sub>2</sub> O
I:E ratio	1:1 to 1:3



## VV-ECMO

ECMO settings		Mechanical ventilation	
Flow	Approx. 2/3 <sup>rd</sup> CO, adjusted to maintain PaO <sub>2</sub> ≥80 mmHg	Mode	Volume controlled
Sweep gas	Initially 80% of flow rate, adjusted to maintain PaCO <sub>2</sub> 30-35 mmHg	Tidal volume	4 mL/kg
Blender O <sub>2</sub>	100%	Respiratory rate	8 breaths per minute
Circuit T <sup>o</sup>	38 degree celsius	FiO <sub>2</sub>	0.3
<b>Anticoagulation</b>		PEEP	10 cmH <sub>2</sub> O
Target ACT 180-210 seconds		Plateau pressure	≤25 cmH <sub>2</sub> O
20 IU/kg i.v. porcine heparin at time of cannulation, then variable rate i.v. heparin infusion		I:E ratio	1:1 to 1:4

## Recruitment

Access cannula clamped, circuit volume returned, return cannula clamped
<b>Recruitment manoeuvre</b>
40 s breath hold at an inspiratory pressure of 40 cmH <sub>2</sub> O and an FiO <sub>2</sub> 1.0
Mechanical ventilation returned to pre-ECMO settings with an initial FiO <sub>2</sub> 1.0 and PEEP 10 cmH <sub>2</sub> O

**Figure 1.** Study schematic. \*Adjusted to maintain pH 7.30–7.45. Permissive hypercapnia was tolerated to a minimum pH of 7.15.

<sup>†</sup>PaO<sub>2</sub> = 55–80 mm Hg. If, despite an FiO<sub>2</sub> of 1.0, oxygenation targets were not met, positive end-expiratory pressure (PEEP) was increased, maintaining plateau pressure ≤32 cm H<sub>2</sub>O. <sup>‡</sup>Total PEEP (extrinsic PEEP + intrinsic PEEP) did not exceed 20 cm H<sub>2</sub>O. PEEP was permitted to be reduced to 5 cm H<sub>2</sub>O to maintain plateau pressure ≤30 cm H<sub>2</sub>O. If, despite a PEEP of 5 cm H<sub>2</sub>O, plateau pressure was >30 cm H<sub>2</sub>O, V<sub>T</sub> was reduced in 1-ml/kg steps until set at 4 ml/kg. ACT = activated clotting time; ECMO = extracorporeal membrane oxygenation; hMSC = human mesenchymal stromal cell; I:E = inspiratory: expiratory time ratio; OA = oleic acid; P/F = PaO<sub>2</sub>/FiO<sub>2</sub> ratio; VV = venovenous.

(±SD) or median (interquartile range [IQR]) if nonnormally distributed. Analysis was undertaken in Graphpad Prism (v 8.1.2; GraphPad Software). Longitudinal data were analyzed by fitting a mixed model. This model uses a compound symmetry covariance matrix and is fit using restricted maximum likelihood. Where a significant interaction was observed, *post hoc* comparisons were undertaken. Correction for multiple comparisons was made using the Benjamini-Hochberg method (false discovery rate restricted to 5%). Nonlongitudinal data were compared using an unpaired *t* test or a Mann-Whitney test, as appropriate. Categorical data were compared using the chi-square test. Statistical significance was assumed if the *P* value was less than 0.05.

## Results

Baseline characteristics at injury (T<sub>0</sub>) are shown in Table 1 and Table E1. All animals completed the study protocol, and were killed at T<sub>24</sub>.

## Respiratory Variables

The use of ECMO facilitated a lower-V<sub>T</sub> ventilation strategy (median [IQR], 4 [4–4] mL/kg ABW). The median (IQR) ECMO flow rate was 2.75 L/min (2.5–3.25 L/min), with a sweep gas flow of 3 L/min (2–3.5 L/min). During VV-ECMO, animals had a median (IQR) PaO<sub>2</sub> of 109 mm Hg (94–131 mm Hg) and a PaCO<sub>2</sub> of 32 mm Hg (30–35 mm Hg). There were no significant differences in these parameters between

groups (Figure E2). Animals were adequately anticoagulated during ECMO, as measured by activated partial thromboplastin time ratios. The dose of heparin was not significantly different between groups (Figure E2).

Because VV-ECMO controls gas exchange, native lung function was assessed 1 hour after cessation of extracorporeal flow and after the performance of a standardized lung recruitment maneuver (T<sub>24</sub>). As shown in Figure 2, both the PaO<sub>2</sub>/FiO<sub>2</sub> ratio (*P* = 0.076) and the oxygenation index (*P* = 0.153) were numerically better in the carrier vehicle-only group; the differences were not statistically significant.

The plateau airway and driving pressures were similar between groups at T<sub>24</sub> (Figure 2). Static lung compliance was

**Table 1.** Baseline Physiological Characteristics

Characteristics	Overall (n = 14)	Vehicle (n = 7)	hMSCs (n = 7)
Weight, kg	52.6 ± 3	52.4 ± 3.2	52.9 ± 2.6
Pre-ECMO Vt, ml/kg	6.0 ± 0.1	6.0 ± 0.1	6.0 ± 0.1
At time of injury (T <sub>0</sub> )			
Peak airway pressure, cm H <sub>2</sub> O	31.5 ± 4.2	31.9 ± 3.3	31.1 ± 4.9
Plateau pressure, cm H <sub>2</sub> O	26.5 ± 3.9	26.1 ± 3.9	26.9 ± 3.8
Driving pressure, cm H <sub>2</sub> O	16.5 ± 3.9	16.1 ± 3.9	16.9 ± 3.8
Static compliance, ml/cm H <sub>2</sub> O	21 ± 5.0	19 ± 4.6	23 ± 4.6
PaO <sub>2</sub> /F <sub>I</sub> O <sub>2</sub>	59 ± 20	58 ± 23	61 ± 17
PaCO <sub>2</sub> , mm Hg	41 (38–46)	38 (38–41)	44 (41–48)
pH	7.36 ± 0.05	7.38 ± 0.04	7.34 ± 0.05
Bicarbonate, mmol/L	23.2 ± 1.5	23.6 ± 1.4	22.9 ± 1.5
Base deficit, mmol/L	1.30 ± 1.31	0.94 ± 1.15	1.66 ± 1.37
Arterial lactate, mmol/L	1.6 ± 0.7	1.8 ± 0.6	1.4 ± 0.7
Heart rate, beats/min	102 (96–117)	108 (98–128)	98 (95–103)
Mean arterial pressure, mm Hg	103 ± 19	96 ± 16	111 ± 19
Central venous pressure, mm Hg	13 (11–13)	12 (11–13)	13 (13–14)
Mean pulmonary artery pressure, mm Hg	25 (16–29)	20 (17–27)	28 (19–29)

Definition of abbreviations: ECMO = extracorporeal membrane oxygenation; hMSCs = human mesenchymal stromal cells.

also similar, both during and after ECMO (Figure 2). The use of a protocolized recruitment maneuver did not improve compliance after cessation of extracorporeal support in either group (Figure 2).

### Hemodynamic Variables

This model of acute lung injury was associated with the development of hyperdynamic shock, which worsened over time (Figure 3). The administration of hMSCs resulted in significantly lower cumulative vasopressor doses (Figure 3). At T<sub>4</sub>, mean arterial pressure (MAP) was significantly higher in the hMSC-treated group ( $P = 0.001$ ), even though these animals received lower doses of noradrenaline (Figure 3). By T<sub>14</sub>, MAP was again similar between groups, although vasopressor requirements continued to be lower in hMSC-treated animals. In addition, there were lower arterial lactate concentrations, higher arterial base excesses, and lower mean pulmonary artery pressures from 12 hours (T<sub>14</sub>) after instillation in the hMSC group (Figure 3); however, these were not statistically significant. Cumulative fluid balance at T<sub>24</sub> was similar in both groups (vehicle control, 2,713 ± 970 ml vs. hMSCs, 2,992 ± 1,237 ml;  $P = 0.648$ ).

### Histopathology and Lung Injury

The blinded assessment of lung tissue was conducted by an independent expert veterinary pathologist. Sections of the right lower lobe were prepared, and a

lung injury score was calculated (19). The administration of hMSCs resulted in significantly lower scores ( $P = 0.04$ ; Figure 4), principally mediated by a reduction in neutrophil infiltration.

There were no significant differences in lung wet/dry ratio or BAL total protein concentration (Figure 4). BAL fluid inflammatory cell counts are detailed in Table E4. There were no significant differences in these counts over time. Similarly, there was no difference in lung tissue homogenate gene expression (as assessed by quantitative PCR) between groups (Figure E3).

In a *post hoc* analysis, pulmonary arterial thrombosis was noted in five hMSC-treated animals, but only one animal receiving carrier vehicle alone ( $P = 0.031$ ).

### Inflammatory Cytokines

BAL and plasma cytokine concentrations were assessed longitudinally (Figures E4 and E5). In BAL, statistically significant differences in IL-8 were observed at T<sub>3</sub>, T<sub>14</sub>, and T<sub>23</sub> ( $P = 0.013$ , 0.016, and 0.028, respectively). In plasma, cytokine trajectories were similar between groups (Figure E5).

### Hematological and Biochemical Measurements


A summary of hematological and biochemical values are provided in Tables E2 and E3. This lung injury model was associated with the development of acute

kidney injury and abnormal liver function, although there were no significant differences in indices between groups. The administration of hMSCs resulted in a significantly lower lymphocyte count at T<sub>24</sub> ( $P = 0.047$ ; Figure E5).

### Cell-ECMO Membrane Interaction and Cell Fate

The administration of hMSCs was associated with a significant increase in the transoxygenator pressure gradient, becoming apparent 4 hours after cell delivery (Figure 5). By T<sub>23</sub>, the mean pressure gradient in the hMSC group reached 64 (±37) mm Hg versus 17 (±9) mm Hg in the vehicle-only group. The instillation of carrier vehicle alone was associated with a reduction in the ECMO pump speed to flow ratio over time, a finding not observed in the hMSC group (Figure 5). During the study, there were no instances of pump or oxygenator failure requiring a component exchange. Likewise, there was no evidence of clotting on the oxygenator surface by visual examination in either group.

Membrane oxygenators from animals treated with hMSCs were isolated and preserved at the termination of ECMO. Subsequent deconstruction and staining of the fiber bundles ( $n = 7$ ) revealed adherent cells exhibiting surface markers consistent with those of hMSCs (Figure E6). Similar cell populations were not apparent in vehicle-only controls ( $n = 3$ ).



**Figure 2.** Oxygenation and respiratory parameters. (A)  $\text{PaO}_2/\text{FiO}_2$  ratio. (B) Oxygenation index. (C) Airway pressures and lung compliance. Data are presented as mean ( $\pm$  95% confidence interval [CI]). Where error bars intersect the x-axis, the 95% CI includes zero. hMSC = human mesenchymal stromal cell; VV-ECMO = venovenous extracorporeal membrane oxygenation.


## Discussion

We performed a trial of clinical-grade iPSC-derived hMSCs, given endobronchially, for




acute lung injury in sheep during VV-ECMO. The main findings of this study can be summarized as follows: 1) with regard to the primary outcome, hMSCs did not

improve oxygenation at 24 hours; 2) hMSCs did not improve pulmonary mechanics, but did improve the severity of histological lung injury and reduced the





**Figure 3.** Hemodynamic variables. (A) Mean arterial pressure and vasopressor dose (mean). (B) Mean pulmonary artery pressure. (C) Cumulative vasopressor dose. (D) Cardiac index, base deficit, and arterial lactate. Data are presented as mean ( $\pm$ 95% confidence interval [CI]). Where error bars intersect the x-axis, the 95% CI includes zero.  $**P < 0.01$ . hMSC = human mesenchymal stromal cell; VV-ECMO = venovenous extracorporeal membrane oxygenation.

**A****B****C**


**Figure 4.** Histopathology and lung injury. (A) Representative images of lung parenchyma. All animals showed evidence of diffuse alveolar damage; however, the frequency and degree of injury differed between groups. Panel 1: extensive alveolar edema with interstitial leukocyte infiltration. Panel 2: marked leukocyte infiltration within alveolar spaces and larger airways. Panel 3: some loss of alveolar structure with edema; however, a reduction in interstitial and alveolar leukocytes. Panel 4: preservation of alveolar architecture with few leukocytes in the alveolar spaces. Panels 5 and 6: representative images of pulmonary arterial and arteriolar emboli (black arrows) in animals receiving human mesenchymal stromal cells (hMSCs). Panels 1–4: scale bars, 100 µm. Panels 5 and 6: scale bars, 200 µm. (B) Composite lung injury score, lung wet/dry ratio (right lower lobe). (C) BAL total protein concentration. Data are presented as mean ( $\pm$ 95% confidence interval). VV-ECMO = venovenous extracorporeal membrane oxygenation.

concentration of BAL IL-8; 3) in spite of endobronchial administration, hMSCs adhered to and impacted the function of a commercial membrane oxygenator *in vivo*, with an increase in the transmembrane oxygenator pressure gradient, and more pulmonary arterial thromboses were noted in hMSC-treated lungs; and 4) hMSCs reduced the depth and severity of shock.

This study was conducted in a large animal model of ARDS and ECMO, which replicates several important clinical features (20). The “double-hit” injury applied in this study resulted in acute severe hypoxemic respiratory failure consistent with modern criteria for the use of VV-ECMO (11). To support the severe acute respiratory failure, we employed a commercial ECMO device that is in widespread clinical use. In

addition, our protocolized intensive care was consistent with clinical best-practice standards (21). A common criticism of preclinical trials of MSCs has been the use of heterogeneous, non-clinical grade cell products (1); however, we tested a commercial hMSC product that is under investigation in clinical trials.

MSCs have been administered to patients with respiratory failure on ECMO



**Figure 5.** Cell–extracorporeal membrane oxygenation (ECMO) interaction. (A) Transmembrane oxygenator pressure gradient. (B) Pump revolutions per minute/flow ratio. Data are presented as mean ( $\pm$ 95% confidence interval [CI]). Where error bars intersect the x-axis, the 95% CI includes zero.

\* $P < 0.05$ . hMSC=human mesenchymal stromal cell.

(22, 23). These reports, which include only three patients, did not describe infusion-related adverse events, and failed to fully characterize the interaction between MSCs and the extracorporeal device. Kocyildirim and colleagues (24) have conducted the only other preclinical study involving both MSCs and ECMO for respiratory failure. Their ovine-based 6-hour pilot study did not report on the impact of MSC administration on the performance of ECMO.

### hMSCs and Pulmonary Function

In this study, the administration of hMSCs failed to improve oxygenation at T<sub>24</sub>. Furthermore, animals receiving hMSCs had a trend for worse oxygenation index values. In a phase 2a study of 60 patients with ARDS, the intravenous administration of hMSCs did not significantly improve PaO<sub>2</sub>/FiO<sub>2</sub> ratio, although there was a signal toward improvement in oxygenation index in a *post hoc* analysis (10). Preclinical studies of MSCs in ARDS have reported improvements in oxygenation, although few have produced lung injury as severe (9). In a recent systematic review of preclinical models combining ARDS and ECMO, only four achieved PaO<sub>2</sub>/FiO<sub>2</sub> values <100 mm Hg (20). The degree of lung injury in this model may explain why oxygenation is impaired in the treated

group, despite improvements in inflammation and lung injury. Emerging research has highlighted the procoagulant effects of transplanted MSCs. These appear to be primarily mediated by MSC expression of tissue factor (25), but also by the secretion of procoagulant microvesicles (26) and by direct enhancement of platelet deposition (27). In preclinical experiments MSCs have been associated with the development of pulmonary emboli *in vivo* (28). In this study, despite the use of heparin, almost all hMSC animals ( $n=6$ ) had histological evidence of pulmonary arterial thrombosis postmortem. The presence of exogenous hMSCs within the disordered pulmonary vasculature may have contributed to impairments in oxygenation, tempered by the fact that there was no increase in mean pulmonary artery pressure in treated animals.

Animals receiving hMSCs had improved composite histologic lung injury scores postmortem. The components of the score most influenced by hMSCs were neutrophil numbers in the alveolar and interstitial space. Multiple studies of MSCs in preclinical models of ARDS have demonstrated their ability to reduce neutrophil infiltration (29) and neutrophil extracellular trap formation (30). In this study, BAL neutrophil counts did not differ between groups. This may reflect the

technical challenges of obtaining and assessing BAL cell counts. In a recent porcine model of ARDS and MSC therapy, a reduction in neutrophil infiltration was correlated with a reduction in BAL IL-8 concentrations (31), a finding confirmed in this study.

### hMSCs and the Systemic Inflammatory Response

Multiple preclinical models (32) and recent clinical trials (33–36) have examined the use of MSCs in the treatment of septic shock. Animals receiving hMSCs required less vasopressor support throughout the experiment to achieve an equivalent or higher MAP. A similar, early, but nonsustained reduction in vasopressor requirement has previously been described in a large-animal model of septic shock treated with MSCs (37). hMSCs did not alter plasma concentrations of proinflammatory cytokines over extended time periods in this study, a finding that has previously been identified in other preclinical (38) and clinical studies (33). A recent phase I dose-escalation study of MSCs in patients with septic shock demonstrated that the maximum effect of cell therapy on plasma cytokine levels occurred at 4 hours after administration and declined with time (35). In this study,



that time period coincides with the maximum separation in vasopressor dose, MAP, and levels of IL-1 $\beta$  and IL-6 between groups. This may indicate that repeat dosing of MSCs will be required for optimal therapeutic efficacy.

### hMSCs and ECMO

The risk posed by MSCs to membrane oxygenators has been postulated for some time (1), but has only recently been shown to have an experimental basis. Our group has previously reported the ability of hMSCs to tightly adhere to the membrane fibers of a commercial oxygenator. This may have been the result of the known plastic avidity of MSCs (13). Recently, Cho and colleagues (15) reported the loss of systemically administered MSCs in an *ex vivo* model of venoarterial ECMO.

Given the emerging signal that systemically administered MSCs may interact with membrane oxygenators, we decided to test endobronchial instillation in this study. Cardenes and colleagues (39) have used 18F-fluorodeoxyglucose labeling to track the fate of both systemically and endobronchially administered MSCs in an ovine model of ARDS. Although systemically administered cells have a wide biodistribution in the first 5 hours, endobronchially administered cells were retained at the site of instillation. There are key differences in our approach, including the means of inducing lung injury and its severity.

### Limitations

This study has some limitations. First, although our model of injury replicates several relevant features, including severe respiratory and hemodynamic failure,

clinical ARDS is usually caused by infection and develops over several days, often in patients with other comorbidities (40). Second, MSCs are known to exhibit different functional responses dependent on the contemporary milieu, which, in some circumstances, may be detrimental (41). This study may have modeled only one phase of acute lung injury, and so hMSCs may have exerted an effect that may differ in other phases. Third, although our model extended 21 hours after cell or vehicle delivery, this may have been too short a period to observe some beneficial effects of the intervention. For example, it may be that the favorable effect of hMSCs on histological injury may have translated to improvements in oxygenation over a longer time period. Conversely, an extension of the study period in the face of rising transoxygenator pressure gradients in the hMSC group may have ultimately led to circuit failures. Fourth, the use of a lung recruitment maneuver and the assessment of native lung function off ECMO may have had several adverse effects, and we cannot be certain that these effects did not differ between groups. This approach was taken due to the challenge of assessing native lung function during ECMO, particularly where a lower V<sub>T</sub> ventilatory strategy has been adopted. The study protocol was designed before the publication of the ART (Alveolar Recruitment for ARDS Trial) randomized controlled trial (42). Fifth, the addition of an uninjured control group may have provided further insights into the distribution of hMSCs during VV-ECMO. Finally, the dose and method of delivery of hMSCs remain a matter of conjecture. Based on the findings of our previous

work (14), we chose not to investigate intravenous administration. Likewise, based on clinical trial experience, we opted to administer a single, fixed dose of hMSCs. It is possible that varying the dose and/or route of administration of hMSCs may alter their efficacy and safety profile during ECMO.

### Conclusions

In a 24-hour ovine model of ARDS and VV-ECMO, we found that hMSC therapy was associated with impairment of the membrane oxygenator. The use of cell therapy did not result in improvements in oxygenation, the primary outcome of this study, but was associated with a reduction in histological evidence of lung injury and inflammation in the lung. Given these data, we cannot currently recommend the administration of hMSCs during ECMO. ■

**Author disclosures** are available with the text of this article at [www.atsjournals.org](http://www.atsjournals.org).

**Acknowledgment:** The authors are grateful for assistance provided during animal experiments by the staff of the Queensland University of Technology Medical Engineering Research Facility and the University of Queensland Animal Science Precinct. They extend their thanks to Ms. Arlanna Esguerra-Lallen for providing expert nursing care and to Ms. Mengyao Yang, Mr. Matthew Wells, Dr. Ai-Ching Boon, Mr. Michael Cavaye, and Ms. Ashlen Garrett for their assistance in study preparation, sampling, and laboratory analyses. They also acknowledge the endorsement of this study by the International ECMO Network. Mesenchymal stromal cells were donated in kind, under a material transfer agreement, by Cynata Therapeutics Ltd., Melbourne, Australia. Cynata Therapeutics Ltd. was not involved in the conception, design, or analysis of the study.

### References

1. Millar JE, Fraser JF, McAuley DF. Mesenchymal stromal cells and the acute respiratory distress syndrome (ARDS): challenges for clinical application. *Thorax* 2015;70:611–612.
2. Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, et al. Multilineage potential of adult human mesenchymal stem cells. *Science* 1999;284:143–147.
3. Curley GF, McAuley DF. Stem cells for respiratory failure. *Curr Opin Crit Care* 2015;21:42–49.
4. Devaney J, Horie S, Masterson C, Elliman S, Barry F, O'Brien T, et al. Human mesenchymal stromal cells decrease the severity of acute lung injury induced by *E. coli* in the rat. *Thorax* 2015;70:625–635.
5. Fang X, Neyrinck AP, Matthay MA, Lee JW. Allogeneic human mesenchymal stem cells restore epithelial protein permeability in cultured human alveolar type II cells by secretion of angiopoietin-1. *J Biol Chem* 2010;285:26211–26222.
6. McAuley DF, Curley GF, Hamid UI, Laffey JG, Abbott J, McKenna DH, et al. Clinical grade allogeneic human mesenchymal stem cells restore alveolar fluid clearance in human lungs rejected for transplantation. *Am J Physiol Lung Cell Mol Physiol* 2014;306:L809–L815.
7. Krasnodembskaya A, Song Y, Fang X, Gupta N, Serikov V, Lee J-W, et al. Antibacterial effect of human mesenchymal stem cells is mediated in part from secretion of the antimicrobial peptide LL-37. *Stem Cells* 2010;28:2229–2238.
8. Jackson MV, Morrison TJ, Doherty DF, McAuley DF, Matthay MA, Kissenpfennig A, et al. Mitochondrial transfer via tunneling nanotubes is an important mechanism by which mesenchymal stem cells enhance macrophage phagocytosis in the *in vitro* and *in vivo* models of ARDS. *Stem Cells* 2016;34:2210–2223.
9. McIntyre LA, Moher D, Fergusson DA, Sullivan KJ, Mei SH, Lalu M, et al.; Canadian Critical Care Translational Biology Group. Efficacy of mesenchymal stromal cell therapy for acute lung injury in preclinical animal models: a systematic review. *PLoS One* 2016;11:e0147170.

10. Matthay MA, Calfee CS, Zhuo H, Thompson BT, Wilson JG, Levitt JE, *et al.* Treatment with allogeneic mesenchymal stromal cells for moderate to severe acute respiratory distress syndrome (START study): a randomised phase 2a safety trial. *Lancet Respir Med* 2019; 7:154–162.
11. Combes A, Hajage D, Capellier G, Demoule A, Lavoué S, Guerville C, *et al.*; EOLIA Trial Group, REVA, and ECMONet. Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome. *N Engl J Med* 2018;378:1965–1975.
12. Hoogduijn MJ, van den Beukel JC, Wiersma LCM, Ijzer J. Morphology and size of stem cells from mouse and whale: observational study. *BMJ* 2013;347:f6833.
13. Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, *et al.* Minimal criteria for defining multipotent mesenchymal stromal cells: the International Society for Cellular Therapy position statement. *Cytotherapy* 2006;8:315–317.
14. Millar JE, von Bahr V, Malferteiner MV, Ki KK, Redd MA, Bartnikowski N, *et al.* Administration of mesenchymal stem cells during ECMO results in a rapid decline in oxygenator performance. *Thorax* 2019; 74:194–196.
15. Cho HJ, Hong H, Kim DW, Lee KS, Han HS, Kim GH, *et al.* Viability of mesenchymal stem cells in an *ex vivo* circulation system. *ASAIO J* 2020;66:433–440.
16. Millar JE, Fanning JP, McDonald CI, McAuley DF, Fraser JF. The inflammatory response to extracorporeal membrane oxygenation (ECMO): a review of the pathophysiology. *Crit Care* 2016;20:387.
17. National Health and Medical Research Council. Australian code for the care and use of animals for scientific purposes, 8th ed. Canberra, Australia: NHMRC; 2013.
18. Kilkeny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving bioscience research reporting: the ARRIVE guidelines for reporting animal research. *Osteoarthritis Cartilage* 2012;20:256–260.
19. Matute-Bello G, Downey G, Moore BB, Groshong SD, Matthay MA, Slutsky AS, *et al.*; Acute Lung Injury in Animals Study Group. An official American Thoracic Society workshop report: features and measurements of experimental acute lung injury in animals. *Am J Respir Cell Mol Biol* 2011;44:725–738.
20. Millar JE, Bartnikowski N, von Bahr V, Malferteiner MV, Obonyo NG, Belliato M, *et al.*; European Extracorporeal Life Support Organisation (EuroELSO) Innovations Workgroup; National Health Medical Research Council Australia Centre of Research Excellence for Advanced Cardio-respiratory Therapies Improving Organ Support (NHMRC CREATIONS). Extracorporeal membrane oxygenation (ECMO) and the acute respiratory distress syndrome (ARDS): a systematic review of pre-clinical models. *Intensive Care Med Exp* 2019;7:18.
21. Griffiths MJD, McAuley DF, Perkins GD, Barrett N, Blackwood B, Boyle A, *et al.* Guidelines on the management of acute respiratory distress syndrome. *BMJ Open Respir Res* 2019;6:e000420.
22. Simonson OE, Mougialakos D, Heldring N, Bassi G, Johansson HJ, Dalen M, *et al.* In vivo effects of mesenchymal stromal cells in two patients with severe acute respiratory distress syndrome. *Stem Cells Transl Med* 2016;5:845.
23. Lin H-C, Wang C-C, Chou H-W, Wu E-T, Lu FL, Ko B-S, *et al.* Airway delivery of bone marrow-derived mesenchymal stem cells reverses bronchopulmonary dysplasia superimposed with acute respiratory distress syndrome in an infant. *Cell Med* 2018;10: 2155179018759434.
24. Kocyildirim E, Cárdenes N, Ting A, Cáceres E, Bermúdez C, Rojas M. The use of GMP-produced bone marrow-derived stem cells in combination with extracorporeal membrane oxygenation in ARDS: an animal model. *ASAIO J* 2017;63:324–332.
25. Christy BA, Herzig MC, Montgomery RK, Delavan C, Bynum JA, Reddoch KM, *et al.* Procoagulant activity of human mesenchymal stem cells. *J Trauma Acute Care Surg* 2017; 83(suppl 1): S164–S169.
26. Fiedler T, Rabe M, Mundkowski RG, Oehmcke-Hecht S, Peters K. Adipose-derived mesenchymal stem cells release microvesicles with procoagulant activity. *Int J Biochem Cell Biol* 2018;100:49–53.
27. Gleeson BM, Martin K, Ali MT, Kumar AH, Pillai MG, Kumar SP, *et al.* Bone marrow-derived mesenchymal stem cells have innate procoagulant activity and cause microvascular obstruction following intracoronary delivery: amelioration by antithrombin therapy. *Stem Cells* 2015;33:2726–2737.
28. Tatsumi K, Ohashi K, Matsubara Y, Kohori A, Ohno T, Kakidachi H, *et al.* Tissue factor triggers procoagulation in transplanted mesenchymal stem cells leading to thromboembolism. *Biochem Biophys Res Commun* 2013;431:203–209.
29. Curley GF, Hayes M, Ansari B, Shaw G, Ryan A, Barry F, *et al.* Mesenchymal stem cells enhance recovery and repair following ventilator-induced lung injury in the rat. *Thorax* 2012;67:496–501.
30. Pedrazza L, Cunha AA, Luft C, Nunes NK, Schmitz F, Gassen RB, *et al.* Mesenchymal stem cells improves survival in LPS-induced acute lung injury acting through inhibition of NETs formation. *J Cell Physiol* 2017;232:3552–3564.
31. Moodley Y, Sturm M, Shaw K, Shimbori C, Tan DB, Kolb M, *et al.* Human mesenchymal stem cells attenuate early damage in a ventilated pig model of acute lung injury. *Stem Cell Res (Amst)* 2016; 17:25–31.
32. Lalu MM, Sullivan KJ, Mei SH, Moher D, Straus A, Fergusson DA, *et al.* Evaluating mesenchymal stem cell therapy for sepsis with preclinical meta-analyses prior to initiating a first-in-human trial. *eLife* 2016;5: e17850.
33. McIntyre LA, Stewart DJ, Mei SHJ, Courtman D, Watpool I, Granton J, *et al.*; Canadian Critical Care Trials Group; Canadian Critical Care Translational Biology Group. Cellular immunotherapy for septic shock: a phase I clinical trial. *Am J Respir Crit Care Med* 2018;197: 337–347.
34. Galstyan G, Makarova P, Parovichnikova E, Kuzmina L, Troitskaya V, Ghemdzian E, *et al.* The results of the single center pilot randomized Russian clinical trial of mesenchymal stromal cells in severe neutropenic patients with septic shock (RUMCESS). *Int J Blood Res Disord* 2018;5:033.
35. Schlosser K, Wang J-P, Dos Santos C, Walley KR, Marshall J, Fergusson DA, *et al.*; Canadian Critical Care Trials Group and the Canadian Critical Care Translational Biology Group. Effects of mesenchymal stem cell treatment on systemic cytokine levels in a phase 1 dose escalation safety trial of septic shock patients. *Crit Care Med* 2019;47:918–925.
36. Perlee D, van Vught LA, Scicluna BP, Maag A, Lutter R, Kemper EM, *et al.* Intravenous infusion of human adipose mesenchymal stem cells modifies the host response to lipopolysaccharide in humans: a randomized, single-blind, parallel group, placebo controlled trial. *Stem Cells* 2018;36:1778–1788.
37. Laroye C, Lemarié J, Boufenzar A, Labroca P, Cunat L, Alauzet C, *et al.* Clinical-grade mesenchymal stem cells derived from umbilical cord improve septic shock in pigs. *Intensive Care Med Exp* 2018;6:24.
38. Laroye C, Boufenzar A, Jolly L, Cunat L, Alauzet C, Merlin JL, *et al.* Bone marrow vs Wharton's jelly mesenchymal stem cells in experimental sepsis: a comparative study. *Stem Cell Res Ther* 2019;10:192.
39. Cardenes N, Aranda-Valderrama P, Carney JP, Sellares Torres J, Alvarez D, Kocyldirim E, *et al.* Cell therapy for ARDS: efficacy of endobronchial versus intravenous administration and biodistribution of MAPCs in a large animal model. *BMJ Open Respir Res* 2019;6: e000308.
40. Matthay MA, Zemans RL, Zimmerman GA, Arabi YM, Beitler JR, Mercat A, *et al.* Acute respiratory distress syndrome. *Nat Rev Dis Primers* 2019;5:18.
41. Islam D, Huang Y, Fanelli V, Delsedime L, Wu S, Khang J, *et al.* Identification and modulation of microenvironment is crucial for effective mesenchymal stromal cell therapy in acute lung injury. *Am J Respir Crit Care Med* 2019;199:1214–1224.
42. Cavalcanti AB, Suzumura EA, Laranjeira LN, Paisani DM, Damiani LP, Guimarães HP, *et al.*; Writing Group for the Alveolar Recruitment for Acute Respiratory Distress Syndrome Trial (ART) Investigators. Effect of lung recruitment and titrated positive end-expiratory pressure (PEEP) vs low PEEP on mortality in patients with acute respiratory distress syndrome: a randomized clinical trial. *JAMA* 2017;318: 1335–1345.