

CHARMM-GUI

Effective Simulation Input Generator and More

CHARMM is a versatile program for atomic-level simulation of many-particle systems, particularly macromolecules of biological interest. - M. Karplus

Some [lectures](#), [job postings](#), and [FAQ](#) are now available. See [update log](#) for update history and [giving](#) for donation. [Contact](#) info is given below.

[Logout](#)

Bilayer Builder

[User Profile](#)

Bookmark this [link](#), if you want to comeback to this page

PDB Info	STEP 1	STEP 2	STEP 3	STEP 4	STEP 5	STEP 6	JOB ID: 3750189754
----------	--------	--------	--------	--------	--------	--------	--------------------

Title	
PDB ID	BTAAC1_CDL_CSTATE_2C3E
Type	Protein
Experimental Method	Unknown

PDB Manipulation Options:

☒ System pH: [Apply](#)☒ Reading Hetero Chain Residues:

CDL ☒ Rename to [CSML Search](#) **Click this if you want to generate your ligand FF using the PDB coordinate.**

☐ Use CHARMM General Force Field to generate CHARMM top & par files (using [ParamChem](#) service) [?](#)

☐ Use Antechamber to generate CHARMM top & par files

☐ Use OpenFF to generate CHARMM top & par files

☐ Upload CHARMM top & par for hetero chain [?](#)

☒ Protonate/Deprotonate based on selected pH

Ligand reading and its FF generation could be tricky, so please read these common [issues](#).

If this residue exists in the CHARMM topology files ([toppar.str](#)), please rename (or retype) it to the CHARMM residue name. Otherwise, CHARMM will be terminated abnormally.

☒ Terminal group patching: [?](#)

First Last
PROA ☐ Cyclic peptide?

☐ Preserve hydrogen coordinates:☐ Mutation:☐ Protonation state:☐ Disulfide bonds:☐ Phosphorylation:☐ Ubiquitylation / SUMOylation:

☐ GPI anchor:

☐ Glycosylation / Glycan Ligand(s): ☐ Use CHARMM MC?

"test_distortion.txt" file.

It is faster than the regular run, but carefully check the output

☐ Heme coordination

☐ Add Lipid-tail 

☐ Peptide Stapling 

☐ Add FRET/LRET fluorophore labels 

☐ Model LBT-loop(s) 

☐ Add MTS reagents: nitroxide spin labels 

☐ Add MTS reagents: chemical modifier 

☐ Non-standard amino acid / RNA substitution:

☐ Lys / Arg PTMs: 

Symmetry Operation Options:

Next Step:
Generate PDB and Orient Molecule



[Lehigh University](#) / [Department of Biological Sciences](#) / [Department of Chemistry](#) / [Department of Bioengineering](#) / [Im Lab](#)
Problems, Questions, & Comments? / [Contact](#) / [Forum](#) / Copyright(c) 2006-2025 by the Im Lab