

The basic outline/plan

- Lefschetz condition for “nice” DM stack, following Hartshorn ASAV chapter 4, begin a definition here
- something

DM Stack basically means that the stabilizers groups are all finite.

## 1 Introduction

**Definition 1.** this is where the definition of Lef for weighted projective stacks will go

**Definition 2.** this is where the definition of Leff for weighted projective stacks will go

## 2 Formal Geometry on Stacks

See this source: <https://math.stanford.edu/~conrad/papers/formalgaga.pdf>

See this source: <https://www.math.uchicago.edu/~emerton/pdffiles/formal-stacks.pdf>

Comment: Probably the most technical part (and the most useful) is going to be here.

**Definition 3.** Formal completion of a stack along a substack

**Proposition 1.** Following 1.1 of Hartshorn ASAV, equivalence of  $\text{cd}(X-Y) < n - 1$  is equivalent to  $\text{Lef}(X, Y)$  and  $Y$  meets every effective Cartier divisor on  $X$

**Definition 4.** Give the correct definition of a complete intersection on a stack, probably intersection of divisors

**Theorem 2.** Following theorem 1.5 of Hartshorn ASAV, give the correct statement of this theorem. As a first attempt take  $X$  to be equal to the weighted projective stack. Something about fake weighted projective stacks (maybe?)

A complete intersection,  $Y$ , has  $\text{Leff}(X, Y)$  assuming that  $Y$  is dimension at least 2.

## 3 Application to the Picard Group

**Theorem 3.** Following theorem 3.1 in Hartshorn ASAV

*Proof.* What we need for the proof is:

- Deformation theory argument
- third point of Hartshorn’s version deals with  $H^i(Y, I^n/I^{n+1})$  this is where the deformation theory comes in
- We have a line bundle  $L$  on  $Y$ , we want to lift it to  $X$
- There are a series of “natural” maps  $\text{Pic}(X) \rightarrow \text{Pic}(U) \rightarrow \text{Pic}(\hat{X}) \rightarrow \text{Pic}(Y)$ . Need to show that they are all isomorphisms and that their composition is the “natural” map  $\text{Pic}(X) \rightarrow \text{Pic}(Y)$  that comes from the pull back  $\mathcal{L} \mapsto i^*(\mathcal{L})$  where  $i : Y \hookrightarrow X$  is the inclusion of  $Y$  into  $X$ .

□

## 4 Cohomological properties of DM Stacks

Serre duality for tame DM stacks. (Tame basically means that if characteristic =  $p$ , then stablizer groups do not have order  $p^n$  for any  $n$ )

See: <https://link.springer.com/article/10.1007/s40687-022-00367-7>

Serre's theorem on global generation on Stacks

See: <https://arxiv.org/abs/1306.5418>

**Theorem 4.** Generalizing theorem 5.18 in Hartshorn's AG

## 5 Calculate the cohomology of line bundles on weighted projective stack

**Theorem 5.** Generalizing theorem 5.1 in Hartshorn's AG