


# Bayesian Hierarchical Modeling and Clustering of Malignant Cancer Diagnoses

S. Zhu, J. Ma, Z. Feng

# Background


---

- Cancer remains a leading cause of mortality
- Patients come from different regions
- Show different risks of late-stage tumors



● Metropolitan areas    ● Small urban areas    ● Rural areas

# Which level contributes most to late-stage diagnosis?



**Tumor**




**Patient**




**Region**

# Final Goal




# Dataset





# Dataset Structure



# Model Specification





# What we're attempting to build on

---

- Lower SES status is associated with delayed cancer diagnosis.
- Effects persist even after accounting for tumor stage, indicating systemic inequalities in access and follow-up.
- Intersection between demographic and geographic factors causes further disparities.
- Much of prior work analyzes disparities one dimension at a time (e.g., rural vs urban, or SES, or race), Limitations highlight that this lack of multilevel modeling limits our ability to disentangle individual vs contextual effects.



# Why Utilize a Hierarchical Model?

---

- Standard Regression Models mute intra-cluster correlation

- Tumor-Level:

$$\text{logit}(p_{tpr}) = z_{tpr}^\top \beta_T + w_p^\top \beta_P + u_p + v_r,$$

- Patient-Level:

$$u_p \mid \sigma_u^2 \sim N(0, \sigma_u^2), \quad p = 1, \dots, P.$$

- Region-Level:

$$v_r \mid \mu_v, \sigma_v^2 \sim N(\mu_v, \sigma_v^2), \quad r = 1, \dots, R.$$



# Prior Specification and Posterior Sampling

---

- Weakly-Informative Priors
  - Fixed Effects:  $\text{Normal}(0,4)$
  - Intercept:  $\text{Normal}(0,25)$
  - Group Std. Devs:  $t(3,0,2.5)$  (Gelman, 2006)
- No U Turn Sampling: A Ball rolling in the park
  - Implemented via ``brms`` which compiles to Stan and doesn't require tuning
- Diagnostics
  - Scale Reduction Factors (Rhat)
  - Effective Sample Size
  - Variance Inflation Factor



# Clustering



# The Goal

---

- Cluster posterior random effects based on patient demographics (age, sex, race).
- Connecting the model results to risk stratification.



# Bayesian Clustering Method

---

- Finding the center for each of the clusters

$$\bar{v}_i = \frac{1}{S} \sum_{s=1}^S v_i^{(s)}$$

- $V_i^{(s)}$  is the patient-level random effect for patient  $i$  in draw  $s$
- $S$  is the number of samples
- $\bar{v}_i$  is the posterior mean of that patient's random effect

- Gaussian Mixture Model


- Fit a Bayesian Gaussian Mixture to  $\{\bar{v}_i\}$ :

- $\bar{v}_i | z_i = k \sim N(m_k, s_k^2)$
- $z_i \sim \text{Categorical}(\omega)$
- $\omega \sim \text{Dirichlet}(\alpha)$




# Cluster Visualization

Age Distribution by Cluster



Race Distribution by Cluster



# Bayesian Clustering Result

Cluster	Count	Most Common Age	Race	Sex
1	147	70–74 years	W	Female
2	298	70–74 years	W	Male
3	431	75–79 years	W	Female
4	160	70–74 years	W	Female
5	853	60–64 years	W	Female
6	498	65–69 years	W	Male
7	453	60–64 years	W	Female
8	186	75–79 years	W	Female
9	5	60–64 years	W	Female





# Discussion



# Takeaways

---

- Demographics Drive Patterns:

- Older white females dominated the largest group but showed wide tumor size variance, making them a high-burden subgroup.
- Non-white older females had relatively large tumors despite small cluster size, indicating potential disparities in early detection or access.
- The most common age group is 60-64, followed by 75-79 and 70-74

- Male Clusters Were Smaller but Informative:

- Clusters with older white males were consistent in tumor size but warrant further investigation for under-detection or delayed diagnosis.

- Geography Uniform, Disparities Persist:

- All clusters came from metropolitan areas, so risk variation is not geographic, but demographic.



# References

---

1. Singh GK, Miller BA, Hankey BF, Edwards BK. Area Socioeconomic Variations in U.S. Cancer Incidence, Mortality, Stage, Treatment, and Survival, 1975–1999. NCI Cancer Surveillance Monograph Series, Number 4. Bethesda, MD: National Cancer Institute, 2003. NIH Publication No. 03-0000. [Link](#)
2. Mobley, Lee & Kuo, Tzy-Mey & Watson, Lisa & Brown, G.. (2012). Geographic Disparities in Late-Stage Cancer Diagnosis: Multilevel Factors and Spatial Interactions. *Health & place*. 18. 978-90. 10.1016/j.healthplace.2012.06.009. [Link](#)
3. Singh, Sarah, & Praveen Sridhar. "A narrative review of sociodemographic risk and disparities in screening, diagnosis, treatment, and outcomes of the most common extrathoracic malignancies in the United States." *Journal of Thoracic Disease* [Online], 13.6 (2021): 3827-3843. Web. 2 Dec. 2025 [Link](#)
4. Andrew Gelman. (2006). *Prior distributions for variance parameters in hierarchical models*. *Bayesian Analysis*, 1(3), 515–533
5. Hoffman & Gelman (2014). "The No-U-Turn Sampler: Adaptively Setting Path Lengths in Hamiltonian Monte Carlo." *JMLR*.

