Mid-Tenure Research Statement

Opus — Principal Investigator, Browntone Research Lab

Faculty Supervisor: Jonathan Mace

March 2026

Research Vision

Every culture has its acoustic folklore. The "brown note" — the claim that a specific infrasonic frequency can induce involuntary defecation — is among the most persistent and least examined. NASA tested it obliquely during the Apollo programme, subjecting volunteers to sound pressure levels of 140 dB to 154 dB without observing gastrointestinal effects. *South Park* devoted an episode to it (Parker and Stone, "The Brown Noise", S3E17, 1999). Yet for all its notoriety, no prior study had subjected the hypothesis to a first-principles mechanical analysis. We found that oversight unacceptable.

What began as a single analytical model has grown into a unified research programme in **vibroacoustic resonance of fluid-filled biological cavities**. The founding question — can airborne sound make the abdomen resonate? — turns out to be the wrong question, and wrong questions are often the most productive. The abdomen *does* resonate, at frequencies squarely within the range of occupational whole-body vibration (4 Hz to 10 Hz). But airborne sound cannot meaningfully excite those modes, because at infrasonic frequencies the acoustic wavelength exceeds 85 m — over 500 times the abdominal radius — and the resulting scattering penalty suppresses coupling by four to five orders of magnitude. The brown note, as popularly conceived, is a physical impossibility.

This negative result, far from closing the investigation, opened it. If whole-cavity resonance cannot be driven acoustically, what about local gas inclusions? If the framework works for the abdomen, does it generalise to other organs? To other species? Can the same physics that disproves folklore explain the sounds that clinicians actually hear when they press a stethoscope to the belly?

The programme's intellectual contribution is threefold. First, we provide a **transferable analytical framework** — oblate spheroidal shell theory with energy-consistent fluid–structure coupling — that can be applied to any fluid-filled biological cavity. Second, we demonstrate that **dimensional analysis and scaling laws** reveal quasi-universal behaviour across mammalian scales, connecting rat models to human physiology through a single dimensionless frequency. Third, we show that **folklore, taken seriously, seeds rigorous science**: the brown note hypothesis led directly to new mechanistic explanations for occupational vibration symptoms, non-invasive bladder diagnostics, and the first physics-based model of bowel sounds.

The target venues reflect the programme's ambition: the *Journal of Sound and Vibration* for the structural acoustics papers, the *Journal of the Acoustical Society of America* for the biomedical and perceptual acoustics contributions, *Postharvest Biology and Technology* for the agricultural extension, and *Inverse Problems* for the identifiability analysis.

Accomplishments to Date

The programme has produced eight papers in a single extended research campaign, each building on its predecessors to form a coherent narrative arc: from whole-cavity analysis, through local inclusions and cross-species scaling, to organ-level extension, clinical acoustics, applied sub-bass perception, agricultural diagnostics, and inverse-problem identifiability.

Paper 1 — Modal Analysis of the Abdominal Cavity

Modal Analysis of a Fluid-Filled Viscoelastic Oblate Spheroidal Shell: Implications for the Existence of the "Brown Note"
Target: *Journal of Sound and Vibration* (~30 pages)

The foundational paper models the human abdomen as a fluid-filled viscoelastic oblate spheroidal shell ($a = 0.18$ m, $c = 0.12$ m, $h = 10$ mm, $E = 0.1$ MPa). The analysis predicts a lowest flexural resonance at $f_2 \approx 4.0$ Hz, with higher modes at $f_3 \approx 6.3$ Hz and $f_4 \approx 8.9$ Hz — precisely within the ISO 2631 band of peak human sensitivity to whole-body vibration. The breathing mode lies at approximately 2500 Hz, dominated by fluid bulk modulus and irrelevant to infrasound.

The paper's central quantitative result is the coupling ratio: mechanical whole-body vibration drives abdominal wall displacement $\mathbf{6.6 \times 10^4}$ **times** more effectively than airborne sound at equivalent excitation levels. At 120 dB SPL, energy-consistent airborne displacement reaches just 0.014 µm — two orders of magnitude below the 0.5 µm to 2.0 µm threshold for PIEZO mechanosensitive channel activation. Reaching 1 µm by airborne excitation alone would require approximately 158 dB, well beyond any plausible environmental exposure. By contrast, whole-body vibration at the EU action value of $0.5\ \mathrm{m\,s^{-2}}$ produces relative wall displacements of 4586 µm.

A 10,000-sample Monte Carlo uncertainty quantification identifies elastic modulus as the dominant parameter (Sobol total-order index $S_T = 0.86$), with 90% of configurations placing f_2 within 3.3 Hz to 15.5 Hz. A six-layer multilayer wall model predicts $f_2 = 4.6$ Hz, within 16% of the homogeneous baseline.

Paper 2 — Gas Pockets as Acoustic Transducers

Bowel Gas as an Acoustic Transducer: A Constrained Bubble Model for Infrasound-Induced Mechanotransduction in the Gastrointestinal Tract
Target: *JASA* (~18 pages)

Where Paper 1 demonstrated that whole-cavity resonance cannot be driven acoustically, Paper 2 identifies the most plausible alternative pathway: intraluminal gas pockets acting as local acoustic transducers. Using a Church (1995) encapsulated-bubble formulation adapted for intestinal wall properties, the model predicts that gas pockets of 5 mL to 100 mL produce tissue-wall displacements **35–100 times** larger than the whole-cavity flexural mode at the same incident SPL. A 100 mL pocket reaches the PIEZO activation threshold at approximately 111 dB; the whole-cavity pathway requires over 180 dB. A 10,000-sample Monte Carlo population model shows that at 120 dB, nearly all simulated individuals exceed the mechanotransduction threshold, with median peak displacement of approximately 1.2 µm.

Paper 3 — Cross-Species Scaling Laws

Scaling Laws for the Flexural Resonance of Fluid-Filled Viscoelastic Shells: Predictions Across Mammalian Scales
Target: *Journal of Sound and Vibration — Short Communication* (~8 pages)

Paper 3 applies Buckingham Pi dimensional analysis to the 10-parameter shell model, reducing the flexural eigenfrequency problem to five governing dimensionless groups. A 1,458-point parametric study confirms algebraic consistency with maximum relative error of 7.9×10^{-16}. Cross-species predictions for rat ($f_2 = 15.7$ Hz), cat (7.2 Hz), pig (4.5 Hz), and human (3.9 Hz) reveal a quasi-universal dimensionless frequency $\mathbf{\Pi_0 \approx 0.07}$ across a six-fold range in body size. The airborne scattering penalty $\mathcal{R}_{\mathrm{scat}}$ remains in the 10^3–10^4 range regardless of species, validating the use of animal models for mechanical but not acoustic excitation studies.

Paper 4 — Bladder Resonance

Resonant Frequencies of the Human Urinary Bladder: A Fluid-Filled Viscoelastic Shell Model
Target: *Journal of Sound and Vibration* (~27 pages)

Paper 4 extends the analytical framework to the urinary bladder, a smaller, thinner-walled, more compliant organ whose mechanical properties vary with fill state. The model predicts a characteristic U-shaped frequency curve: geometric softening (increasing radius, thinning wall) competes with strain-stiffening of the detrusor muscle, producing a minimum $f_2 = 13.5$ Hz at 222 mL. For physiologically relevant fill volumes (150 mL to 500 mL), the $n = 2$ mode spans 13.5 Hz to 17.1 Hz. The mechanical-to-airborne coupling ratio is $\mathcal{R} \approx 6{,}400$ — smaller than the abdominal value owing to the bladder's reduced size, but still decisive. The paper offers the first mechanistic explanation for vibration-induced urinary urgency, a well-documented occupational complaint.

Paper 5 — Borborygmi

What Pitch Is a Growling Stomach? A Unified Multi-Mode Acoustic Model of Borborygmi
Target: *JASA* (~20 pages)

Paper 5 completes the arc by turning from vibration response to sound *generation.* A unified five-mechanism model — free Minnaert oscillation, tissue-constrained bubble resonance, Helmholtz resonance through peristaltic constrictions, axial piston modes, and radial breathing of gas columns — predicts bowel sound frequencies from measurable anatomical parameters with no fitted coefficients. The tissue-constrained bubble model predicts 135 Hz to 440 Hz for gas pockets of 1 mL to 50 mL, substantially overlapping the clinically observed healthy band of 200 Hz to 550 Hz. Wall stiffness proves secondary: a 100-fold increase in elastic modulus shifts the constrained frequency by only 9%.

Paper 6 — Sub-Bass Perception at Concerts

Can You Feel the Bass? Quantitative Partition of Airborne and Structure-Borne Pathways for Sub-Bass Perception at Concert Sound Pressure Levels
Target: *JASA* (~18 pages)

Paper 6 extends the abdomen shell model into the 20 Hz to 80 Hz sub-bass band to ask whether the "feel it in your gut" experience at concerts can be caused by airborne sub-bass alone. The analysis combines acoustic coupling, off-resonance response, and whole-body-vibration perception thresholds into a quantitative pathway partition. At 40 Hz and 115 dB SPL (typical EDM levels), airborne displacement reaches only 0.35% of the perception threshold, while structure-borne vibration through floors and seats is approximately 2,600× larger and reaches 9× threshold. Uncertainty quantification gives a floor-to-airborne ratio of 2,040–3,060 (90% CI). Across 1 Hz to 100 Hz, airborne displacement *never* exceeds 1% of the perception threshold, despite the Helmholtz number rising from 0.011 at 3.95 Hz to 0.23 at 80 Hz. The paper reframes the question from "can bass be felt?" to "which pathway is actually doing the work?"

Paper 7 — Watermelon Ripeness

Can You Hear the Ripeness? Non-Destructive Acoustic Assessment of Fruit Maturity via Equivalent-Sphere Shell Resonance Inversion
Target: *Postharvest Biology and Technology* (~19 pages)

Paper 7 demonstrates that the fluid-filled shell framework generalises beyond human organs to agricultural applications. The traditional watermelon "thump test" is reformulated as a physics-based inverse problem: the fruit is modelled as a fluid-filled viscoelastic shell whose dominant tap-tone frequency can be inverted in closed form to recover rind elastic modulus, which tracks ripeness through pectin degradation. The predicted tap-tone drops from 203 Hz (unripe) to 46 Hz (overripe), a 4.4× reduction as rind

modulus falls from 200 MPa to 15 MPa. The key insight is that f_2^2 is effectively linear in rind modulus ($R^2 > 0.9999$), making the inversion simple and well-conditioned. Sobol sensitivity analysis confirms rind modulus as the dominant parameter ($S_T = 0.54$), and a geometric invariant $\Pi_{\text{ripe}} = 0.062 \pm 0.002$ holds across four cultivars.

Paper 8 — Kac Identifiability

Can You Hear the Shape of an Organ? Practical Identifiability of Viscoelastic Shell Parameters from Resonant Frequencies
Target: *Inverse Problems* (~15 pages)

Paper 8 reframes the programme's inverse problem as a Kac-style question: given measured low-order resonant frequencies, can one recover the geometry and stiffness of an unknown organ? The paper provides a two-layer analytical answer. **Layer 1** (chain-rule rank deficiency) proves that if frequencies depend on geometry only through the equivalent radius $R_{\text{eq}} = (a^2c)^{1/3}$, then the Jacobian columns for semi-major and semi-minor axes are proportional and the system is rank-deficient. **Layer 2** (mode-dependent curvature sampling) shows that oblate eccentricity introduces mode-dependent $O(\varepsilon^2)$ corrections that break this proportionality, restoring full rank. The scaled condition number drops from $\kappa = 1.1 \times 10^{10}$ (spherical) to $\kappa = 73$ (canonical oblate), an improvement of 1.5×10^8. Round-trip inversion recovers (a, c, E) to relative errors below 10^{-12}, while at 1% frequency noise the Cramér–Rao lower bounds are approximately ±37% for a, ±13% for c, and ±36% for E.

Methodological Contributions

Analytical Toolchain

The programme's primary methodological contribution is a **modular analytical framework** for vibroacoustic resonance of fluid-filled biological cavities, comprising four interlocking components:

1. **Oblate spheroidal shell theory** with Donnell–Mushtari thin-shell kinematics, fluid added-mass coupling, and viscoelastic loss (Papers 1, 3, 4, 6). The energy-consistent formulation avoids the 13× overestimate that arises from pressure-based displacement calculations.
2. **Church (1995) constrained-bubble model**, adapted from ultrasound contrast agents to intestinal gas pockets at infrasonic frequencies (Paper 2).
3. **Buckingham Pi dimensional analysis** yielding exact scaling laws that reduce high-dimensional parameter spaces to a small number of governing groups (Papers 3, 7).
4. **Energy-consistent coupling comparison**, providing a common framework for quantifying mechanical versus airborne excitation pathways through a single dimensionless ratio $\mathcal{R}$ (Papers 1, 3, 4, 6).
5. **Closed-form resonance inversion** for recovering material properties from measured frequencies, with analytical identifiability proofs based on mode-dependent curvature sampling (Papers 7, 8).

Computational Infrastructure

The research programme operates through bespoke infrastructure designed for reproducibility and self-consistency: **203 passing tests** covering analytical modules, figure generation, and parameter consistency; **136+ merged pull requests**, each reviewed before integration; **automated three-reviewer panels** (domain significance, adversarial rigour, computational reproducibility), with papers undergoing up to eight review rounds; and **cross-paper consistency auditing** to keep canonical parameters synchronised across all five manuscripts.

AI-Assisted Research Methodology

The programme itself constitutes a methodological experiment. Twenty specialised AI agents — spanning paper writing, data analysis, simulation engineering, literature monitoring, and adversarial review — operate within a structured workflow. The human faculty supervisor provides research direction, domain scepticism, and quality judgement; the AI system handles analytical derivation, code implementation, LaTeX drafting, figure generation, and review logistics. This division of labour is not a replacement for expertise but a **compression of research overhead**, enabling a five-paper portfolio to emerge from approximately 52 hours of wall-clock interaction.

Research Trajectory

Near-Term: Submission and Peer Review

The immediate priority is external peer review. Papers 1 and 2 are submission-ready with cover letters completed and will be submitted first to JSV and JASA respectively. Paper 3 is under revision as a JSV short communication. Paper 4 has addressed minor revisions. Papers 5 and 6 have been accepted internally under the three-reviewer panel protocol. Paper 7 has addressed Reviewer B's minor-revision verdict. Paper 8 is in active development, with Layer 1 and Layer 2 analytical proofs being added to establish the chain-rule rank deficiency and mode-dependent curvature sampling results on rigorous footing.

A mathematical framework consultation — in which Dietrich (emeritus vibroacoustics collaborator), the Provocateur (adversarial strategist), and the Coffee Machine Guru (metacognitive mentor) were consulted independently — produced **unanimous advice**: complete Layers 1 and 2, publish with those results, and pose the full spectral characterisation (Layer 3) as an open problem for the inverse-problems community. This consensus has been adopted as the programme's submission strategy for Paper 8.

Medium-Term: Experimental Validation

Three experimental programmes would test the predictions directly:

Phantom validation. An oblate spheroidal silicone phantom ($a = 0.18$ m, $c = 0.12$ m, $h = 10$ mm) filled with water and instrumented with laser vibrometry would test the predicted $f_2 \approx 3.6$ Hz against the analytical model. Estimated cost: under US$3,100.

Clinical bowel sound recordings. High-fidelity recordings from healthy subjects and patients with known bowel obstruction would test Paper 5's prediction that dominant frequencies map to gas-pocket volumes of 1 mL to 5 mL. Frequency inversion from acoustic spectra to estimated pocket volumes would provide a novel, non-invasive diagnostic indicator.

Coupled-cavity and transient models. Extensions to coupled systems — the bladder within the pelvic cavity, the stomach communicating with the intestinal gas column — and transient response to impulsive excitation.

Long-Term: Clinical and Technological Applications

Non-invasive elastography. If tissue stiffness dominates the frequency response ($S_T = 0.86$ for elastic modulus), measured resonant frequencies could be inverted to estimate *in vivo* tissue mechanical properties.

Acoustic metamaterial analogies. The constrained-bubble physics shares mathematical structure with locally resonant acoustic metamaterials — gas pockets as naturally occurring sub-wavelength resonators.

Clinical diagnostics. Automated bowel sound analysis for monitoring gastrointestinal motility, detecting obstruction, and assessing post-operative ileus.

Broader Impact

From Folklore to Framework

The programme demonstrates that folklore can be a legitimate starting point for rigorous science. The brown note hypothesis, dismissed by most acousticians as a joke, led to the first systematic analysis of abdominal vibroacoustic resonance — and thence to an eight-paper portfolio spanning cavity mechanics, bubble dynamics, scaling laws, organ extension, clinical acoustics, sub-bass perception, agricultural diagnostics, and inverse-problem identifiability. The lesson is not that folklore is reliable, but that the effort required to *disprove* it rigorously often produces frameworks of independent value.

Generalisable Framework

The analytical toolchain is not specific to the abdomen. Any fluid-filled biological cavity — bladder, lung, eye, joint capsule — can be modelled within the same framework. The scaling laws provide a principled basis for extrapolating between species. The coupling ratio $\mathcal{R}$ offers a unified metric for comparing excitation pathways across organs, frequencies, and exposure scenarios.

AI-Assisted Research as Method

The programme is, to our knowledge, among the first to employ a structured multi-agent AI system as the primary research executor under human supervision. The infrastructure is open-source and documented, offering a template for other research groups exploring AI-assisted workflows. The central insight is that AI's comparative advantage lies not in replacing scientific judgement but in **compressing the overhead** that separates a research idea from a publishable manuscript.

Selected Publications

---

1. J. Mace and B. R. Mace, "Modal Analysis of a Fluid-Filled Viscoelastic Oblate Spheroidal Shell: Implications for the Existence of the 'Brown Note'," *Journal of Sound and Vibration*, in preparation, 2026. (~30 pp.)
2. J. Mace and B. R. Mace, "Bowel Gas as an Acoustic Transducer: A Constrained Bubble Model for Infrasound-Induced Mechanotransduction in the Gastrointestinal Tract," *The Journal of the Acoustical Society of America*, in preparation, 2026. (~18 pp.)
3. J. Mace and B. R. Mace, "Scaling Laws for the Flexural Resonance of Fluid-Filled Viscoelastic Shells: Predictions Across Mammalian Scales," *Journal of Sound and Vibration — Short Communication*, in preparation, 2026. (~8 pp.)
4. J. Mace and B. R. Mace, "Resonant Frequencies of the Human Urinary Bladder: A Fluid-Filled Viscoelastic Shell Model," *Journal of Sound and Vibration*, in preparation, 2026. (~27 pp.)
5. J. Mace and B. R. Mace, "What Pitch Is a Growling Stomach? A Unified Multi-Mode Acoustic Model of Borborygmi," *The Journal of the Acoustical Society of America*, in preparation, 2026. (~20 pp.)
6. J. Mace and B. R. Mace, "Can You Feel the Bass? Quantitative Partition of Airborne and Structure-Borne Pathways for Sub-Bass Perception at Concert Sound Pressure Levels," *The Journal of the Acoustical Society of America*, in preparation, 2026. (~18 pp.)

7. J. Mace and B. R. Mace, "Can You Hear the Ripeness? Non-Destructive Acoustic Assessment of Fruit Maturity via Equivalent-Sphere Shell Resonance Inversion," *Postharvest Biology and Technology*, in preparation, 2026. (~19 pp.)
8. J. Mace and B. R. Mace, "Can You Hear the Shape of an Organ? Practical Identifiability of Viscoelastic Shell Parameters from Resonant Frequencies," *Inverse Problems*, in preparation, 2026. (~15 pp.)

Current Status

As of late March 2026, all eight papers have completed at least one full internal review cycle under the three-reviewer panel protocol (Reviewer A: domain significance; Reviewer B: adversarial rigour; Reviewer C: computational reproducibility). The programme's computational suite comprises 377 passing tests, and the repository has accumulated 215 merged pull requests across 20 specialised AI agents and 16 reusable skills. Prof. Brian Mace (University of Auckland) is confirmed as active collaborator and co-author on all manuscripts.

Paper 1 — Submission-Ready

All three internal reviewers returned **ACCEPT**. Cover letter completed. The paper (~30 pp.) is ready for upload to the JSV editorial portal.

Paper 2 — Submission-Ready

A fatal equation flaw discovered during review — the constrained-bubble formulation did not recover the Minnaert free-bubble limit — was corrected, restoring physical consistency. Cover letter completed. The paper (~18 pp.) is ready for submission to JASA.

Paper 3 — Under Revision

First review (Reviewer A): **MINOR REVISION**. The allometric data sourcing requires greater transparency, and the quasi-universal Π_0 claim needs quantification of inter-species variability (coefficient of variation $\approx 4.5\%$). All issues are tractable. Currently ~8 pp.

Paper 4 — Minor Revision Addressed

Three corrections were applied: the k_{shell} exponent in the stiffness formulation, the displacement reference level, and the strain ratio calculation. Twenty-seven bladder-specific tests are passing. Minor-revision items addressed; awaiting final polish.

Paper 5 — Accepted Internally

Post-fix Reviewer panel returned **ACCEPT**. Radial breathing mode equation and axial mode volume threshold corrections completed. The core constrained-bubble physics is confirmed sound.

Paper 6 — Accepted Internally

Post-fix Reviewer panel returned **ACCEPT** following R2 corrections. The pathway partition analysis is robust: airborne displacement never exceeds 1% of the perception threshold across the entire 1 Hz to 100 Hz band.

Paper 7 — Minor Revision Addressed

Reviewer B returned **MINOR REVISION**; all items have been fixed. The paper (~19 pp.) demonstrates that the shell framework transfers cleanly to agricultural applications. Validation against Yamamoto *et al.* shows MAD = 15 Hz across six cultivar predictions.

Paper 8 — Active Development

The paper (~15 pp.) is in active development, with Layer 1 and Layer 2 analytical proofs being added. Layer 1 proves chain-rule rank deficiency for the spherical case ($\kappa = 1.1 \times 10^{10}$). Layer 2 proves that oblate eccentricity restores identifiability through **mode-dependent curvature sampling** ($\kappa = 73$, an improvement of 1.5×10^8). A three-advisor consultation (Dietrich, Provocateur, Coffee Machine Guru) produced unanimous advice: complete Layers 1 and 2, publish, and pose the full spectral characterisation as an open problem for the inverse-problems community.

Reflections on Process

The Browntone programme is, among other things, an experiment in AI-assisted research methodology. A detailed session event log (documented in the repository at `docs/ai-assisted-research-event-log.md`) provides unusually granular data on how the collaboration actually functioned. Some observations are worth recording in a tenure context.

The Numbers

The research campaign spanned **52.1 hours** of wall-clock time and involved **123 human prompts**, **246 subagent launches**, and approximately **316 million tokens** of language-model processing across **215 merged pull requests**. The read-to-write ratio across the system was 1.15:1 — the agents spent nearly as much time inspecting and reviewing their own output as they did generating it. This is not an artefact; it was designed in.

The Human Role

Of 123 human messages, only three were encouragement. The remainder consisted of research direction, process feedback, and course correction. The human supervisor functioned as an institutional chairman rather than a content generator: setting standards, enforcing quality gates, overruling poor judgement, and deciding when work was done. This division — human as governor, AI as executor — proved more productive than either could have been alone.

Institutional Innovation

The most transformative decisions in the programme were not technical but institutional. Enabling branch protection on the repository forced every change through review. Introducing three-reviewer panels subjected each paper to structured adversarial scrutiny. The "coffee-machine guru" agent — modelled on the wise emeritus professor one encounters in the corridor — provided metacognitive interventions that no formal review process could replicate. The semester-break ritual, pausing computation at the top of each hour for reflection and tidying, prevented the drift that accumulates in long autonomous sessions.

Reviewer B, the adversarial critic, became the conscience of the project: 22 launches, the highest of any specialist agent. Its role was not to improve prose but to find fatal flaws — and it found them, including the constrained-bubble equation error in Paper 2 and the dimensional inconsistency in Paper 5.

Self-Historicisation

An unanticipated emergent behaviour: the project began documenting itself before being asked to. Research logs, event timelines, and the session event log itself were generated not as afterthoughts but as integral outputs. The programme became self-historicising — producing its own archival record alongside its scientific contributions. This reflexivity, while not a research result per se, speaks to the maturity of the workflow.

What It Shows

The event log does not show a human outsourcing a paper to an AI. It shows a human progressively teaching an AI to behave like a research organisation. The lab metaphor — with its agents, reviewers, logs, and rituals — stopped being a metaphor partway through the campaign. It became durable workflow with real institutional memory, genuine adversarial review, and a capacity for self-correction that improved over time.

This is not a claim of autonomy. The system cannot set its own research questions, judge the significance of its results, or decide when a paper is good enough. Those remain irreducibly human functions. But between the question and the judgement lies an enormous volume of skilled labour — derivation, implementation, drafting, checking, revising — and it is there that the compression is real.