Highlights

Scaling Laws for the Flexural Resonance of Fluid-Filled Viscoelastic Shells: Predictions Across Mammalian Scales

Jonathan Mace

- Mammalian abdominal resonance governed by quasi-universal $\Pi_0 \approx 0.07$
- 486-point parametric study confirms algebraic consistency of non-dimensionalisation
- Cross-species f_2 scales as $(1/a)\sqrt{E/\rho_f}$ when other Π-groups are approximately constant
- Coupling ratio $\mathcal{R} \sim 10^3$–10^4 is size-independent: animal models are valid proxies
- Breathing mode never reaches infrasound for any organism

Scaling Laws for the Flexural Resonance of Fluid-Filled Viscoelastic Shells: Predictions Across Mammalian Scales

Jonathan Mace[a,*]

[a] *Department of Engineering, University of Example*

Abstract

Fluid-filled viscoelastic shells arise in contexts ranging from abdominal biomechanics to underwater acoustics. Modal frequencies of such shells depend on eleven physical parameters, making systematic comparison across geometries and species impractical without dimensional reduction. We apply Buckingham Pi analysis to the governing equations of a fluid-filled oblate spheroidal shell, identifying eight dimensionless groups of which six govern the flexural mode frequencies. The resulting scaling law, $f_n = \sqrt{E/\rho_f}\,(1/a)\,\Phi_n(\Pi_1, \ldots, \Pi_5)$, where Φ_n absorbs the $1/(2\pi)$ factor, admits a closed-form expression from the shell equations. A 486-point parametric study spanning physiological ranges of all parameters collapses onto the analytical curves to within numerical round-off (5.8×10^{-16}), confirming the algebraic consistency of the non-dimensionalisation. Cross-species predictions for four mammals (rat, cat, pig, human) reveal a quasi-universal dimensionless frequency $\Pi_0 \approx 0.07$, implying that the airborne coupling ratio $\mathcal{R} \sim 10^3$–10^4 is approximately size-independent and that animal models are valid proxies for human abdominal resonance studies. The breathing mode ($n = 0$), dominated by the fluid bulk modulus, would require a body radius of order 410 m to reach 20 Hz and is therefore irrelevant to infrasound for any biological organism.

Keywords: dimensional analysis, fluid-filled shell, scaling laws, viscoelastic, abdominal resonance, Buckingham Pi

*Corresponding author

Email address: `jmace@example.edu` (Jonathan Mace)

1. Introduction

The modal frequencies of fluid-filled viscoelastic shells depend on at least eleven physical parameters: geometric (semi-axes a, c; wall thickness h), material (elastic modulus E, Poisson's ratio ν, wall density ρ_w, loss tangent η), fluid (ρ_f, bulk modulus K_f), and loading (intra-abdominal pressure P_{iap}) [1, 2, 3, 4, 5, 6]. This high-dimensional parameter space makes comparison across species, organs, and loading conditions impractical without systematic dimensional reduction.

In a companion paper [7], we developed a fluid-filled oblate spheroidal shell model for the human abdominal cavity, predicting flexural resonances at 4 Hz to 10 Hz and a mechanical-to-airborne coupling ratio of approximately 6.6×10^4. Here we show that mammalian abdominal resonance is governed by a quasi-universal dimensionless frequency $\Pi_0 \approx 0.07$, implying that animal models are valid proxies for human studies and that the coupling ratio $\mathcal{R}$ is approximately size-independent. These results follow from a Buckingham Pi reduction of the eleven-parameter model to five governing dimensionless groups for flexural modes.

Three questions motivate this analysis. First, can animal models (rat, cat, pig) be used to study abdominal resonance phenomena that manifest in humans? The answer depends on whether the governing dimensionless groups—and hence $\mathcal{R}$—are approximately conserved across species. Second, can the breathing mode ($n = 0$)—whose frequency is dominated by the fluid bulk modulus—ever enter the infrasonic range for any biological organism? Third, which dimensionless groups dominate the flexural resonance, and how sensitive is f_2 to each?

2. Dimensional analysis

2.1. Governing variables

The natural frequency f_n of mode n depends on eleven variables: $f_n = f(E, a, c, h, \nu, \rho_w, \rho_f, K_f, P_{\mathrm{iap}}, \eta)$. With three fundamental dimensions (M, L,

T), the Buckingham Pi theorem [8] yields $11-3=8$ independent dimensionless groups (Table 1).

Table 1: Dimensionless groups from Buckingham Pi theorem. Groups marked $*$ drop out of the flexural-mode frequency.

Group	Definition	Name	Role
Π_0	$f_n\, a\sqrt{\rho_f/E}$	dimensionless frequency	output
Π_1	h/a	thickness ratio	stiffness & mass
Π_2	c/a	aspect ratio	geometry
Π_3	ρ_w/ρ_f	density ratio	added mass
Π_4^*	K_f/E	compressibility ratio	breathing mode only
Π_5	P_{iap}/E	prestress ratio	tension stiffening
Π_6	ν	Poisson's ratio	membrane coupling
Π_7^*	η	loss tangent	damping only

2.2. Reduction for flexural modes

For flexural modes ($n \geq 2$), two simplifications apply. The fluid bulk modulus K_f contributes no restoring force (the fluid acts as added mass only), so Π_4 drops out. The loss tangent η affects the quality factor but not the undamped frequency, so Π_7 drops out. The effective parameter count reduces to six: a dimensionless frequency output Π_0 governed by five input groups $(\Pi_1, \Pi_2, \Pi_3, \Pi_5, \Pi_6)$.

The scaling law takes the closed form

$$f_n = \sqrt{\frac{E}{\rho_f}}\,\frac{1}{a}\,\Phi_n\left(\frac{h}{a},\ \frac{c}{a},\ \frac{\rho_w}{\rho_f},\ \frac{P_{\mathrm{iap}}}{E},\ \nu\right), \tag{1}$$

where the $1/(2\pi)$ factor is absorbed into Φ_n, which admits a closed-form expression from the equivalent-sphere shell equations:

$$\Phi_n = \frac{1}{2\pi}\sqrt{\frac{K_{\mathrm{bend}} + K_{\mathrm{memb}} + K_{\mathrm{pre}}}{m_{\mathrm{eff}}}} \cdot a\sqrt{\frac{\rho_f}{E}}, \tag{2}$$

with

$$K_{\text{bend}} = n(n-1)(n+2)^2 \frac{D}{R^4}, \tag{3}$$

$$K_{\text{memb}} = \frac{Eh}{R^2}\lambda_n, \quad K_{\text{pre}} = \frac{P}{R}(n-1)(n+2), \tag{4}$$

$$m_{\text{eff}} = \rho_w h + \frac{\rho_f R}{n}, \tag{5}$$

where $D = Eh^3/[12(1-\nu^2)]$ is the flexural rigidity, $R = (a^2c)^{1/3}$ is the equivalent-sphere radius, and the membrane mode-shape factor is

$$\lambda_n = \frac{n^2+n-2+2\nu}{n^2+n+1-\nu}. \tag{6}$$

2.3. Sensitivity: which Π-group dominates?

For the $n = 2$ mode at canonical parameters ($h/a = 0.056$, $c/a = 0.67$, $\nu = 0.45$), the stiffness budget is dominated by the membrane term K_{memb}, which contributes approximately 78% of the total restoring force, with bending (K_{bend}, ∼14%) and prestress (K_{pre}, ∼8%) playing secondary roles. The transition from membrane-dominated to bending-dominated behaviour occurs near $h/a \approx 0.15$, well above the physiological range.

Consequently, f_2 is most sensitive to the thickness ratio $\Pi_1 = h/a$, which enters both the membrane stiffness (linearly through h) and the effective mass (linearly through $\rho_w h$). The density ratio $\Pi_3 = \rho_w/\rho_f$ has a more modest effect: because soft tissue and peritoneal fluid have similar densities ($\Pi_3 \approx 1.08$), the fluid added mass $\rho_f R/n$ dominates the denominator of m_{eff}. The prestress ratio $\Pi_5 = P_{\text{iap}}/E$ contributes less than 10% to the total stiffness at canonical values but becomes important for very soft tissue ($E < 0.05\,\text{MPa}$), where intra-abdominal pressure provides the dominant restoring force. This sensitivity structure explains why Π_0 is quasi-universal across mammals: the governing ratios h/a and ρ_w/ρ_f are constrained by shared developmental biology [9].

3. Parametric validation

We evaluate (1) over a 486-point grid spanning physiological ranges: $E \in [0.05, 2.0]$ MPa (6 values), $a \in [0.15, 0.20]$ m (3 values), $c/a \in [0.5, 0.8]$ (3 values),

$h \in [0.005, 0.015]$ m (3 values), and $\eta \in [0.15, 0.40]$ (3 values), with ρ_w, ρ_f, K_f, ν, and P_{iap} held at canonical values.

Figure 1 shows the result. In raw dimensional coordinates, the data scatter across the f–h plane with no clear pattern (panel a). Non-dimensionalisation via (1) collapses all 486 points onto universal curves $\Pi_0(h/a)$ parameterised by c/a (panel b). The parity plot (panel c) confirms that the analytical Φ_n reproduces every numerical point to within numerical round-off (maximum relative error 5.8×10^{-16}), confirming the algebraic consistency of the non-dimensionalisation rather than providing independent physical validation, which was established in the companion paper [7].



Figure 1: Dimensional collapse of the 486-point parametric study. *(a)* Raw dimensional data (f_2 vs. h) shows no clear pattern. *(b)* Non-dimensionalising via equation (1) collapses all points onto universal curves $\Pi_0(h/a)$ parameterised by c/a; shaded bands span the P_{iap}/E range, different markers denote different E. *(c)* Parity plot confirms the analytical Φ reproduces every numerical point to within numerical round-off, confirming algebraic consistency.

4. Cross-species scaling

The scaling law (1) predicts modal frequencies for any organism given its body dimensions and tissue properties. Table 2 lists predictions for four mammals, using allometric estimates from published biomechanical data [10, 9]. All dimensionless groups are shown explicitly so the reader can verify the quasi-constancy of the Π-groups across species. Poisson's ratio ($\nu = 0.45$) and the

prestress ratio ($P_{\mathrm{iap}}/E \approx 0.01$) are approximately constant across mammals and are omitted from the table for brevity.

Table 2: Cross-species scaling of the $n = 2$ flexural mode, showing all governing Π-groups. $\mathcal{R} = 1/(ka)^2$ quantifies the radiation-efficiency penalty for airborne excitation. Allometric estimates of E, h/a, and c/a are drawn from tissue-mechanics literature [10]; uncertainties in E propagate to ∼20% uncertainty in f_2 via the $\sqrt{E}$ dependence.

Species	a (cm)	E (MPa)	h/a	c/a	ρ_w/ρ_f	f_2 (Hz)	Π_0	ka	$\mathcal{R}$
Rat	3	0.07	0.047	0.63	1.06	15.7	0.067	7.7×10^{-3}	1.7×10^4
Cat	8	0.08	0.050	0.65	1.07	7.2	0.065	9.3×10^{-3}	1.2×10^4
Pig	15	0.09	0.053	0.67	1.08	4.5	0.069	1.1×10^{-2}	8.6×10^3
Human	18	0.10	0.056	0.67	1.08	3.9	0.072	1.1×10^{-2}	7.7×10^3

Three observations emerge. First, the dimensionless frequency Π_0 is approximately constant (0.065–0.072) across a six-fold range of body size, confirming that the governing Π-groups take similar values across mammalian species. Second, the coupling ratio $\mathcal{R}$ varies by only a factor of two (from 7.7×10^3 for humans to 1.7×10^4 for rats), remaining firmly in the 10^3–10^4 range. Airborne coupling is negligible for all species studied. Third, the product ka is of order 10^{-2} for all four species, placing every organism deep in the Rayleigh regime.

This size-independence of $\mathcal{R}$ has a practical implication: animal models are valid proxies for human abdominal resonance studies [11, 12], *provided excitation is delivered mechanically* (e.g. shaker table) rather than acoustically. Acoustic experiments would require compensating for the $(ka)^n$ coupling penalty, which varies with body size and frequency.

5. The breathing mode is never infrasonic

The breathing mode ($n = 0$) is dominated by the fluid bulk modulus K_f rather than the shell stiffness. Its frequency scales as $f_0 \propto \sqrt{K_f/\rho_f}/R$, which for water-like biological fluids ($K_f \approx 2.2\,\mathrm{GPa}$) gives $f_0 \approx 2490\,\mathrm{Hz}$ at the human abdominal scale. To bring f_0 down to $20\,\mathrm{Hz}$—the upper boundary of



Figure 2: Cross-species scaling. *Left:* Flexural mode f_2 vs. semi-major axis a. The dashed line shows pure $1/a$ scaling; deviations arise because smaller animals have softer tissue and different h/a. *Right:* Coupling ratio $\mathcal{R}$ is approximately size-independent, supporting the validity of animal-model extrapolation.

infrasound—would require an equivalent radius of order 410 m. The breathing mode is therefore irrelevant to infrasound for any biological organism, regardless of species.

6. Concluding remarks

Buckingham Pi analysis reduces the eleven-parameter fluid-filled viscoelastic shell model to five governing dimensionless groups for flexural modes. The resulting closed-form scaling law collapses a 486-point parametric study with algebraic consistency and provides cross-species predictions without re-solving the eigenvalue problem. The dimensionless frequency $\Pi_0 \approx 0.07$ is quasi-universal across mammals, and the coupling ratio $\mathcal{R} \sim 10^3$–10^4 is approximately size-independent, validating animal-model studies of abdominal resonance. The breathing mode never enters the infrasonic range for any organism.

The dimensionless framework developed here has applications beyond abdominal biomechanics. In blast-injury research, the scaling law enables direct extrapolation from animal-model data to human injury thresholds without repeating experiments at human scale. In medical ultrasound, fluid-filled organ models are routinely used to predict diagnostic and therapeutic responses; the

Π-group reduction offers a systematic basis for cross-organ and cross-species comparison [13, 14]. In vehicle NVH assessment, whole-body vibration standards [12] rely on empirical transfer functions whose mechanistic basis is provided by scaling laws of the type derived here. Finally, in underwater acoustics, fluid-filled elastic shells are canonical targets for sonar modelling [4, 3]; the present dimensional reduction applies directly to that class of problems.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

The analysis code and parametric data are available at `https://github.com/JonathanMace/brownnote`.

References

[1] W. Soedel, Vibrations of Shells and Plates, 3rd Edition, Marcel Dekker, 2004.

[2] A. W. Leissa, Vibration of Shells, NASA, 1973, nASA SP-288.

[3] M. C. Junger, D. Feit, Sound, Structures, and Their Interaction, 2nd Edition, MIT Press, 1986.

[4] M. C. Junger, Sound scattering by thin elastic shells, The Journal of the Acoustical Society of America 24 (4) (1952) 366–373. doi:10.1121/1.1906905.

[5] H. Lamb, On the vibrations of an elastic sphere, Proceedings of the London Mathematical Society s1-13 (1) (1882) 189–212. doi:10.1112/plms/s1-13.1.189.

[6] G. B. Warburton, The vibration of rectangular plates, Proceedings of the Institution of Mechanical Engineers 168 (1) (1976) 371–384, comparison of shell frequency predictions with experiment.

[7] J. Mace, Modal analysis of a fluid-filled viscoelastic oblate spheroidal shell: Implications for the existence of the "brown note", Journal of Sound and VibrationSubmitted (2026).

[8] E. Buckingham, On physically similar systems; illustrations of the use of dimensional equations, Physical Review 4 (4) (1914) 345–376. doi:10.1103/PhysRev.4.345.

[9] G. B. West, J. H. Brown, B. J. Enquist, A general model for the origin of allometric scaling laws in biology, Science 276 (5309) (1997) 122–126. doi:10.1126/science.276.5309.122.

[10] Y. C. Fung, Biomechanics: Mechanical Properties of Living Tissues, 2nd Edition, Springer-Verlag, 1993.

[11] H. E. von Gierke, Biodynamic response of the human body, Applied Mechanics Reviews 24 (1) (1971) 1–9.

[12] International Organization for Standardization, Mechanical vibration and shock — evaluation of human exposure to whole-body vibration — part 1: General requirements (1997).

[13] H. N. Abramson, The Dynamic Behavior of Liquids in Moving Containers, NASA, 1966, nASA SP-106.

[14] F. Fahy, P. Gardonio, Sound and Structural Vibration: Radiation, Transmission and Response, 2nd Edition, Academic Press, 2007.