

## *Hybrid Partial Least Squares*

Axiom [axiom]  
rem]Lemma [theorem]  
Fact Remark

BIOMETRICS

000,

000—

000

DOI:

**Hybrid**

000

000

**Par-**

0000

**tial**

**Least**

**Squares**

**Re-**

**gres-**

**sion**

**with**

**Mul-**

**ti-**

**ple**

**Func-**

**tional**

**and**

**Scalar**

**Pre-**

**dic-**

**tors**

re-

duc-

tion,

func-

**Keywords:**

di-

men-

sion

1. Introduction

Biomedical studies often collect multiple modal data, as in the Emory University renal study (Chang et al., 2020; Jang, 2021), which records multiple renogram curves (functional

tional data analysis, multiple time data modalities, multi-variate data analysis, multi-variate functional data, partial least square

and data)  
 scalar- along  
 valued with  
 i.i.d. scalar  
 co- side  
 vari- in-  
 ates for-  
 in ma-  
 a tion  
 lin- for  
 ear each  
 re- kid-  
 gres- ney.  
 sion Such  
 model: dis-

$$Y_i = \boldsymbol{\beta}^\top \mathbf{Z}_i + \sum_{k=1}^K \int_0^1 \beta_k(t) X_{ik}(t) dt + \epsilon_i,$$

re-  
 where lated  
 $Y_i$  phys-  
 is i-  
 a o-  
 scalar log-  
 re- i-  
 sponse, cal  
 $X_{i1}(t), \dots, X_{iK}(t) \in$  sig-  
 $\mathbb{L}^2[0, 1]$  nals  
 are can  
 func- be  
 tional rep-  
 pre- re-  
 dic- sented  
 tors, as  
 $\mathbf{Z}_i =$  functional-  
 $(Z_{i1}, \dots, Z_{ip})^\top$   
 is

dic- a  
 tors Eu-  
 are clidean  
 cen- vec-  
 tered, tor  
 so co-  
 the vari-  
 in- ate,  
 ter- and  
 cept  $\epsilon_i$   
 term de-  
 can notes  
 be ob-  
 omit- ser-  
 ted. va-  
 In tional  
 ad- noise.  
 di- For  
 tion no-  
 to ta-  
 the tional  
 ill- con-  
 posedness ve-  
 of nience,  
 the we  
 infinite- as-  
 dimensional sume  
 slope through-  
 func- out  
 tions that  
 $\beta_k(t)$  the  
 and re-  
 the sponses  
 high and  
 di- pre-

per  
ad-  
dresses  
all  
three  
is-  
sues  
in  
a  
uni-  
fied  
man-  
ner  
by  
in-  
tro-  
duc-  
ing  
a  
hy-  
brid  
par-  
tial  
least  
squares  
(PLS)  
re-  
gres-  
sion  
frame-  
work  
de-  
fined  
on  
a  
men-  
sion-  
al-  
ity  
of  
the  
pre-  
dic-  
tors  
( $K +$   
 $p$ ),  
a  
fur-  
ther  
chal-  
lenge  
comes  
from  
the  
strong  
cor-  
re-  
la-  
tions  
be-  
tween  
func-  
tional  
and  
scalar  
pre-  
dic-  
tors.  
This  
pa-

in- novel  
 clude hy-  
 ba- brid  
 sis Hilbert  
 ex- space.  
 pan-  
 sion, 1.1 *Previous*  
 us- *works*  
 ing *and*  
 power- *lim-*  
 series *i-*  
 (Gold- *ta-*  
 smith *tions*  
 et al., To  
 2011), ad-  
 B- dress  
 splines the  
 (Car- ill-  
 dot posedness  
 et al., stem-  
 2003; ming  
 Cai from  
 and the  
 Hall, infinite-  
 2006), dimensionality  
 or of  
 wavelets the  
 (Zhao func-  
 et al., tional  
 2012), com-  
 and po-  
 structure- nents,  
 aware com-  
 reg- mon  
 u- reme-  
 dies

rec- lar-  
tion iza-  
is tion,  
prin- such  
ci- as  
pal rough-  
com- ness  
po- penal-  
nent ties  
anal- For  
y- the  
sis high-  
(PCA) dimensionality,  
re- a  
gres- ma-  
sion, jor  
which so-  
ap- lu-  
plies tion  
PCA is  
sep- us-  
a- ing  
rately de-  
to rived  
the in-  
functional- puts.  
A  
(by, straight-  
for for-  
ex- ward  
am- ap-  
ple, proach  
[Happ](#) in  
[and](#) this  
[Greven](#) di-



ear (2018)  
re- and  
gres- scalar  
sion pre-  
with dic-  
func- tors,  
tional and  
pre- then  
dic- per-  
tors forms  
alone a  
(Hall clas-  
and si-  
Horowitz, cal  
2007; mul-  
Reiss ti-  
and vari-  
Og- ate  
den, re-  
2007; gres-  
Febrero- sion  
Bande on  
et al., the  
2017). com-  
How- bined  
ever, scores.  
since PCA  
the re-  
PCA- gres-  
derived sion  
in- is  
puts well-  
are studied  
not for  
in- lin-

In terms of pre-dic-tive power, par-tial least squares (PLS) re-gres-sion is a pow-er-ful al-ter-na-tive. It it-er-a-tively con-structs a set of or-formed by the re-sponse vari-able, they are not guar-an-teed to cap-ture the core re-gres-sion re-la-tion-ship with a small num-ber of de-rived in-puts.

## *Hybrid Partial Least Sq*

for thog-  
re- o-  
gres- nal  
sion. la-  
PLS tent  
for com-  
lin- po-  
ear nents  
re- from  
gres- the  
sion pre-  
in dic-  
the tors  
con- that  
text have  
of the  
func- max-  
tional i-  
pre- mal  
dic- co-  
tor vari-  
alone ance  
was with  
in- the  
tro- re-  
duced sponse  
by vari-  
[Preda](#) able,  
[and](#) and  
[Saporta](#) use  
([2005](#)) the  
for re-  
the sult-  
case ing  
of scores

the a  
 frame- sin-  
 work gle  
 by pre-  
 in- dic-  
 cor- tor.  
 po- Mo-  
 rat- ti-  
 ing vated  
 ba- by  
 sis re-  
 ap- gres-  
 prox- sion  
 i- on  
 ma- chemo-  
 tions met-  
 and ric  
 rough- spec-  
 ness tra,  
 penal- [Reiss](#)  
 ties [and](#)  
 to [Og-](#)  
 pro- [den](#)  
 mote [\(2007\)](#),  
 smooth- [Aguil-](#)  
 ness. [era](#)  
[De-](#) [et al.](#)  
[laigle](#) [\(2010\)](#),  
[and](#) [and](#)  
[Hall](#) [Aguil-](#)  
[\(2012\)](#) [era](#)  
 pro- [et al.](#)  
 vided [\(2016\)](#)  
 the ex-  
 first tended

o- thor-  
 nal- ough  
 iza- the-  
 tion. o-  
 For ret-  
 a i-  
 com- cal  
 pre- anal-  
 hen- y-  
 sive sis,  
 re- while  
 view, [Sari-](#)  
 see [cam](#)  
[Febrero-](#) [et al.](#)  
[Bande](#) (2022)  
[et al.](#) pro-  
 (2017). posed  
 More a  
 re- com-  
 cently, pu-  
[Beyaz-](#) ta-  
[tas](#) tion-  
[and](#) ally  
[Lin Shang](#) ef-  
 (2022) fi-  
 ex- cient  
 tended pro-  
 the ce-  
 frame- dure  
 work based  
 to on  
 ac- Golub-  
 com- Kahan  
 mo- bidi-  
 date ag-

which can lead to multi-collinearity and sub-optimal predictive performance. Multivariate modal correlations have mostly been addressed within an unsu- mul- ti- ple func- tional pre- dic- tors. However, these ex- ist- ing ap- proaches over- look the po- ten- tially strong cor- re- la- tions be- tween func- tional and scalar com- po- nents,

data, while [Geng et al. \(2020\)](#) considered joint pre-cision matrix estimation for brain measurements and confounding scalar co-variate. [Jang \(2021\)](#) proposed per-vised learning framework. For instance, [Kolar et al. \(2014\)](#) studied the estimation of joint undirected graphical models for functional and vector

re-  
 sponse  
 and  
 may  
 fail  
 to  
 cap-  
 ture  
 cor-  
 re-  
 la-  
 tion  
 with  
 the  
 out-  
 come.

a  
 joint  
 PCA  
 method  
 that  
 ac-  
 counts  
 for  
 cor-  
 re-  
 la-  
 tions  
 be-  
 tween  
 func-  
 tional  
 and

1.2 *Our*  
*con-*  
*tri-*  
*bu-*  
*tions*

scalar  
 data.  
 How-  
 ever,  
 the

To  
 ad-  
 dress  
 the  
 gaps  
 men-  
 tioned  
 above,  
 we  
 unify  
 the  
 reme-  
 dies  
 pro-

re-  
 sult-  
 ing  
 com-  
 po-  
 nents  
 are  
 still  
 not  
 in-  
 formed  
 by  
 the



## *Hybrid Partial Least Squares*

pro- posed  
pose for  
a these  
hy- three  
brid is-  
PLS sues:  
re- ba-  
gres- sis  
sion ex-  
frame- pan-  
work sion,  
that par-  
in- tial  
te- least  
grates squares,  
func- and  
tional ac-  
and count-  
vec- ing  
tor for  
pre- cor-  
dic- re-  
tors la-  
in tions  
a be-  
prin- tween  
ci- the  
pled func-  
and tional  
co- and  
her- scalar  
ent com-  
man- po-  
ner. nents.  
To We

and extract pre-dic-tive struc-ture from these jointly ob-served and po-tentially cor-re-lated data types, we de-fine a Hilbert space that treats the tu-ple of func-tional

is it-  
read-er-  
ily a-  
ap-tively  
pli-max-  
ca-i-  
ble miz-  
to ing  
dense the  
and em-  
ir-pir-  
reg-i-  
u-cal  
lar co-  
func-vari-  
tional ance  
data with  
and the  
sup-re-  
ports sponse,  
reg-sub-  
u-ject  
lar-to  
iza-a  
tion unit-  
tech-norm  
niques con-  
to straint  
pre-in  
vent this  
over-Hilbert  
fit-space.  
ting Our  
and frame-  
re-work

**2. Background**

on partial least squares and its extension to hybrid predictors

duce variance. We also provide the mathematical identities that

For intuition, let us return to the high-dimensional Euclidean predictor set

justify our algorithm.

vector  $Y_i = (\hat{\rho}_1^{[1]}, \dots, \hat{\rho}_1^{[L]})^\top \in \mathbb{R}^L$ .  $\mathbf{S}^\top \mathbf{Z}_i + \epsilon_i$ . To A common way to address ill-posedness and correlation, the  $l$ -th PLS is to approximate  $\hat{\boldsymbol{\xi}}_l$  by  $\mathbf{Z}_i$  using a low-dimensional where the two  $\widehat{\text{Cov}}^2$  using a low-dimensional cross-

tial covariance  
least and  
squares sam-  
(NI- ple  
PALS) co-  
is vari-  
pre- ance,  
sented re-  
in spec-  
Al- tively.  
go- A  
rithm stan-  
1. dard  
al-  
To go-  
ex- rithm  
tend for  
Al- solv-  
go- ing  
rithm this  
1 op-  
to ti-  
the miza-  
hy- tion  
brid prob-  
re- lem,  
gres- called  
sion non-  
model lin-  
(1), ear  
we it-  
must er-  
com- a-  
pute tive  
hy- par-

brid  
ob-  
jects  
in  
both  
the  
PLS  
di-  
rec-  
tion  
es-  
ti-  
ma-  
tion  
and  
pre-  
dic-  
tor  
resid-  
u-  
al-  
iza-  
tion  
steps  
(lines  
4  
and  
15).  
These  
com-  
pu-  
ta-  
tions  
can-  
not

# Algorithm

1

Scalar

par-

tial

least

squares

re-

gres-

sion

1:

Stan-

dard-

ize

each

$\mathbf{Z}_1, \dots, \mathbf{Z}_n$

so

that

each

fea-

ture

have

mean

zero

and

vari-

ance

one.

Stan-

dard-

ize

$Y_1, \dots, Y_n$ .

2:

**for**

$l =$

$1, 2, \dots, L$

**do**

3:

**PLS**

**di-**

**rec-**

**tion**

**and**

**score**

**es-**

**ti-**

**ma-**

**tion:**

4:

$\hat{\boldsymbol{\xi}}^{[l]} \leftarrow$

$\arg \max_{\boldsymbol{\alpha}} \widehat{\text{Cov}}^2(\{$

1

$\triangleright$

PLS

di-

rec-

tion

5:

$\hat{\rho}_i^{[l]} \leftarrow$

$\langle \hat{\boldsymbol{\xi}}^{[l]}, \mathbf{Z}_i^{[l]} \rangle, i =$

$1, \dots, n$

$\triangleright$

PLS

score

6:

**Residualization:**

7:

$\nu^{[l]} \leftarrow$

$\frac{\sum_{i=1}^n Y_i^{[l]} \hat{\rho}_i^{[l]}}{\sum_{i=1}^n \hat{\rho}_i^{[l]^2}}$

$\triangleright$

tion be  
 points sep-  
 $t$  a-  
 to rated  
 lines into  
 4 func-  
 and tional  
 15, and  
 treat- scalar  
 ing parts,  
 as  
 $(X_{i1}(t), \dots, X_{iK}(t), Z_{i1}, \dots, Z_{ip}) \in \mathbb{R}$   
 their  
 as cor-  
 a re-  
 Eu- la-  
 clidean tion  
 pre- must  
 dic- be  
 tor, ad-  
 ap- dressed.  
 ply- A  
 ing naive  
 steps ap-  
 4 proach  
 and is  
 15, to  
 and add  
 ag- loops  
 gre- over  
 gat- the  
 ing ob-  
 the served  
 re- eval-  
 sults. u-  
 This a-



## *Hybrid Partial Least Squares*

vari- point-  
abil- wise  
ity method,  
across how-  
the ever,  
do- is  
main, com-  
re- pu-  
sult- ta-  
ing tion-  
in ally  
over- pro-  
fit- hibitive  
ting for  
and densely  
un- ob-  
sta- served  
ble func-  
pre- tional  
dic- pre-  
tions. dic-  
The tors,  
main in-  
chal- fea-  
lenge si-  
of ble  
ex- for  
ten- ir-  
sion reg-  
to u-  
the lar  
hy- data,  
brid and  
set- prone  
ting to

size lies  
is in  
small the  
com- fol-  
pared low-  
to ing  
the points.  
di- First,  
men- in-  
sion de-  
and pen-  
num- dent  
ber vari-  
of ables  
func- con-  
tional sist  
and of  
scalar mul-  
pre- ti-  
dic- ple  
tors. highly  
Third, struc-  
ex- tured  
ist- im-  
ing ages  
par- and  
tial scalar  
least pre-  
squares dic-  
(PLS) tors.  
meth- Sec-  
ods ond,  
can our  
only sam-  
ac- ple

et al., 2017; Beyaztas and Shang, 2020); or (ii) a uni-variate functional pre-dictor with other scalar pre-dictors (Wang, 2018). The next section proposes a new ex-

com-mo-date (i) uni-variate or mul-ti-variate functional pre-dictors with-out any scalar pre-dictors (Preda and Saporta, 2005; De-laigle and Hall, 2012; Febrero-Bande

are ten-  
arith- sion  
metic frame-  
op- work  
er- that  
a- ad-  
tions dresses  
in these  
an is-  
in- sues.

ner  
prod-  
uct  
space.

We  
there-  
fore

### 3. Proposed PLS Al- go- rithm

for- Our  
mally new  
de- frame-  
fine work  
the builds  
Hilbert on  
space the  
of fact  
hy- that  
brid all  
ran- com-  
dom pu-  
pre- ta-  
dic- tions  
tors in  
along Al-  
with go-  
its rithm

Our  
new  
frame-  
work  
builds  
on  
the  
fact  
that  
all  
com-  
pu-  
ta-  
tions  
in  
Al-  
go-  
rithm

defined as the Cartesian product of  $K$  copies of the space of square-integrable functions on  $[0, 1]$  and the  $p$ -dimensional Euclidean space:

$$\mathbb{H} := (\mathbb{L}^2[0, 1])^K \times \mathbb{R}^p.$$

An element  $h \in$

corresponding to the complete inner product, and leverage them to extend Algorithm 1.

definition

[  
•  
Hybrid  
space]  
Let  
 $\mathbb{H}$   
be  
a  
product  
space

ca-  $\mathbb{H}$   
tion. is  
We an  
de- or-  
fine dered  
an tu-  
in- ple  
ner  $h =$   
prod-  $(f_1, \dots, f_K, \mathbf{u})$ ,  
uct where  
on  $f_k \in$   
 $\mathbb{H}$   $\mathbb{L}^2[0, 1]$   
for for  
any  $k =$   
two  $1, \dots, K$   
el- and  
e-  $\mathbf{u} \in$   
ments  $\mathbb{R}^p$ .  
 $h_1 =$  This  
 $(f_1, \dots, f_K, \mathbf{u})$  space  
and is  
 $h_2 =$  equipped  
 $(g_1, \dots, g_K, \mathbf{v})$  with  
as: element-

$$\langle h_1, h_2 \rangle_{\mathbb{H}} := \sum_{k=1}^K \int_{\text{vec-}}^{\text{wise}} f_k(t) g_k(t) dt + \omega \mathbf{u}^T \mathbf{v}$$

where tor  
 $\omega$  ad-  
is di-  
a tion  
and  
pos- scalar  
i- mul-  
tive ti-  
con- pli-  
stant

is a positive weight that needs to be pre-specified or estimated. It is mainly used to take into account heterogeneous behavior between functional and scalar

$(\omega > 0)$ . The inner product induces a norm  $\|\cdot\|_{\mathbb{H}}$  on the space, defined as  $\|h\|_{\mathbb{H}} := \langle h, h \rangle_{\mathbb{H}}^{1/2}$ , and a corresponding inner metric  $d(h_1, h_2) = \|h_1 - h_2\|_{\mathbb{H}}$ .

In (3),  $\omega$

tion, parts  
 all in  
 the terms  
 fol- of  
 low- mea-  
 ing sure-  
 the- ment  
 o- scale  
 ret- and/or  
 i- amount  
 cal of  
 re- vari-  
 sults a-  
 will tion  
 be (see  
 de- Sec-  
 rived tion  
 for [3.2](#)).  
 $\omega =$  With-  
 1. out  
 The loss  
 re- of  
 sults gen-  
 re- er-  
 main al-  
 valid ity  
 for and  
 any for  
 pos- the  
 i- clar-  
 tive ity  
 weights. of

lemma il-  
 lus-  
 tra-

[



Pre-  
dic-  
tor]

For  
the  
Hilbert  
space  
 $\mathbb{H}$   
de-  
fined  
in  
Def-  
i-  
ni-  
tion  
[3.1](#),  
The  
Borel  
 $\sigma$ -  
field  
on  
 $\mathbb{H}$ ,  
de-  
noted  
 $\mathfrak{B}(\mathbb{H})$ ,  
is  
the  
small-  
est  
 $\sigma$ -  
field  
con-  
tain-  
ing  
the

•  
Hy-  
brid  
Hilbert  
space]  
The  
hy-  
brid  
space  
 $\mathbb{H}$   
is  
a  
sep-  
a-  
ra-  
ble  
Hilbert  
space.  
  
Proof  
of  
Lemma  
[3.2](#)  
is  
pro-  
vided  
in  
Ap-  
pendix  
[??](#).  
  
definition  
  
[  
•  
Hy-  
brid

ing class  
 and  $\mathfrak{M}$   
 Eu- of  
 bank all  
 2015). sets  
 A of  
 hy- the  
 brid form  
 pre-  $\{q \in$   
 dic-  $\mathbb{H} \mid$   
 tor  $\langle q, h \rangle \in$   
 $W_i = O\},$   
 $(X_{i1}(t), \dots, X_{iK}(\mathfrak{t}), \mathbf{Z}_i)$   
 is any  
 a  $h \in$   
 mea-  $\mathbb{H}$   
 sur- and  
 able any  
 map- open  
 ping sub-  
 from set  
 a  $O \subseteq$   
 prob-  $\mathbb{R}$   
 a- (de-  
 bil- tails  
 ity can  
 space be  
 $(\Omega, \mathfrak{F}, P)$  found  
 into in  
 $(\mathbb{H}, \mathfrak{B}(\mathbb{H})).$  The-  
 o-  
 Then rem  
 the 7.1.1  
 joint of  
 re- [Hs-](#)  
 gres-

4)

	<i>Hybrid Partial Least Squares</i>
to	sion
a	model
dif-	(1)
fer-	can
ent	be
Hilbert	con-
space,	cisely
pos-	writ-
si-	ten
bly	as
de-	
fined	$Y_i = \langle \beta, W_i \rangle_{\mathbb{H}} + \epsilon_i$ , wh
over	
a	remark
dis-	
tinct	<b>O</b>
com-	.
pact	ur
do-	method
main	is
in	ap-
$\mathbb{R}^d$ ,	pli-
for	ca-
ar-	ble
bi-	to
trary	set-
$d$ ,	tings
and	where
ob-	each
served	func-
at	tional
dif-	pre-
fer-	dic-
ent	tor
time	be-
	longs

provides an efficient and robust means of producing PLS component scores in the presence of multiple dense and/or irregular functional points. However, for non-tational simplicity, our discussion assumes a common Hilbert space over the domain  $[0, 1]$  for all functional predictors. Our approach

## *Hybrid Partial Least Squares*

exploit structural relationships within and between the functions to avoid overfitting of the PLS components and to improve the generalizability of the model. This is achieved by combining the strengths of PLS and Ridge Regression. Ridge Regression is a statistical method that adds a penalty term to the least squares regression equation, which helps to reduce the variance of the coefficient estimates and thus prevents overfitting. By combining the two methods, the Hybrid PLS method can leverage the predictive power of PLS while also benefiting from the regularization properties of Ridge Regression. This hybrid approach is particularly useful in situations where the data is noisy or the number of predictors is large relative to the number of observations. The resulting model is more robust and generalizable than either PLS or Ridge Regression alone. The Hybrid PLS method is implemented in the following R code snippet:

tion  
of  
smoothed  
PLS  
com-  
po-  
nents  
and  
or-  
thog-  
o-  
nal-  
iza-  
tion,  
de-  
tailed  
in  
Sec-  
tions 3.3.1  
and 3.3.2,  
re-  
spec-  
tively.  
Af-  
ter  
a  
suit-  
able  
num-  
ber  
of  
it-  
er-  
a-  
tions,  
ity  
and  
in-  
ter-  
pretabil-  
ity  
of  
the  
pre-  
dic-  
tive  
model.  
Each  
it-  
er-  
a-  
tion  
of  
our  
ap-  
proach  
con-  
sists  
of  
two  
sub-  
rou-  
tines:  
reg-  
u-  
lar-  
ized  
es-  
ti-  
ma-

denotes the hybrid regression coefficient is estimated, as described in Section 3.4. For notational simplicity, we omit the iteration operation in the following discussion, with the understanding that the sub-routines apply to any iteration. The complete algorithm

lin- is  
 early sum-  
 in- ma-  
 de- rized  
 pen- in  
 dent, Al-  
 for go-  
 ex- rithm 2.  
 am-  
 ple,

### 3.1 *Preliminary*

cu- *step*  
 bic *l*:  
 B- *finite-*  
 splines, *basis*  
 the *ap-*  
 Fourier *prox-*  
 ba- *i-*  
 sis, *ma-*  
 or *tion*

an Let  
 or-  $b_m(t)$   
 thonor- be  
 mal a  
 poly- twice-  
 no- differentiable  
 mial ba-  
 ba- sis  
 sis of  
 of  $\mathbb{L}^2([0, 1])$   
 de- whose  
 gree sec-  
 greater ond  
 than deriva-  
 three. tives  
 Us- are  
 also



co- ing  
 ef- this  
 fi- ba-  
 cient sis,  
 (with the  
 it-  $j$ th  
 er- func-  
 a- tional  
 tion pre-  
 in- dic-  
 dices tor,  
 sup- re-  
 pressed) gres-  
 are sion  
 rep- co-  
 re- ef-  
 sented fi-  
 as cient,  
 fol- PLS  
 lows: com-

$$X_{ij}(t) = \sum_{m=1}^{\infty} \theta_{ijm} \underset{\text{po-}}{\overset{\text{nent}}{b_m}}(t), \beta_j(t) = \sum_{m=1}^{\infty} \beta_{jm} \underset{\text{po-}}{\overset{\text{nent}}{b_m}}(t).$$

In di-  
 prac- rec-  
 tice, tion,  
 the and  
 full or-  
 set thog-  
 of o-  
 co- nal-  
 ef- iza-  
 fi- tion  
 cients re-  
 can gres-  
 not sion

mod- be  
er- ob-  
ately tained  
large with  
 $M$  fi-  
(e.g., nite  
15 sam-  
or ple  
20) size,  
to as  
cap- func-  
ture tional  
func- data  
tional are  
vari- mea-  
a- sured  
tion on  
with- a  
out fi-  
fine- nite  
tuning, grid.  
as Thus,  
smooth- we  
ness trun-  
is cate  
han- the  
dled ex-  
via pan-  
pe- sion  
nal- at  
iza-  $M$   
tion terms.  
(see We  
Sec- choose  
tion 3.3.1). a

rec-  
tion  
and  
or-  
thog-  
o-  
nal-  
iza-  
tion  
re-  
gres-  
sion  
co-  
ef-  
fi-  
cient  
to  
ad-  
mit  
the

The  
trun-  
cated  
ex-  
pan-  
sions  
of  
the  
pre-  
dic-  
tor  
and  
co-  
ef-  
fi-  
cient  
are  
de-  
noted  
as

fol-  
low-  
ing  
ex-  
pan-  
sion:

$$\tilde{X}_{ij}(t) := \sum_{m=1}^M \theta_{ijm} b_m$$

and  
our  
sug-  
gest

$$\xi_j(t) = \sum_{m=1}^M \gamma_{jm} b_m(t), \quad \delta_j(t) = \sum_{m=1}^M \tau_{jm} b_m(t)$$

This  
im-  
plies  
that  
all  
com-  
pu-

method  
re-  
stricts  
each  
PLS  
com-  
po-  
nent  
di-

tu- ta-  
ple tions

$\widetilde{W}_i := (\widetilde{X}_{i1}, \dots, \widetilde{X}_{iK}, \mathbf{Z}_i)$ .  
in  
this

Let pa-

$\boldsymbol{\theta}_{ij}$ , per

$\boldsymbol{\eta}_j$ , are

$\boldsymbol{\gamma}_j$ , car-

and ried

$\boldsymbol{\pi}_j$  out

de- en-

note tirely

the within

$M$ - the

dimensional sub-

vec- space

tors

$\widetilde{\mathbb{H}} := \text{span}(b_1(t), \dots,$

of

6)

co-

The

ef-

$i$ th

fi-

hy-

cients:

brid

$\boldsymbol{\theta}_{ij} := (\theta_{ij1}, \dots, \theta_{ijM})^\top$ ,  $\boldsymbol{\eta}_j := (\eta_{j1}, \dots,$

dic-

For

tor,

the

pro-

pre-

jected

dic-

on

tors,

$\widetilde{\mathbb{H}}$ ,

we

is

stack

rep-

the

re-

co-

sented

ef-

by

fi-

the

as coefficient

$\mathbf{y} :=$  vec-

$(y_1, \dots, y_n)^\top$ . tors

Let across

$B, B'' \in$  ob-

$\mathbb{R}^{M \times M}$  ser-

de- va-

note tions

the into

Gram the

ma- ma-

tri- trix

ces

of  $\Theta_j := (\boldsymbol{\theta}_{1j}, \dots, \boldsymbol{\theta}_{nj})^\top$

8) the and

ba- con-

sis struct

func- the

tions full

and de-

their sign

sec- ma-

ond trix

deriva-

tives,  $\Theta := (\Theta_1, \dots, \Theta_K, \mathbf{Z})$

9) with

en- Let

tries us

$B_{m,m'} := \int_0^1 b_m(t) b_{m'}(t) dt,$  de-  
note  $B''_{m,m'}$

for the

$m, m' =$  re-

$1, \dots, M.$  sponse

We vec-

then tor

the de-  
tu- fine  
ple the

$(\mathbb{B}, \mathbb{B}'', \Theta, \mathbf{y}) \in \mathbb{R}^{\text{block-}(MK+p) \times (MK+p) \times \mathbb{R}^{(l)}}$   
diagonal

with ma-  
the tri-  
in- ces

dex  $\mathbb{B} := \text{blkdiag}(B, \dots, I)$

1)  $l$

omit- Then

ted the  
for full  
brevity. data

remark for

the  
hy-  
W brid

• PLS

hile prob-  
dif- lem

fer- at  
ent the

bases  $l$ -

could th

be it-

used er-

for a-

each tion

func- can

tional be

pre- rep-

dic- re-

tor, sented

we by

adopt

### 3.2 Preliminary

*step*

2

:

*data*

*pre-*

*pro-*

*cess-*

*ing*

Functional

and

scalar

el-

e-

ments

of

the

hy-

brid

pre-

dic-

tors

of-

ten

have

in-

com-

pat-

i-

ble

units

and/or

ex-

hibit

dif-

a com-

mon

ba-

sis

for

sim-

plic-

ity.

The

def-

i-

ni-

tions

of

$\mathbb{B}$

and

$\mathbb{B}''$

re-

main

gen-

eral

enough

to

ac-

com-

mo-

date

dis-

tinct

bases

if

needed.

chance of contributing to the predictor/response structure; and ii) a predictor with high correlation to  $Y$  but relatively low variance may be different amounts of variation. This can be problematic for our PLS framework which is not scale invariant as:

i) each predictor has different



The first step is to account for discrepancies *within* respective functional and scalar parts, if needed. If the functional parts  $\tilde{X}_{i1}(t), \dots, \tilde{X}_{iK}(t)$  are measured in different over-looked. To obtain PLS components that have a meaningful interpretation, we standardize the predictor data via the following steps.

dic- ent  
 tors units  
 $\mathbf{Z}_i =$  or  
 $(Z_{i1}, \dots, Z_{ip})^\top$  have  
 ex- quite  
 hibit dif-  
 dif- fer-  
 fer- ent  
 ent do-  
 amounts mains,  
 of one  
 vari- can  
 a- stan-  
 tion, dard-  
 one ize  
 can them  
 stan- to  
 dard- have  
 ize mean  
 them zero  
 to and  
 have in-  
 mean te-  
 zero grated  
 and vari-  
 unit ance  
 vari- of  
 ance. one.  
 The If  
 sec- mul-  
 ond ti-  
 step vari-  
 is ate  
 to scalar  
 elim- pre-

## Hybrid Partial Least Squares

product (3) that ensures functional and vector parts have comparable variance. A sensible data-driven approach to choosing an appropriate  $\omega$  in the hybrid inner

the weight

hy- is

brid to

ob- set

ject  $\omega = \frac{\sum_{i=1}^n \sum_{k=1}^K \|\tilde{X}_i\|}{\sum_{i=1}^n \|\mathbf{Z}_i\|}$

as

$\widetilde{W}_i =$  In

$(\tilde{X}_{i1}(t), \dots, \tilde{X}_{iK}(t)) \omega^{1/2} \mathbf{Z}_i$ ,

whose tice,

vec- in-

tor stead

part of

has us-

been ing

scaled in-

by ner

a prod-

fac- uct

tor weighted

of by

$\omega^{1/2}$ .  $\omega^{1/2}$ ,

one

### 3.3 *Iterative*

can

*steps*

im-

The ple-

it- ment

er- this

a- weight-

tive ing

pro- scheme

cess by

pre- for-

sented mu-

here lat-

yields ing

an

### Hybrid Partial Least Sq

tion of the PLS component direction (Section 3.3.1) and residualization (Section 3.3.2). The proofs are deferred to Appendix ??.

or-  
thonor-  
mal  
hy-  
brid  
ba-  
sis  
that  
ef-  
fec-  
tively  
cap-  
tures  
the  
predictor-  
response  
re-  
la-  
tion-  
ships.  
It  
pro-  
ceeds  
through  
two  
in-  
ter-  
me-  
di-  
ate  
steps:  
the  
es-  
ti-  
ma-

*nent* of  
*di-* the  
*rec-* re-  
*tion.* sult-  
 ing  
 We es-  
 be- ti-  
 gin mates  
 by are  
 for- in-  
 mally tro-  
 in- duced  
 tro- in  
 duc- Sec-  
 ing tion  
 the 4.  
 core

op- 3.3.1  
 ti- *It-*  
 miza- *er-*  
 tion *a-*  
 prob- *tive*  
 lem *step*  
 per- *l:*  
 ti- *reg-*  
 nent *u-*  
 to *lar-*  
 the *ized*  
 PLS *es-*  
 di- *ti-*  
 rec- *ma-*  
 tion *tion*  
 es- *of*  
 ti- *PLS*  
 ma- *com-*  
*po-*

PLS regression, which is formulated as a generalized Rayleigh quotient (Proposition 3.4). Building upon this foundational concept, we present our regularized component for simulating as a generalized Rayleigh quotient (Proposition 3.5). Furthermore, we detail an efficient computational

rec- scheme  
tion. (Propo-  
It si-  
fully tion  
lever- 3.6).  
ages  
the *Core*  
con- *op-*  
tin- *ti-*  
u- *miza-*  
ous *tion*  
na- *prob-*  
ture *lem*. We  
of present  
the the  
func- core  
tional op-  
com- ti-  
po- miza-  
nents tion  
and prob-  
the lem  
function- that  
scalar di-  
hy- rectly  
brid es-  
struc- ti-  
ture. mates  
We the  
de- hy-  
scribe brid  
the PLS  
strat- com-  
egy po-  
at nent  
di-



the  
 $l$ -  
 th  
 it-  
 er-  
 a-  
 tion.  
 The  
 PLS  
 com-  
 po-  
 nent  
 di-  
 rec-  
 tion  
 is  
 es-  
 ti-  
 mated  
 by  
 the

$$\langle \widetilde{W}_1, \xi \rangle_{\mathbb{H}}, \dots, \langle \widetilde{W}_{\text{unif}}, \xi \rangle_{\mathbb{H}}$$

and norm  
 the di-  
 rec-  
 tion  
 $y_1, \dots, y_n,$   $\xi^{[l]} \in$   
 $\widetilde{\mathbb{H}}$   
 that  
 as max-  
 i-

$$\widehat{\text{Cov}}(\langle \widetilde{W}, \xi \rangle_{\mathbb{H}}, Y) \stackrel{\text{the}}{:=} \frac{1}{n} \sum_{i=1}^n y_i \langle \widetilde{W}_i, \xi \rangle_{\mathbb{H}}.$$

squared  
 We  
 em-  
 de-

tain-  
ing  
these  
co-  
ef-  
fi-  
cients  
is  
equiv-  
a-  
lent  
to  
solv-  
ing  
the  
max-  
i-  
miza-  
tion  
prob-  
lem  
(14).  
The  
fol-  
low-  
ing  
propo-  
si-  
tion  
for-  
mu-  
lates  
this  
co-  
ef-

note  
the  
es-  
ti-  
mated  
PLS  
com-  
po-  
nent  
di-  
rec-  
tion  
as

$$\hat{\xi} := \arg \max_{\xi \in \tilde{\mathbb{H}}} \widehat{\text{Cov}}^2(\langle \mathbf{V}, \xi \rangle)$$

Here,

$$\hat{\xi} \in \tilde{\mathbb{H}}$$

is  
an  
or-  
dered  
pair  
ex-  
panded

as:

$$\hat{\xi} = (\hat{\xi}_1(t), \dots, \hat{\xi}_K(t),$$

where

$$\hat{\zeta}$$

is  
the  
scalar

part.

Ob-

iteration of the PLS algorithm, the covariance matrix of the squared multiple correlation coefficients of the squared co-vari-  
 ance maximizer defined in (14), is obtained as

following procedure as a generalized Rayleigh quotient:

# Proposition

1.

Let  $(\mathbb{B}, \mathbb{B}'', \Theta, \mathbf{y})$  denote the observed data defined in (12).

At the  $l$ -th

$\left(\hat{\gamma}_{11}, \dots, \hat{\gamma}_{1M}, \dots, \hat{\gamma}_{K1}, \dots, \hat{\gamma}_{KM}, \hat{\boldsymbol{\zeta}}^\top\right)$

of-  
fers  
an  
ef-  
fi-  
cient  
way  
to  
es-  
ti-  
mate  
the  
PLS  
com-  
po-  
nent  
di-  
rec-  
tion  
 $\hat{\xi}$ ,  
its  
func-  
tional  
com-  
po-  
nents,  
 $\hat{\xi}_1, \dots, \hat{\xi}_K$ ,  
may  
not  
be  
smooth.  
which  
com-  
pli-  
cates

where

$$V := \frac{1}{n^2} (\mathbb{B}\Theta^\top \mathbf{y})(\mathbb{B}\Theta$$

The

proof  
of  
Propo-  
si-  
tion  
[3.4](#)  
is  
pro-  
vided  
in  
Ap-  
pendix  
??.

*Proposed*  
*reg-*  
*u-*  
*lar-*  
*ized*  
*es-*  
*ti-*  
*ma-*  
*tion*  
*pro-*  
*ce-*  
*dure*. Al-  
though  
Propo-  
si-  
tion  
[3.4](#)

balances predictive performance with smoothness. Specifically, we penalize the roughness of each  $\hat{\xi}_j$  using its integrated squared second derivative in-  
 terpretation and can lead to overfitting and unstable predictions. To address this, we propose a regularized extension that

con-  
trol  
the  
com-  
plex-  
ity  
of  
the  
es-  
ti-  
mated  
func-  
tional  
com-  
po-  
nents.  
One  
pos-  
si-  
ble  
ap-  
proach  
is  
to  
ex-  
tend  
the  
smoothed  
func-  
tional  
PCA  
frame-  
work  
of  
[Rice](#)

tive

$$\text{PEN}(\hat{\xi}_j) := \int_0^1 \{\hat{\xi}_j''(t)$$

Instead

of  
solely  
max-  
i-  
miz-  
ing  
the  
squared  
em-  
pir-  
i-  
cal  
co-  
vari-  
ance  
 $\widehat{\text{Cov}}^2(\langle \widetilde{W}, \xi \rangle_{\mathbb{H}}, Y)$ ,  
we  
in-  
cor-  
po-  
rate  
this  
rough-  
ness  
penalty  
to  
si-  
mul-  
ta-  
ne-  
ously

max-  
i-  
miz-  
ing  
co-  
vari-  
ance  
and  
pe-  
nal-  
iz-  
ing  
rough-  
ness.  
How-  
ever,  
this  
ap-  
proach  
Rice  
and  
Sil-  
ver-  
man  
(1991)  
as-  
sumes  
that  
the  
func-  
tional  
pre-  
dic-  
tors  
ad-

and  
Sil-  
ver-  
man  
(1991)  
by  
mod-  
i-  
fy-  
ing  
the  
ob-  
jec-  
tive  
in  
(14)  
to

$$\widehat{\text{Cov}}^2\left(\langle \widetilde{W}, \xi \rangle_{\mathbb{H}}, Y\right) - \sum_{j=1}^K$$

where  
the  
smooth-  
ing  
pa-  
ram-  
e-  
ters  
 $\{\lambda_j\}_{j=1}^K$   
con-  
trol  
the  
trade-  
off  
be-  
tween

strat- mit  
 egy an  
 of or-  
[Sil-](#) thog-  
[ver-](#) o-  
[man](#) nal  
 (1996), ex-  
 which pan-  
 re- sion  
 places in  
 the the  
 stan-  $\mathbb{L}^2$   
 dard sense.

or- To  
 thonor- avoid  
 mal- the  
 ity or-  
 con- thog-  
 straint o-  
 with nal  
 a ba-  
 weaker sis  
 one as-  
 based sump-  
 on tion  
 a of  
 mod- [Rice](#)  
 i- [and](#)  
 fied [Sil-](#)  
 in- [ver-](#)  
 ner [man](#)  
 prod- (1991),  
 uct we  
 that adopt  
 in- the



ing the observed serv-  
 va- tions have been resid-  
 u- al- ized in pre-  
 vi- ous steps, solves the  
 fol- low- ing op-  
 ti- miza- tion prob-  
 lem:

$$\hat{\xi} := \arg \max_{\xi \in \widetilde{\mathbb{H}}} \widehat{\text{Cov}}_{\text{dex}}^2(\langle \widetilde{W}, \xi \rangle_{\mathbb{H}}, Y) \text{ s.t. } \|\xi\|_{\mathbb{H}} \leq 1$$

Here,  $\hat{\xi} \in \widetilde{\mathbb{H}}$  is an omit-  
 ted and as-  
 sum-

tain-  
 ing  
 these  
 co-  
 ef-  
 fi-  
 cients  
 is  
 equiv-  
 a-  
 lent  
 to  
 solv-  
 ing  
 the  
 max-  
 i-  
 miza-  
 tion  
 prob-  
 lem  
 (18).  
 The  
 fol-  
 low-  
 ing  
 propo-  
 si-  
 tion  
 for-  
 mu-  
 lates  
 this  
 co-  
 ef-

or-  
 dered  
 pair  
 ex-  
 panded  
 as:  

$$\hat{\xi} = (\hat{\xi}_1(t), \dots, \hat{\xi}_K(t),$$
 where  

$$\hat{\zeta}$$
 is  
 the  
 scalar  
 part.  
 This  
 for-  
 mu-  
 la-  
 tion  
 max-  
 i-  
 mizes  
 the  
 squared  
 co-  
 vari-  
 ance  
 over  
 a  
 class  
 of  
 smooth  
 func-  
 tions.  
 Ob-

Let  $(\mathbb{B}, \mathbb{B}'', \Theta, \mathbf{y})$  denote the data given at the  $l$ -th iteration, as a generalized Rayleigh quotient:  $n^{-2}(\mathbb{B}\Theta^\top \mathbf{y})(\mathbb{B}\Theta^\top \mathbf{y})^\top$ .

**Proposition 2** (Regularity). Recall from (16) that  $V = n^{-2}(\mathbb{B}\Theta^\top \mathbf{y})(\mathbb{B}\Theta^\top \mathbf{y})^\top$ . Let  $\Lambda \in \mathbb{R}^{(MK+p) \times (MK+p)}$  be defined as the PLS component direction).  $\Lambda$  can be defined as the PLS component direction).

as

as:

$$\left(\hat{\gamma}_{11}, \dots, \hat{\gamma}_{1M}, \dots, \Lambda, \hat{\gamma}_{K+1}, \dots, \hat{\gamma}_{KM}, \hat{\mathbf{g}}_{KM}, \hat{\mathbf{g}}_{KM}^{\top}\right)$$

9)

The proof of Proposition 3.5 is provided in Appendix ??.

The constraint  $\boldsymbol{\xi}^{\top}(\mathbb{B} + \Lambda \mathbb{B}'')\boldsymbol{\xi} = 1$  enforces the orthonormality of the estimated

The

Here,

$$0_{p \times p}$$

denotes

the

$$p \times$$

$$p$$

zero

matrix.

The

co-

ef-

fi-

cients

of

the

squared

co-

vari-

ance

max-

i-

mizer

de-

fin-

in

(18),

are

ob-

tained

## Hybrid Partial Least Squares

ances PLS  
good- com-  
ness po-  
of nent  
fit di-  
and rec-  
smooth- tions  
ness with  
in re-  
 $\hat{\xi}_j$ . spect  
Smaller to  
 $\lambda_k$  a  
yields mod-  
com- i-  
po- fied  
nents in-  
that ner  
bet- prod-  
ter uct  
fit (see  
the Sec-  
data tion  
but 4.2  
risks for  
over- de-  
fit- tails).  
ting; The  
set- smooth-  
ting ing  
 $\lambda_k =$  pa-  
0 ram-  
re- e-  
cov- ter  
ers  $\lambda_k$   
the bal-

practice, both  $\{\lambda_k\}$  and the number of components  $L$  can be selected via cross-validation using a predictive criterion such as mean squared error.

unregularized solution in Proposition 3.4. Larger  $\lambda_k$  enforces greater smoothness, and in the limit  $\lambda_k \rightarrow \infty$ ,  $\hat{\xi}_j(t)$  approaches a linear form  $a + bt$ . In

one  
 struc-  
 ture  
 of  
 the  
 ma-  
 trix  
 $V$   
 Propo-  
 si-  
 tion  
 3.6  
 de-  
 rives  
 a  
 closed-  
 form  
 so-  
 lu-  
 tion  
 that  
 re-  
 quires  
 only  
 the  
 so-  
 lu-  
 tion  
 of  
 lin-  
 ear  
 sys-  
 tems.

**Proposition**  
**3**

*Computation.* The  
 gen-  
 er-  
 al-  
 ized  
 eigen-  
 prob-  
 lem  
 pre-  
 sented  
 in  
 Propo-  
 si-  
 tion 3.5  
 may  
 be  
 com-  
 pu-  
 ta-  
 tion-  
 ally  
 un-  
 sta-  
 ble  
 in  
 prac-  
 tice.  
 How-  
 ever,  
 by  
 lever-  
 ag-  
 ing  
 the  
 rank-

served (Closed–  
 data form  
 but so–  
 are lu–  
 not tion).  
 de- Con-  
 ci- sider  
 sion the  
 vari- op-  
 ables: ti-

$\mathbf{u}_j := B\Theta_j^\top \mathbf{y} \in \mathbb{R}^M$  for  $j = 1, \dots, K$ ,  
 miza-  
 tion

Let prob-

$$q := \sum_{j=1}^K \mathbf{u}_j^\top (B + \lambda_j B'')^{-1} \mathbf{u}_j + \mathbf{v}^\top \mathbf{v}.$$

lem  
 de-

Then scribed

the in  
 unique Propo-  
 (up si-  
 to tion **3.5**.  
 sign) De-  
 so- fine  
 lu- the  
 tion fol-  
 to low-  
 the ing  
 reg- quan-  
 u- ti-  
 lar- ties,  
 ized which  
 max- de-  
 i- pend  
 miza- on  
 tion the  
 prob- ob-



the  
func-  
tional  
and  
scalar  
com-  
po-  
nents  
sep-  
a-  
rately,  
fol-  
lowed  
by  
nor-  
mal-  
iza-  
tion  
by  
a  
com-  
mon  
fac-  
tor.  
Al-  
though  
the  
un-  
nor-  
mal-  
ized  
co-  
ef-  
fi-  
cients

*lem*  
*is*  
*given*  
*in*  
*closed*  
*form*  
*by*

$$\hat{\gamma}_j = \frac{1}{\sqrt{q}}(B + \lambda_j B'')^{-1}$$

The  
proof  
of  
Propo-  
si-  
tion 3.6  
is  
pro-  
vided  
in  
Ap-  
pendix ??.  
The  
ex-  
pres-  
sions  
above  
in-  
volve  
solv-  
ing  
lin-  
ear  
sys-  
tems  
for

en- are  
ables ob-  
the tained  
pro- in-  
ce- de-  
dure pen-  
to dently,  
cap- the  
ture nor-  
the mal-  
cor- iza-  
re- tion  
la- step  
tion cou-  
be- ples  
tween the  
the func-  
func- tional  
tional and  
and scalar  
scalar parts,  
com- al-  
po- low-  
nents ing  
of them  
the to  
PLS in-  
di- flu-  
rec- ence  
tion. one

### 3.3.2

*It-* other.  
*er-* This  
*a-* cou-  
*tive* pling

dic-  
tors  
and  
re-  
sponses.  
We  
first  
com-  
pute  
the  
in-  
di-  
vid-  
ual  
PLS  
score:

$step$   
 $2:$   
 $resid-$   
 $u-$   
 $al-$   
 $iza-$   
 $tion$   
 $via$   
 $hybrid-$   
 $on-$   
 $scalar$   
 $re-$   
 $gres-$   
 $sion.$

The

$l-$   
th  
it-  
er-  
a-  
tion's  
sec-  
ond  
step  
in-  
volves  
resid-  
u-  
al-  
iza-  
tion  
of  
both  
pre-

$\hat{\rho}_i^{[l]} := \langle \widetilde{W}_i^{[l]}, \hat{\xi}^{[l]} \rangle_{\mathbb{H}},$

Propo-

1)- si-  
th tions  
it- 3.5  
er- and  
a- 3.6.  
tion's Since  
re-  $\widetilde{W}_i^{[l]}$   
sponses are  
and as-  
hy- sumed  
brid to  
pre- have  
dic- a  
tors, sam-  
we ple  
regress mean  
the of  
( $l$ )- zero,  
th these  
it- PLS  
er- scores  
a- will  
tion's also  
re- have  
sponses a  
and sam-  
hy- ple  
brid mean  
pre- of  
dic- zero.  
tors To  
on ob-  
these tain  
PLS the  
scores ( $l+$

model: by least squares

$$\widetilde{W}_i^{[l]} = \widehat{\rho}_i^{[l]} \widehat{\delta}^{[l]} + \epsilon_i,$$

where and

$$\widehat{\delta}^{[l]} \in \widetilde{\mathbb{H}}$$

is the residual u-  
the al-  
re-ize.  
gres-Specif-  
sion i-  
co-cally,  
ef-the  
fi- $(l+$   
cient.1)-  
In th  
the pre-  
same dic-  
spirit tor  
as is  
the com-  
PLS puted  
com-as  
po-a  
nent resid-  
di-ual  
rec-of  
tion hybrid-  
es-on-  
ti-scalar  
ma-lin-  
tion ear  
step, re-  
rather gres-  
than sion

ex-  
pan-  
sion  
ap-  
proach  
to  
fit  
the  
en-  
tire  
hy-  
brid  
ob-  
ject  
in  
one  
step.  
There-  
fore,  
our  
method  
is  
com-  
pu-  
ta-  
tion-  
ally  
ef-  
fi-  
cient,  
and  
ap-  
pli-  
ca-  
ble  
treat-  
ing  
the  
hy-  
brid  
ob-  
ject  
as  
a  
long  
vec-  
tor  
of  
con-  
cate-  
nated  
func-  
tion  
eval-  
u-  
a-  
tions  
at  
time  
points  
and  
scalar  
vec-  
tors,  
we  
em-  
ploy  
a  
ba-  
sis

$\beta$  for  
seems dense  
to or  
so-ir-  
ley reg-  
rely u-  
on lar  
the func-  
smooth- tional  
ness data.  
of Con-  
 $\xi$ . se-

See quently,  
Sec-  $\delta^{[l]}$   
tion is  
**3.4.** ob-

$$\hat{\delta}^{[l]} := \arg \min_{\delta \in \tilde{\mathbb{H}}} \sum_{i=1}^n \|\widetilde{W}_i^{[l]} - \hat{\rho}_i^{[l]} \delta\|_{\mathbb{H}}^2.$$

On min-  
the i-  
other miz-  
hand, ing  
the a  
( $l+$  least  
1)-squares  
th cri-  
re- te-  
sponse rion:  
is im-  
com- pos-  
puted si-  
as ble.  
a Smooth-  
resid- ness  
ual of

ply, of  
 anal- a  
 o- scalar-  
 gous on-  
 to scalar  
 scalar lin-  
 PLS. ear

lemma re-  
 gres-  
 sion

[ model:

•  $Y_i^{[l]} = \hat{\nu}^{[l]} \hat{\rho}_i^{[l]} + \epsilon_i.$

Closed-

form The

so- fol-

lu- low-

tion] ing

Let propo-

us si-

de- tion

note demon-

$\hat{\boldsymbol{\rho}}^{[l]} :=$  strates

$(\hat{\rho}_1^{[l]}, \dots, \hat{\rho}_n^{[l]})^\top.$  that

The this

$(l+$  resid-

1)- u-

th al-

it- iza-

er- tion

a- step

tion's can

pre- be

dic- per-

tors formed

and sim-

re-



zero, their re-spec-tive resid-u-als,  $\widetilde{W}_i^{[l+1]}$  and  $Y_i^{[l+1]}$ , also main-tain a zero sam-ple mean.

sponses are com-puted as

$$\widetilde{W}_i^{[l+1]} := \widetilde{W}_i^{[l]} - \widehat{\rho}_i^{[l]} \widehat{\delta}_i^{[l]}$$

and

$$Y_i^{[l+1]} = Y_i^{[l]} - \widehat{\nu}^{[l]} \widehat{\rho}_i^{[l]},$$

Proof of Lemma 3.7 is provided in Appendix ??.

Since  $\widetilde{W}_i^{[l]}$  and  $Y_i^{[l]}$  are as-sumed to have a sam-ple mean of

a 3.4 *Final*  
 lin- *step:*  
 ear *es-*  
 com- *ti-*  
 bi- *mat-*  
 na- *ing*  
 tion *the*  
 of *hy-*  
 PLS *brid*  
 di- *re-*  
 rec- *gres-*  
 tions: *sion*

lemma *co-*  
*ef-*  
*fi-*  
*cient*

• The  
 lemma:recursive hy-  
 Let brid  
 use re-  
 de- gres-  
 fine sion  
 $\hat{\iota}^{[1]} :=$  co-  
 $\hat{\xi}^{[1]}$ . ef-  
 For fi-  
 $l \geq$  cient  
 2,  $\beta$   
 we in  
 re- model  
 cur- (4)  
 sively can  
 de- be  
 fine: writ-

$$\hat{\iota}^{[l]} = \hat{\xi}^{[l]} - \sum_{u=1}^{l-1} \langle \hat{\delta}^{[u]}, \hat{\xi}^{[l]} \rangle_{\mathbb{H}} \hat{\iota}^{[u]}.$$

Then

as:

we

$$Y_i = \sum_{l=1}^L \hat{\nu}^{[l]} \langle W_i^{[l]}, \hat{\xi}^{[l]} \rangle_{\mathbb{H}} + \epsilon_i = \langle W_i, \sum_{l=1}^L \hat{\rho}_i^{[l]} \rangle_{\mathbb{H}} + \epsilon_i$$

which,

have:

$$\hat{\rho}_i^{[l]} = \langle W_i^{[l]}, \hat{\xi}^{[l]} \rangle_{\mathbb{H}} = \langle W_i, \hat{\xi}^{[l]} \rangle_{\mathbb{H}}$$

given

Proof

the

of

unique-

Lemma

ness

3.8

of

is

$\beta$ ,

pro-

leads

vided

to

in

$$\hat{\beta} = \sum_{l=1}^L \hat{\nu}^{[l]} \hat{\iota}^{[l]} \sum_{l=1}^L \left( \hat{\nu}^{[l]} - \sum_{k=l+1}^L \hat{\nu}^{[k]} \langle \hat{\delta}^{[k]}, \hat{\delta}^{[l]} \rangle \right)$$

Thus

??.

the

Next,

num-

(23)

ber

leads

of

to

PLS

the

com-

fol-

po-

low-

nents

ing

$L$

model:

to

$$Y_i = \sum_{l=1}^L \hat{\nu}^{[l]} \hat{\rho}_i^{[l]} + \epsilon_i.$$

be

es-

This

ti-

model

mated

lets

can

us

be

to

cho-

ex-

sen

press

by

$Y_i$

**Algorithm****2**

Hy-brid

par-tial

least

squares

re-

gres-

sion

1:

**Ini-****tial-****ize:** $(\mathbb{B}, \mathbb{B}'', \Theta^{[1]}, \mathbf{y}^{[1]})$ 

as

the

data

ob-

jects

af-

ter

ba-

sis

ex-

pan-

sion,

fol-

low-

ing

Sec-

tion

**3.1.**

2:

 $\widetilde{W}_1^{[1]}, \dots, \widetilde{W}_n^{[1]}, \widetilde{Y}_1^{[1]}, \dots, Y_n^{[1]} \leftarrow$ 

stan-

dard-

ized

ver-

sions

of

 $W_1, \dots, W_n, Y_1, \dots, Y_n,$ 

fol-

low-

ing

Sec-

tion

**3.2**

3:

**for** $l =$  $1, 2, \dots, L$ **do**

4:

**PLS****di-****rec-****tion****and****score****es-****ti-****ma-****tion****(Propo-****si-****tion****3.6):**

5:

 $\mathbf{u}^{[l]} \leftarrow$ 

cross-

validation.

**4. Properties****of****the****hy-****brid****PLS**

This

sec-

tion

pro-

vides

the

math-

e-

mat-

i-

cal

er-

ties

that

sup-

port

the

al-

go-

rithm

sug-

gested

in

Sec-

tion

3.4, is well-defined under mild conditions. Section 4.2 demonstrates that our algorithm preserves the core properties of PLS, namely the orthogonality

3. Section 4.1 shows that the core optimization problem for the Partial Least Squares (PLS) discretization estimates the information step, presented in Proposition

go- of  
 rithm **2** the  
 and de-  
 es- rived  
 tab- di-  
 lish rec-  
 its tions  
 key and  
 math- the  
 e- or-  
 mat- thog-  
 i- o-  
 cal nal-  
 prop- ity  
 er- of  
 ties, the  
 namely scores.

the

or- 4.1

thog- *Tucker's*

o- *Cri-*

nal- *te-*

ity *tion*

of We

the now

PLS de-

di- rive

rec- the

tions pop-

and u-

scores, la-

as tion

well ana-

as logue

con- of

Al-

simply work with  $Y$  and  $W = (X_1, \dots, X_K, Z_1, \dots, Z_p)$ .

vergence to the true response.

Throughout this section we assume  $\mathbb{E}[Y] = 0 \in \mathbb{R}$  and  $\mathbb{E}[W] = 0 \in \mathbb{H}$ .

the population covariance level, where the random domain objects are fully observable.

The servable, we may drop the sample in-  
 dex  $i$  and

core optimization problem in PLS is for-

er-  
a-  
tors,  
which  
we  
de-  
fine  
next.  
For  
no-  
ta-  
tional  
sim-  
plic-  
ity,  
we  
sup-  
press  
the  
it-  
er-  
a-  
tion  
in-  
dex  
 $l$ ,  
with  
the  
un-  
der-  
stand-  
ing  
that  
for  
 $l =$

mu-  
lated  
in  
terms  
of  
max-  
i-  
miz-  
ing  
the  
squared  
cross-  
covariance.  
At  
the  
pop-  
u-  
la-  
tion  
level,  
this  
quan-  
tity  
is  
nat-  
u-  
rally  
char-  
ac-  
ter-  
ized  
by  
cross-  
covariance  
op-



$\mathbb{E}[W \otimes W]$ , so that for  $u, v \in \mathbb{H}$ , we have
 
$$\Sigma_W u = \mathbb{E}[\langle W, u \rangle_{\mathbb{H}} W], \quad \langle \Sigma_W u, v \rangle_{\mathbb{H}} = \mathbb{E}[\langle W, u \rangle_{\mathbb{H}} \langle W, v \rangle_{\mathbb{H}}].$$
 The original cross-covariances between the  $l \geq 2$  response  $Y$  and the predictors are
 
$$\sigma_{YX} := (\mathbb{E}[Y X_1], \dots, \mathbb{E}[Y X_K]) \in \mathcal{F}^K$$
 The versions of the cross-covariance between  $Y$  and  $X$  are defined as follows.

def:cross\_cov\_terms The

Let the  
 $W^{[1]} :=$  hy-  
 $W$  brid  
and pre-  
 $Y^{[1]} :=$  dic-  
 $Y$ . tor  
For  $W$   
any is  
in- then  
te-  
ger  $\Sigma_{YW} := \mathbb{E}[YW] = (\sigma$   
 $l \geq$  At  
2, the  
to pop-  
be u-  
the la-  
resid- tion  
u- level,  
als Al-  
of go-  
the rithm  
where 2  
and with-  
which out  
is reg-  
at- u-  
tained lar-  
by izaiton  
the can  
lead- be  
ing ex-  
eigen- pressed  
func- as  
tion fol-  
of lows.

the  
 operation  
 a-  
 tor  
 $\mathcal{U}^{[l-1]}$ ,  
 de-  
 fined  
 anal-  
 o-  
 gously  
 to  
 Lemma 4.6  
 us-  
 ing  
 $W^{[l-1]}$   
 and  
 $Y^{[l-1]}$ ,  
 pro-  
 vided  
 the  
 con-  
 di-  
 tions  
 in  
 Lemma 4.4  
 are  
 sat-  
 is-  
 fied.  
 definition  
 the  
 con-  
 strained  
 gen-  
 Pop-

**Proposition****4**

(Prop-

er-

ties

of

pop-

u-

la-

tion

PLS).

*The**so-**lu-**tion**of**op-**ti-**miza-**tion**prob-**lem*

(24)

*of**Def-**i-**ni-**tion*

4.2

*is**equiv-**a-**lent**to**that*

er-

al-

ized

eigen-

prob-

lem:

$$\xi^{[l]} = \arg \max_{h \in \mathbb{H}} \text{Cov}^2(\langle$$

As

in

scalar

case,

this

prob-

lem

can

be

solved

by

an

it-

er-

a-

tive

al-

go-

rithm,

and

yields

or-

thog-

o-

nal

scores:

is of  
 provided population  
 in the  
 Appendix A  
 ??.

We go-  
 show rithm  
 that 3).  
 problem For  
 lem every  
 (24)  $s \geq$   
 and 1  
 Algorithm and  
 rithm every  
 3 are  $j \leq$   
 well-  $s -$   
 defined 1,  
 under we  
 der have  
 mild  $\mathbb{E}[\rho^{[j]} \rho^{[s]}] =$   
 con- 0  
 di- and  
 tions  $\mathbb{E}[Y^{[s]} \rho^{[j]}] =$   
 by 0.  
 es-  
 tab- Proof  
 lish- of  
 ing Propo-  
 that si-  
 line tion

3  
of  
Al-  
go-  
rithm

3  
is  
well-  
defined  
un-  
der  
mild  
con-  
di-  
tions.

For  
clar-  
ity,  
we  
de-  
fine  
the  
rel-  
e-  
vant  
op-  
er-  
a-  
tors  
while  
omit-  
ting  
the  
it-  
er-

### Algorithm

3

Pop-  
u-  
la-  
tion  
NI-  
PALS

```

1:  $(W^{[1]}, Y^{[1]}) \leftarrow (W, Y)$ 
2: for  $l = 1, 2, \dots, L$  do
3:  $\xi^{[l]} \leftarrow \arg \max_{h \in \mathbb{H}} \text{Cov}^2$ 
   1,
    $\triangleright$  PLS
   di-
   rec-
   tion
4:  $\rho^{[l]} \leftarrow \langle \xi^{[l]}, W^{[l]} \rangle$ 
    $\triangleright$  PLS
   score
5:  $\delta^{[l]} \leftarrow \frac{1}{\mathbb{E}[(\rho^{[l]})^2]} \mathbb{E}[W^{[l]} \rho^{[l]}]$ 
    $\triangleright$  Lin-
   ear
   re-
   gres-
   sion
   of
    $W^{[l]}$ 
   on
    $\rho^{[l]}$ 
6:  $\nu^{[l]} \leftarrow \frac{1}{\mathbb{E}[(\rho^{[l]})^2]} \mathbb{E}[Y^{[l]} \rho^{[l]}]$ 
    $\triangleright$  Lin-
   ear
   re-
   gres-
   sion
   of
    $Y^{[l]}$ 
   on
    $\rho^{[l]}$ 
7:  $W^{[l+1]} \leftarrow W^{[l]} - \rho^{[l]} \delta^{[l]}$ 
    $\triangleright$  Resid-
   u-
   al-
   ized
   pre-
   dic-
   tor
8:  $Y^{[l]} \leftarrow$ 
```

are a-  
 bounded tion  
 in in-  
 the dex  
 sense for  
 that the  
 there resid-  
 ex- u-  
 ist al-  
 fi- ized  
 nite pre-  
 con- dic-  
 stants tors  
 $Q_1$  and  
 and re-  
 $Q_2$  sponses.  
 such lemma  
 that

$$\max_{k=1,\dots,K} \sup_{t \in [0,1]} \mathbb{E}[Y X_k](t)^2 < Q_1 \quad \text{and} \\ \bullet$$

the lemma:cross<sub>cov</sub>function  
 op-  $C_{YW}$   
 er- such  
 a- that  
 tor for  
 $C_{YW}$  any  
 is  $h =$   
 a  $(f_1, \dots, f_K, \mathbf{v}) \in$   
 com-  $\mathbb{H}$ ,  
 pact  $C_{YW}h := \mathbb{E} [\langle W, h \rangle_{\mathbb{H}} Y]$   
 op-  
 er- If  
 a-  $\sigma_{YX}$   
 tor. and  
 $\sigma_{YZ}$

tor  
of  
 $\mathcal{C}_{YW}$ .  
That  
is,  
 $\mathcal{C}_{WY} =$   
 $\mathcal{C}_{YW}^*$ .

The  
proof  
of  
Lemma  
[4.5](#)

is  
pro-  
vided  
in  
Ap-  
pendix  
[??](#).

[lemma](#)  
[  
.

Com-  
pos-  
ite  
cross-  
covariance  
op-  
er-  
a-  
tor]  
De-  
fine  
 $\mathcal{U} :=$

Proof  
of  
Lemma

[4.4](#)  
is  
pro-  
vided

in  
Ap-  
pendix  
[??](#).

[lemma](#)

•  
[lemma:cross\\_cov\\_operat](#)

$\mathbb{E}[Y \otimes$   
 $W],$

such  
that  
for

any  
 $d \in$

$\mathbb{R},$

$\mathcal{C}_{WY}d := \mathbb{E}[\langle Y, d \rangle W]$

Then

$\mathcal{C}_{WY}$

is

an

ad-

joint

op-

er-

a-



semidefinite	$\mathcal{C}_{WY} \circ$
op-	$\mathcal{C}_{YW} :$
er-	$\mathbb{H} \rightarrow$
a-	$\mathbb{H}$
tor.	as
Un-	an
der	op-
the	er-
con-	a-
di-	tor
tions	which
of	per-
Lemma	forms
4.4,	the
$\mathcal{U}$	fol-
is	low-
a	ing
com-	map-
pact	ping:
op-	$\mathcal{U}h = \mathcal{C}_{WY}(\mathcal{C}_{YW}h) =$
er-	
a-	In
tor.	other
	words,
The	$\mathcal{U} =$
proof	$\Sigma_{YW} \otimes$
of	$\Sigma_{YW}.$
Lemma	Then
4.6	$\mathcal{U}$
is	is
pro-	a
vided	self-
in	adjoint
Ap-	and
pendix	positive-
??.	

sys- By  
 tem the  
 of Hilbert-  
 eigen- Schmidt  
 func- the-  
 tions o-  
 $\{\xi_{(u)}\}_{u \in \mathbb{N}}$  rem  
 of (e.g.,  
 $\mathcal{U}$  The-  
 in o-  
 $\mathbb{H}$  rem  
 such 4.2.4  
 that in  
 $\mathcal{U}\xi_{(u)} =$  [Hs-](#)  
 $\kappa_{(u)}\xi_{(u)},$  [ing](#)  
 where [and](#)  
 $\{\kappa_{(u)}\}_{u \in \mathbb{N}}$  [Eu-](#)  
 are [bank,](#)  
 the [2015\),](#)  
 cor- Lemma  
 re- [4.6](#)  
 spond- guar-  
 ing an-  
 se- tees  
 quence the  
 of ex-  
 eigen- is-  
 val- tence  
 ues of  
 that a  
 goes com-  
 to plete  
 zero or-  
 as thonor-  
 $u \rightarrow$  mal

that  $\infty$ ,  
 line that  
 3 is,  
 of  $\kappa_{(1)} \geq$   
 Al-  $\kappa_{(2)} \geq$   
 go-  $\dots \geq$   
 rithm 0.

3 The  
 is fol-  
 well- low-  
 defined ing  
 un- the-  
 der o-  
 mild rem  
 con- presents  
 di- population-  
 tions. level

# theoremPLS

and  
 Tucker's  
 [ Cri-  
 • te-  
 Tucker's rion  
 cri- for  
 te- our  
 rion] scalar-  
 Un- on-  
 der hybrid  
 the re-  
 con- gres-  
 di- sion  
 tions model  
 of and  
 Lemma en-  
 4.4, sures  
 the

o- con-  
rem strained

4.7 max-

is i-

pro- mum

vided

in  $\max_{\|\xi\|_{\mathbb{H}}=1} \text{Cov}^2(\langle W, \xi \rangle_{\mathbb{H}},$

Ap- is

pendix at-

?? tained

Finally, by

the the

fol- eigen-

low- func-

ing tion

the- as-

o- so-

rem ci-

states ated

that with

the the

fit- largest

ted eigen-

val- value

ues of

from the

hy- op-

brid er-

PLS a-

con- tor

verge  $\mathcal{U}$ .

to

those The

of proof

or- of

The-

of  
a  
full  
lin-  
ear  
ear  
model  
when  
a  
suf-  
fi-  
cient  
num-  
ber  
of  
com-  
po-  
nents  
is  
used.

di-  
nary  
lin-  
ear  
re-  
gres-  
sion  
in  
the  
mean-  
squared  
sense.  
This  
shows  
that  
hy-  
brid  
PLS,  
as

theorema

dimension-  
reduction  
[  
•  
 $L^2$   
Con-  
ver-  
gence  
of  
Hy-  
brid  
PLS  
at  
Pop-  
u-  
la-

dimension-  
reduction  
tech-  
nique,  
can  
ef-  
fec-  
tively  
cap-  
ture  
the  
pre-  
dic-  
tive  
power

sense:

tion

$$\lim_{L \rightarrow \infty} \mathbb{E} \left[ \|Y_{\text{PLS},L} - Y\|_2^2 \right] = 0$$

Level]  
Let

The

$$Y_{\text{PLS},L} =$$

proof

$$\sum_{l=1}^L \nu^{[l]} \rho^{[l]}$$

of

be

The-

the

o-

Hy-

rem

brid

**4.8**

PLS

is

fit-

pro-

ted

vided

value

in

with

Ap-

$L$

pendix

com-

??.

po-

nents.

4.2 *Geometric*

Then,

*prop-*

the

*er-*

Hy-

*ties*

brid

A

PLS

fun-

fit-

da-

ted

men-

value

tal

con-

prop-

verges

erty

to

of

$Y$

par-

in

tial

the

least

mean-

squares,

squared

as

seem

onal. Our regularization estimates preserve this property, with respect to a modified inner product that incorporates the roughness

in Definition 4.2 and Proposition 4.3, is that between iterations, its derived discrepancies are orthonormal and PLS scores are orthogonal

defined in Definition 3.3, and a roughness penalty matrix  $\Lambda = \text{blkdiag}(\lambda_1 I_M, \dots, \lambda_K I_M, 0_{p \times p})$ , the roughness-sensitive inner product between  $W_1$  and  $W_2$  is defined as:

penalty, defined as follows: definition 3.3, and a [ . Roughness-sensitive inner product  $\Lambda = \text{blkdiag}(\lambda_1 I_M, \dots, \lambda_K I_M, 0_{p \times p})$ , Given two hybrid predictors  $W_1 = (X_{11}, \dots, X_{1K}, \mathbf{Z}_1)$  and  $W_2 = (X_{21}, \dots, X_{2K}, \mathbf{Z}_2)$ , both elements

$$\langle W_1, W_2 \rangle_{\mathbb{H}, \Lambda} := \sum_{k=1}^K \int_0^1 X_{1k}(t) X_{2k}(t) dt$$

as



thonormal. Based on this **Proposition 5** (Orthogonality of estimated PLS components  $\hat{\xi}^{[1]}, \hat{\xi}^{[2]}, \dots, \hat{\xi}^{[L]}$ , Proposition 3.5 via Proposition

si-

tion

4.10

is

pro-

vided

in

Ap-

pendix

??.

The

next

propo-

si-

tion

states

that

the

vec-

tors

of

es-

ti-

mated

PLS

scores

for

dif-

fer-

ent

it-

er-

a-

tion

num-

3.5

with

a

rough-

ness

penalty

ma-

trix

$\Lambda =$

$\text{blkdiag}(\lambda_1 I_M, \dots, \lambda_P$

are

mu-

tu-

ally

or-

thonor-

mal

with

re-

spect

to

the

in-

ner

prod-

uct

$\langle \cdot, \cdot \rangle_{\mathbb{H}, \Lambda}$ .

That

is,

$\langle \widehat{\xi}^{[l_1]}, \widehat{\xi}^{[l_2]} \rangle_{\mathbb{H}, \Lambda} = \mathbb{1}(l_1$

The

proof

of

Propo-

es- bers  
 ti- are  
 mated mu-  
 PLS tu-  
 scores ally  
 ( $l =$  or-  
 $1, \dots, L$ ) thog-  
 of o-  
 $n$  nal.  
 ob-

ser- **Proposition**

va- **6.**

tions. Re-

The call

vec- from

tors Lemma

$\hat{\boldsymbol{\rho}}^{[1]}, \hat{\boldsymbol{\rho}}^{[2]}, \dots, \hat{\boldsymbol{\rho}}^{[L]}$  3.7

are that

mu-  $\hat{\boldsymbol{\rho}}^{[l]}$

tu- de-

ally note

or- the

thog-  $n$ -

o- dimensional

nal vec-

in tor

the whose

sense el-

that e-

$\hat{\boldsymbol{\rho}}^{[l_1]\top} \hat{\boldsymbol{\rho}}^{[l_2]} = 0$  ments  
 for  $l_1, l_2 \in \{1, \dots, L\}$   
 con-

The sist

proof of

of the

Propo-  $l$ -

th

tures  
—  
specif-  
i-  
cally,  
de-  
pen-  
den-  
cies  
among  
func-  
tional

si-  
tion  
**4.11**  
is  
pro-  
vided  
in  
Ap-  
pendix  
**??**.

pre-  
dic-  
tors,  
among  
scalar  
pre-  
dic-  
tors,  
and  
be-  
tween  
scalar  
and  
func-  
tional  
pre-  
dic-  
tors  
—

## **5. Simulation studies**

To  
eval-  
u-  
ate  
the  
su-  
pe-  
ri-  
or-  
ity  
of  
our  
method  
un-  
der  
com-  
plex

### **5.1 *Synthetic data***

To  
gen-  
struc-

de-  
pen-  
dency  
struc-

*Hybrid Partial Least Squares*

$3 \times 100$  matrix, where each row represents the  $i$ th observation of a functional pre-dictor with 100 evaluations. This method gives us pre-er-ate functional pre-dictors, we use a matrix-normal distribution, also known as a Kronecker-separable covariance model. For each of  $i = 1, \dots, n$ , we create a

in- cise  
 sight con-  
 be- trol  
 hind over  
 Kro- cross-  
 necker modal  
 sep- (between-  
 a- row)  
 ra- de-  
 bil- pen-  
 ity dence  
 is and  
 that smooth-  
 if ness  
 $\mathbf{Y} \sim$  of  
 $\mathcal{MN}(\mathbf{M}, \mathbf{R}, \mathbf{C})$ , the  
 then curves  
 its (between-  
 vec- column  
 tor- de-  
 ized pen-  
 form dence).  
 fol-  
 lows  $\mathbf{X} \sim \mathcal{MN}_{m \times n}(\mathbf{M}; \mathbf{R}$   
 a and  
 mul- its  
 ti- log-  
 vari- density  
 ate is  
 nor- given  
 mal by  
 dis-  
 tri-  $\log p(\mathbf{X} \mid \mathbf{M}, \mathbf{R}, \mathbf{C}) =$   
 bu- The  
 tion: key

work  
 lever-  
 ag-  
 ing  
 the  
 matrix-  
 normal  
 dis-  
 tri-  
 bu-  
 tion,  
 an  
 ap-  
 proach  
 pre-  
 vi-  
 ously  
 sug-  
 gested  
 by  
 our  
 ad-  
 vi-  
 sor.  
 The  
 goal  
 was  
 to  
 cre-  
 ate  
 a  
 dataset  
 with  
 a  
 spe-

$\text{vec}(\mathbf{Y}) \sim$   
 $\mathcal{N}(\text{vec}(\mathbf{M}), \mathbf{C} \otimes$   
 $\mathbf{R}),$   
 where  
 $\otimes$   
 de-  
 notes  
 the  
 Kro-  
 necker  
 prod-  
 uct  
 and  
 $\text{vec}$   
 is  
 the  
 $\text{vec}$ -  
 tor-  
 iza-  
 tion  
 op-  
 er-  
 a-  
 tor.  
 We  
 de-  
 vel-  
 oped  
 a  
 data  
 gen-  
 er-  
 a-  
 tion  
 frame-

To  
in-  
tro-  
duce  
a  
struc-  
tured  
de-  
pen-  
dence  
within  
each  
row,  
the  
row  
co-  
vari-  
ance  
ma-  
trix  
was  
set  
to  
mimic  
an  
AR(1)  
pro-  
cess,  
which  
is  
a  
spe-  
cial  
case  
of  
cific,  
con-  
trol-  
lable  
cor-  
re-  
la-  
tion  
struc-  
ture  
be-  
tween  
func-  
tional  
and  
scalar  
pre-  
dic-  
tors.  
cor-  
re-  
sponds  
to  
the  
num-  
ber  
of  
eval-  
u-  
a-  
tion  
points  
for  
each  
curve.



### *Hybrid Partial Least Squares*

each a  
ma- Gaus-  
trix sian  
were pro-  
treated cess.  
as This  
func- en-  
tional sures  
data. that  
The each  
third row  
row ex-  
was hibits  
con- the  
verted smooth,  
into curve-  
a like  
set char-  
of ac-  
scalar ter-  
pre- is-  
dic- tics  
tors. of  
This a  
was func-  
ac- tional  
com- ob-  
plished ser-  
by va-  
par- tion.  
ti-  
tion-  
ing  
the  
third of

The

first  
two  
rows  
of

tors, row  
 we into  
 would blocks  
 par- and  
 ti- then  
 tion av-  
 the er-  
 row ag-  
 into ing  
 (1,3), the  
 (2,4), val-  
 and ues  
 (3,4), within  
 yield- each  
 ing block.  
 the For  
 scalar in-  
 val- stance,  
 ues if  
 2.0,3.0, the  
 and third  
 3.5. row

Building was

on (1,3,2,4,3,4)  
 the and  
 func- we  
 tional wanted  
 graph- to  
 i- gen-  
 cal er-  
 model ate  
 sim- three  
 u- scalar  
 la- pre-  
 tion dic-

as of  
 Gaus- [Zhu](#)  
 sian [et al.](#)  
 pro- (2016),  
 cesses we  
 us- gen-  
 ing er-  
 a ate  
 trun- a  
 cated mixed  
 Karhunen- graph-  
 Loève i-  
 ex- cal  
 pan- model  
 sion, with  
 where five  
 the nodes,  
 eigen- de-  
 ba- scribed  
 sis as  
 con- fol-  
 sists lows:  
 of  
 Fourier •  
 ba- Nodes  
 sis F1  
 func- and  
 tions F2:  
 with Two  
 a func-  
 fixed tional  
 num- pre-  
 ber dic-  
 of tors  
 ba- mod-  
 eled

the functional predictors, these scalar predictors are modeled directly without basis expansion.

sis functions,  $M = 9$ .

•

scalar predictors are modeled directly without basis expansion.

Nodes S1, S2, and S3: Three scalar predictors, each following an  $s$ -dimensional

To capture dependence among predictors, we

multivariate normal distribution. Unlike

con-  
nected  
graph.  
In  
the  
Gaus-  
sian  
pro-  
cess  
frame-  
work,  
the  
pre-  
ci-  
sion  
ma-  
trix  
 $\mathbf{R}_0^{-1}$   
en-  
codes  
con-  
di-  
tional  
in-  
de-  
pen-  
dence  
re-  
la-  
tion-  
ships,  
while  
its  
in-  
verse,

in-  
tro-  
duce  
a  
graph  
struc-  
ture  
that  
gov-  
erns  
their  
con-  
di-  
tional  
cor-  
re-  
la-  
tions.  
We  
con-  
sider  
two  
types  
of  
graph  
struc-  
tures:  
a  
weakly  
con-  
nected  
graph  
and  
a  
strongly

tions  $\mathbf{R}_0$ ,  
 be- rep-  
 tween re-  
 FPC sends  
 scores marginal  
 and co-  
 scalar vari-  
 pre- ances.  
 dic- This  
 tor struc-  
 val- ture  
 ues. ex-  
 Each tends  
 block to  
 $(i, j)$  a  
 of block-  
 $\mathbf{R}$  wise  
 is cor-  
 given re-  
 by la-  
 $(R_0)_{ij} \mathbf{I}_{M_i, M_j}$ , tion  
 where ma-  
 $\mathbf{I}_{M_i, M_j}$  trix  
 is  $\mathbf{R} \in$   
 a  $\mathbb{R}^{(2M+3s) \times (2M+3s)}$ ,  
 rect- where  
 an- off-  
 gu- diagonal  
 lar blocks  
 iden- rep-  
 tity re-  
 ma- sent  
 trix. cor-  
 Here, re-  
 $M_i =$  la-

tor, 9

we if

ass- node

gin  $i$

$M$  cor-

ref- re-

er- sponds

ence to

eigen- a

val- func-

ues tional

(or pre-

FPC dic-

score) tor

drawn and

in-  $M_j =$

de-  $s$

pen- if

dently node

from  $j$

gamma cor-

dis- re-

tri- sponds

bu- to

tions a

with scalar

de- pre-

creas- dic-

ing tor.

means. For

These each

ref- func-

er- tional

ence pre-

eigen- dic-

are  
fixed  
over  
all  
sam-  
ples  
and  
all  
in-  
de-  
pen-  
dent  
rep-  
e-  
ti-  
tion  
of  
the  
ex-  
per-  
i-  
ment.  
We  
draw  
it  
one  
ran-  
domly  
just  
to  
en-  
sure  
the  
dif-  
fer-

val-  
ues  
will  
later  
be  
mul-  
ti-  
plies  
by  
ran-  
domly  
mul-  
ti-  
plier  
drawn  
from  
cor-  
re-  
lated  
mul-  
ti-  
variate  
Nor-  
mal  
dis-  
tri-  
bu-  
tion.  
These  
ref-  
er-  
ence  
eigen-  
val-  
ues



ti-  
 vari-  
 ate  
 nor-  
 mal  
 data  
 from  
 the  
 co-  
 vari-  
 ance  
 ma-  
 trix  
**R.**  
 The  
 last  
 $3s$   
 com-  
 po-  
 nents  
 are  
 as-  
 signed  
 as  
 scalar  
 pre-  
 dic-  
 tors,  
 while  
 the  
 first  
 $2M$   
 com-  
 po-  
 nents,

en-  
 ti-  
 a-  
 tion  
 be-  
 tween  
 the  
 two  
 func-  
 tional  
 pre-  
 dic-  
 tors.  
 we  
 in-  
 de-  
 pen-  
 dently  
 draw  
 for  
 each  
 func-  
 tional  
 pre-  
 dic-  
 tor  
 from  
 .  
 We  
 then  
 sam-  
 ple  
 zero-  
 mean  
 mul-

ing scaled  
 a by  
 Fourier their  
 ba- cor-  
 sis. re-

Finally, spond-

we ing  
 in- eigen-  
 tro- val-  
 duce ues,  
 struc- serve  
 tured as  
 vari- FPC  
 abil- scores.  
 ity These  
 by scores  
 adding are  
 a then  
 com- ex-  
 mon panded  
 mean into  
 func- func-  
 tion, tional  
 de- data,  
 fined eval-  
 as u-  
 a ated  
 scaled over  
 and 100  
 shifted equally  
 sine spaced  
 func- points  
 tion. on  
 To  $[0, 1]$   
 vi- us-

graph	su-
struc-	al-
tures,	ize
il-	the
lus-	gen-
trat-	er-
ing	ated
the	data,
dis-	Fig-
tinct	ure
de-	??
pen-	plots
dency	func-
pat-	tional
terns	pre-
in-	dic-
duced	tor
by	re-
the	al-
pre-	iza-
ci-	tions
sion	and
ma-	heatmaps
tri-	of
ces.	sam-
	ple
5.2 Simulation	cor-
<i>shown</i>	re-
<i>in</i>	la-
<i>febru-</i>	tion
<i>ary</i>	ma-
<i>2025</i>	tri-
<i>meet-</i>	ces
<i>ing</i>	for
To	both
ef-	

Markov  
ran-  
dom  
field  
(GMRF)  
within  
the  
frame-  
work  
of  
Gaus-  
sian  
undi-  
rected  
graph-  
i-  
cal  
mod-  
els.  
In  
a  
GMRF,  
the  
off-  
diagonal  
el-  
e-  
ments  
of  
the  
pre-  
ci-  
sion  
ma-  
trix

fec-  
tively  
man-  
age  
the  
de-  
pen-  
den-  
cies  
among  
func-  
tional  
pre-  
dic-  
tors,  
scalar  
co-  
vari-  
ates,  
and  
be-  
tween  
func-  
tional  
and  
scalar  
pre-  
dic-  
tors,  
we  
uti-  
lize  
a  
Gaus-  
sian

we adopt the simulation setup proposed by [Kolar et al. \(2014\)](#), which focuses on mixed attribute Gaussian graphical models. We construct a graph con-  
capture the conditional correlations between the corresponding components. Given that our setting involves both functional and scalar co-variates,

tion  
pro-  
cess  
for  
each  
node.

sist-  
ing  
of  
two  
func-  
tional

pre-  
*Functional* dic-  
*Pre-* tor  
*dic-* nodes  
*tors.* and

We  
con-  
sider  
two  
func-  
tional  
pre-  
dic-  
tors  
that  
share  
the  
same  
pre-  
ci-  
sion  
ma-  
trix.

three  
vec-  
tor  
pre-  
dic-  
tor  
nodes.

Be-  
low,  
we  
se-  
quen-  
tially  
de-  
scribe  
the  
graph  
struc-

ture  
and  
the  
gen-  
er-  
a-

De-  
noted  
as  
 $\Theta :=$   
 $(\theta_{t\tilde{t}}) \in$   
 $\mathbb{R}^{p \times p},$

this

a	pre-
smooth	ci-
func-	sion
tional	ma-
tra-	trix
jec-	fol-
tory.	lows
More	an
for-	AR(1)
mally,	struc-
each	ture
func-	with
tional	a
pre-	white
dic-	noise
tor	vari-
is	ance
eval-	of
u-	$0.1^2$
ated	and
at	an
$p$	au-
equally	tore-
spaced	gres-
points	sive
on	co-
$[0, 1]$ ,	ef-
with	fi-
val-	cient
ues	of
de-	=
fin	0.95,
re-	en-
cur-	sur-
sively	ing

by: as:

$$\theta_{tt} = \begin{cases} \frac{1}{\sigma^2(1 - 0.95^2)} X_i^{(k)}(0) & t=0, 1, p, X_i^{(k)}(t) \\ \frac{1 + 0.95^2}{\sigma^2(1 - 0.95^2)}, & 2 \leq t \leq p - 1 \end{cases}$$

The this  
off- set-  
diagonal ting,  
en- the  
tries pre-  
are: ci-

$$\theta_{t,t+1} = \theta_{t+1,t} = -\frac{\rho}{\sigma^2(1 - \rho^2)}, \quad 1 \leq$$

Finally, trix  
the  $\Theta$   
func- ex-  
tional hibits  
pre- a  
dic- tridi-  
tors ag-  
are o-  
smoothed nal  
us- Toeplitz  
ing struc-  
a ture,  
B- where  
spline the  
ba- di-  
sis ag-  
with o-  
15 nal  
ba- en-  
sis tries  
func- are  
tions, given  
as



a shared pre-cision matrix. This pre-cision matrix, denoted as  $\Gamma := (\gamma_{ij}) \in \mathbb{R}^{d \times d}$ , follows a Toeplitz structure with exponentially decaying entries, described in Section 3.3.1. *Vector Pre-dictors.* We consider  $s$  vector predictors, each following a  $d$ -dimensional zero-mean Gaussian distribution with

to given  
the by:

first,  
as  $\gamma_{ij} := 0.5^{|i-j|}$ .

il-  
lus- *Graph*  
trated *Struc-*  
in *ture.*

Fig- We  
ure ??.

An range

edge five

in nodes

the in

graph a

in- chain

di- struc-

cates ture,

that where

the each

con- node

nected fol-

nodes lows

re- a

main se-

cor- quen-

re- tial

lated or-

when der,

the and

val- the

ues last

of node

all con-

other nects

back

$F_2$  nodes  
as are  
func- fixed.  
tional The  
pre- re-  
dic- sult-  
tors ing  
and marginal  
nodes de-  
 $V_1$ , pen-  
 $V_2$ , dence  
and struc-  
 $V_3$  ture  
as is  
vec- sig-  
tor nif-  
pre- i-  
dic- cantly  
tors. more  
To com-  
in- plex  
tro- than  
duce the  
con- chain  
di- struc-  
tional ture  
de- it-  
pen- self.  
den- We  
cies des-  
be- ig-  
tween nate  
the nodes  
com-  $F_1$   
po- and

er- nents,  
wise. we  
Here, set  
**1** the  
de- off-  
notes diagonal  
a blocks  
ma- of  
trix the  
of pre-  
ap- ci-  
pro- sion  
pri- ma-  
ate trix  
di- to  
men- be  
sions 0.51  
where if  
all the  
el- cor-  
e- re-  
ments spond-  
are ing  
equal com-  
to po-  
1. nents  
Over- are  
all, con-  
we nected  
gen- in  
er- the  
ate graph,  
a and  
 $2p+$  zero  
 $3d-$  oth-

observed from this multivariate Gaussian distribution, we process the regression coefficients for the first functional predictive

dimensional multivariate Gaussian distribution, with mean zero and a precision matrix  $\Omega$ , structured as follows:

$$\Omega = \begin{pmatrix} \Omega_F & 0.5\mathbf{1} & 0 \\ 0.5\mathbf{1} & \Omega_F & 0 \\ 0 & 0.5\mathbf{1} & 0 \\ 0 & 0 & 0 \\ 0.5\mathbf{1} & 0 & 0 \end{pmatrix}$$

For each

dic- tor,  
 tor,  $\beta_{F_1}$ ,  
 $\beta_{F_2}$ , are  
 are drawn  
 drawn from  
 in- a  
 de- mul-  
 pen- ti-  
 dently vari-  
 from ate  
 the nor-  
 same mal  
 dis- dis-  
 tri- tri-  
 bu- bu-  
 tion tion  
 and  $N(0, 5\mathbf{I}_p)$ ,  
 are with  
 also a  
 smoothed fixed  
 us- ran-  
 ing dom  
 a seed.  
 B- The  
 spline co-  
 ba- ef-  
 sis fi-  
 with cients  
 10 for  
 ba- the  
 sis sec-  
 func- ond  
 tions. func-  
 For tional  
 the pre-

*Hybrid Partial Least Sq*

their	vec-
cor-	tor
re-	co-
spond-	vari-
ing	ates,
co-	the
ef-	re-
fi-	gres-
cients,	sion
in-	co-
de-	ef-
pen-	fi-
dent	cients
Gaus-	are
sian	sam-
noise	pled
from	from
$N(0, 0.1^2)$	$N(0, I_d)$ .
is	Af-
added	ter
to	cal-
the	cu-
gen-	lat-
er-	ing
ated	the
re-	in-
sponses	ner
to	prod-
sim-	uct
u-	of
late	the
mea-	co-
sure-	vari-
ment	ates
noise.	and

pre-  
dic-  
tors [Happ  
and  
Greven  
\(2018\)](#)  
and  
scalar  
PCA  
and  
run  
OLS  
on  
the  
PC  
scores.

The  
base-  
line  
meth-  
ods  
are:

- Penalized  
func-  
tional  
re-  
gres-  
sion  
(pfr) [Gold-  
smith  
et al.  
\(2011\)](#)

We

com-  
pare  
our  
method  
with  
these  
base-  
line  
meth-  
ods  
us-  
ing  
 $p =$   
100.

- Principal  
com-  
po-  
nent  
re-  
gres-  
sion  
(fpcr):  
run  
both  
of  
PCA  
for  
mult-  
ple  
func-  
tional

We  
con-  
sider  
sce-



*Hybrid Partial Least Sq*

ing nar-  
30% ios  
test with  
set,  $d =$   
us- 1, 2, 3, 4, 5  
ing and  
the  $n =$   
root 100, 200, 300, 400.  
pre- For  
dic- each  
tion sce-  
mean nario,  
squared we  
er- use  
ror 70%  
as of  
the the  
eval- data  
u- for  
a- train-  
tion ing  
met- and  
ric. eval-  
For u-  
our ate  
method pre-  
and dic-  
PCR, tion  
the per-  
max- for-  
i- mance  
mum on  
num- the  
ber re-  
of main-

strate that our method consistently outperforms the baseline methods across all scenarios.

com-  
po-  
nents  
are  
set  
as  
20.  
The  
num-  
ber  
of  
com-  
po-  
nents  
is  
cho-  
sen  
by  
5-  
fold

## 6. Data

### Ap- pli- ca- tion

*Renal study data.* We applied our proposed hy-

cross  
val-  
i-  
da-  
tion.  
The  
re-  
sults,  
sum-  
ma-  
rized  
in  
Ta-  
ble ??,  
demon-

113	brid
sub-	func-
jects,	tional
in-	PLS
clud-	re-
ing:	gres-
(i)	sion,
base-	along
line	with
renogram	other
curves;	re-
(ii)	gres-
post-	sion
furosemide	meth-
renogram	ods,
curves;	to
(iii)	the
or-	Emory
di-	re-
nal	nal
rat-	study
ings	data.
of	The
kid-	study
ney	col-
ob-	lected
struc-	data
tion	on
sta-	226
tus	kid-
(non-	neys
obstructed,	(left
equiv-	and
o-	right)
cal,	from

(v) or two subject-level variables (age and gender). The subjects had a mean age of 57.8 years (SD = 15.5; range = 18–83), with 54 males (48%) and 59 females (52%).

ob-structed) in-de-pen-dently as-sessed by three nu-clear medicine ex-perts; (iv) eight kidney-level phar-ma-coki-netic vari-ables de-rived from ra-dionu-clide imag-ing; and

	The	The
two		three
renogram		ex-
curves,		perts
(i)		unan-
and		i-
(ii),		mously
were		clas-
treated		si-
as		fied
func-		153
tional		kid-
pre-		neys
dic-		as
tors		non-
and		obstructed,
smoothed		5
us-		as
ing		equiv-
a		o-
B-		cal,
spline		and
ba-		40
sis		as
of		ob-
or-		structed,
der		while
15.		28
The		kid-
re-		neys
main-		had
ing		dis-
vari-		crepant
ables,		rat-
ex-		ings.

not including the redundant, no-sis, they were may treated as scalar additional predictors. Given the nature of these variables, we assume they are correlated with the renogram curves but

prediction performance by root mean squared error on the test data, normalized by the range of the test data response.

6.1 ...

error-  
aged  
and  
trans-  
formed  
us-  
ing  
a  
min-  
max  
logit  
trans-  
for-  
ma-  
tion.  
We  
split-  
ted  
the  
data  
into  
70%  
of  
train-  
ing  
data  
and  
30%  
of  
test-  
ing  
data,  
and  
evaluated  
the

*Sys-  
tems*

**154,**  
80–  
92.

Aguilera,  
A. M.,  
Es-  
cabras,  
M.,  
Preda,  
C.,  
and  
Saporta,  
G.  
(2010).

Us-  
ing  
ba-  
sis  
ex-  
pan-  
sions  
for  
es-  
ti-  
mat-  
ing  
func-  
tional  
PLS  
re-  
gres-  
sion:

## References

Aguilera,  
A. M.,  
Aguilera-  
Morillo,  
M. C.,  
and  
Preda,  
C.  
(2016).  
Pe-  
nal-  
ized  
ver-  
sions  
of  
func-  
tional  
PLS  
re-  
gres-  
sion.  
*Chemo-  
met-  
rics  
and  
In-  
tel-  
li-  
gent  
Lab-  
o-  
ra-  
tory*



	<i>Hybrid Partial Least Squares</i>
bust	Ap-
func-	pli-
tional	ca-
par-	tions
tial	with
least	chemo-
squares	met-
for	ric
scalar-	data.
on-	<i>Chemo-</i>
multiple-	<i>met-</i>
function	<i>rics</i>
re-	<i>and</i>
gres-	<i>In-</i>
sion.	<i>tel-</i>
<i>Jour-</i>	<i>li-</i>
<i>nal</i>	<i>gent</i>
<i>of</i>	<i>Lab-</i>
<i>Chemo-</i>	<i>o-</i>
<i>met-</i>	<i>ra-</i>
<i>rics</i>	<i>tory</i>
<b>36,</b>	<i>Sys-</i>
e3394.	<i>tems</i>
	<b>104,</b>
	289–
Beyaztas,	305.
U.	
and	
Shang,	Beyaztas,
H. L.	U.
(2020).	and
On	Lin Shang,
function-	H.
on-	(2022).
function	A
re-	ro-

- tional  
lin-  
ear  
re-  
gres-  
sion.  
*The*  
*An-*  
*nals*  
*of*  
*Statis-*  
*tics*  
**34**,  
2159–  
2179.
- Cardot,  
H.,  
Fer-  
raty,  
F.,  
and  
Sarda,  
P.  
(2003).  
Spline  
es-  
ti-  
ma-  
tors  
for  
the  
func-  
tional  
lin-
- gres-  
sion:  
Par-  
tial  
least  
squares  
ap-  
proach.  
*En-*  
*vi-*  
*ron-*  
*men-*  
*tal*  
*and*  
*Eco-*  
*log-*  
*i-*  
*cal*  
*Statis-*  
*tics*  
**27**,  
95–  
114.
- Cai,  
T. T.  
and  
Hall,  
P.  
(2006).  
Pre-  
dic-  
tion  
in  
func-



- González-Manteiga, W. (2017). Functional principal component regression and partial least squares regression: An overview and a comparative study. *In* P. Febrero-Bande, M., Galeano, P., and
- P. (2012). Methodology and theory for partial least squares regression applied to functional data. *The Annals of Statistics* **40**, 322–352.

tion	ter-
with	na-
Con-	tional
found-	Sta-
ing.	tis-
In	ti-
<i>Pro-</i>	<i>cal</i>
<i>ceed-</i>	<i>Re-</i>
<i>ings</i>	<i>view</i>
<i>of</i>	<b>85,</b>
<i>The</i>	61–
<i>35th</i>	83.
<i>Un-</i>	
<i>cer-</i>	
<i>tainty</i>	Geng,
<i>in</i>	S.,
<i>Ar-</i>	Ko-
<i>ti-</i>	lar,
<i>fi-</i>	M.,
<i>cial</i>	and
<i>In-</i>	Koyejo,
<i>tel-</i>	O.
<i>li-</i>	(2020).
<i>gence</i>	Joint
<i>Con-</i>	Non-
<i>fer-</i>	para-
<i>ence,</i>	met-
pages	ric
378–	Pre-
388.	ci-
PMLR.	sion
	Ma-
	trix
Goldsmith,	Es-
J.,	ti-
Bobb,	ma-

- Hall, J.,  
P. Crainiceanu,  
and C. M.,  
Horowitz, Caffo,  
J. L. B.,  
(2007).  
Method- Re-  
ol- ich,  
ogy D.  
and (2011).  
con- Pe-  
ver- nal-  
gence ized  
rates func-  
for tional  
func- re-  
tional gres-  
lin- sion.  
ear *Jour-*  
re- *nal*  
gres- *of*  
sion. *Com-*  
*The pu-*  
*An- ta-*  
*nals tional*  
*of and*  
*Statis- Graph-*  
*tics i-*  
**35,** *cal*  
70– *Statis-*  
91. *tics*  
**20,**  
830–  
Happ, 851.  
C.  
and



sis *Anal-*  
of *y-*  
hy- *sis,*  
brid *with*  
func- *an*  
tional *In-*  
and *tro-*  
vec- *duc-*  
tor *tion*  
data. *to*  
*Statis-* *Lin-*  
*tics* *ear*  
*in* *Op-*  
*Medicine* *er-*  
**40,** *a-*  
5152– *tors.*  
5173. Wi-  
ley,  
1st  
Kolar, edi-  
M., tion  
Liu, edi-  
H., tion.  
and  
Xing,  
E. P. Jang,  
(2014). J. H.  
Graph (2021).  
es- Prin-  
ti- ci-  
ma- pal  
tion com-  
from po-  
multi- nent  
attribute anal-  
data. y-





vari-  
 ance  
 struc-  
 ture  
 non-  
 para-  
 met-  
 ri-  
 cally  
 when  
 the  
 data  
 are  
 curves.  
*Jour-  
 nal  
 of  
 the  
 Royal  
 Sta-  
 tis-  
 ti-  
 cal  
 So-  
 ci-  
 ety:  
 Se-  
 ries  
 B*  
**53,**  
 233–  
 243.  
 Saricam,  
 S.,


*nal  
 of  
 the  
 Amer-  
 i-  
 can  
 Sta-  
 tis-  
 ti-  
 cal  
 As-  
 so-  
 ci-  
 a-  
 tion*  
**102,**  
 984–  
 996.  
 Rice,  
 J. A.  
 and  
 Sil-  
 ver-  
 man,  
 B. W.  
 (1991).  
 Es-  
 ti-  
 mat-  
 ing  
 the  
 mean  
 and  
 co-

	<i>Hybrid Partial Least Squares</i>
Silverman, B. W. (1996). Smoothed functional principal components by choice of norm.	Beyaztas, U., Asikgil, B., and Shang, H. L. (2022). On partial least-squares estimation of the scalar-on-function regression model.
<i>The Annals of Statistics</i>	<i>Journal of Chemometrics</i>
<b>24,</b> 1–24.	<b>36,</b> e3452.
Wang, Y. (2018). <i>Partial least squares meth-</i>	

in *ods*  
 func- *for*  
 tional *func-*  
 lin- *tional*  
 ear *re-*  
 re- *gres-*  
 gres- *sion*  
 sion. *mod-*  
*Journal* *els.*  
*of* PhD  
*Com-* the-  
*pu-* sis,  
*ta-* Uni-  
*tional* ver-  
*and* sity  
*Graph-* of  
*i-* North  
*cal* Car-  
*Statis-* olina  
*tics* at  
**21,** Chapel  
 600– Hill.  
 617.  
 Zhao,  
 Y.,  
 Zhu, Og-  
 H., den,  
 Strawn, R. T.,  
 N., and  
 and Reiss,  
 Dun- P. T.  
 son, (2012).  
 D. B. Wavelet-  
 (2016). based  
 Bayesian LASSO

**Tables** graph-  
**and** i-  
**Fig-** cal  
**ures** mod-  
els  
Table 1: for  
Ex- mul-  
am- ti-  
ple Ta- vari-  
ble ble  
Cap- ate  
tion.

Col1	Col2	Col3	func-
a	b	c	tional



*Re-*  
**Figure** *search*  
**1:** **17,**  
Ex- 1–  
am- 27.  
ple  
Fig-  
ure  
Cap-  
tion.