

Data-Integrated Simulation Science A
Elsa Bunz, Schmitt Group

Lecture 2: Modelling muscles and ligaments





University of Stuttgart
Institute for Modelling and Simulation
of Biomechanical Systems



SimTech

cbb Computational
Biophysics
and Biorobotics

What is required for such a model?



What is required for such a model?



What is required for such a model?

Model of high-level motor control

EP control

$$\Lambda_i^{\text{move}} = \{\vec{\lambda}_1, \vec{\lambda}_2, \vec{\lambda}_3, \dots, \vec{\lambda}_n\}$$

Joint space control

continuous vs. intermittent

model-based control

learning-based control

Model of low-level motor control

monosynaptic reflex


alpha-gamma co-activation

$$u_j = u_j^{\text{open}} + u_j^{\text{closed}}$$

$$u_j^{\text{closed}} = f_n(\kappa, \lambda_j^{\text{CE}}, l_j^{\text{CE}}, i_j^{\text{CE}})$$

Parameters

Initial conditions





Model of muscle-tendon unit


What we will look at today:

- ▶ Muscle physiology + morphology
- ▶ Microscopic vs. macroscopic model
- ▶ Muscle experiments
- ▶ Important muscle properties
- ▶ Hill-type muscle model

Muscle morphology + physiology



in Brickmann et al. (2012) mod. di Prampero (1985)



Video source: Hasudungan [2]

Microscopic modelling


Huxley-type muscle models

Andrew Fielding Huxley

Nobel prize 1963



- ▶ Explain the mechanism of force generation known as the "sliding filament theory" based on microstructure
- ▶ Look at one muscle in detail
- ▶ Require a lot of parameters
- ▶ Difficult to scale to a bigger system with several muscles



$$F = \frac{m \cdot s \cdot k}{2 \cdot l} \int_{-\infty}^{\infty} n(x) \cdot x \cdot dx$$

m: number of cross bridges per unit
s: sarcomere length
k: spring constant
l: distance between two M positions
x: elongation from rest length
 $n(x)$: Anzahl der gebundenen Zustände bei einer Auslenkung x

Huxley and Niedergerke [3]

Macroscopic modelling


Hill-type muscle models

Archibald Vivian Hill

Nobel prize 1922




- ▶ Include a contractile element which represents empirical data
- ▶ Phenomenological models
- ▶ Model the important characteristics
- ▶ Reduced number of necessary parameters
- ▶ Can be used in a system with several muscles




l^{CE} : length of contractile element (muscle fibre)

l^{MTU} : length of the muscle-tendon unit

Macroscopic modelling



Macroscopic modelling




Macroscopic modelling

simplified view

Excitation /
Contraction coupling

- (1) Electro-magnetic activation of the **motoneurons**
- (3) Electro-magnetic activation of the **muscle fibre (EMG)**
- (5) Bio-chemical activation of the thin **filaments (Ca^{2+})**



Activation Dynamics (Hatze)

Sought-after: a relation of the neural stimulation \mathbf{u} and muscle activity \mathbf{a}

$$\dot{\gamma} = m \cdot (u - \gamma) \quad \text{free Calcium ion concentration}$$


$$a_{(Hatze)}(l_{CE}\gamma) = \frac{a_0 + (\rho(l_{CE})\gamma)^3}{1 + (\rho(l_{CE})\gamma)^3}$$

with

$$\rho(l_{CE}) = c \cdot \eta \cdot \frac{(k-1)}{(k - \frac{l_{CE}}{l_{CE,opt}})} \frac{l_{CE}}{l_{CE,opt}}$$


After Hatze [4]. A simplification can be found in Rockenfeller and Günther [5].

Macroscopic modelling




Muscle experiments

Volume constancy




Erasistratus (304-250 BC)




Swammerdam (1663)

Muscle experiments


Quick release experiments




Hill [6, 7]



Muscle contraction



Heat production



Force-velocity-relation

Muscle experiments








Image sources: [8, 9]


Muscle experiments



Muscle experiments



Rassier, MacIntosh, and Herzog [11]



Gordon, Huxley, and Julian [12]







Image source: [13]



Gordon, Huxley, and Julian [12]


Muscle experiments



$$\text{Efficiency} = \frac{\text{power output}}{\text{power input}}$$

Günther and Schmitt [14]

Muscle experiments




Enthalpy rate = (mech. power output) + (maint. heat rate) + (contract. heat rate)


Günther and Schmitt [14]

Muscle properties


... that need to be reproduced in a macroscopic model




Force-length
relationship



Force-velocity
relation



Mechanical
efficiency




Enthalpy rate

$$\mu = \frac{\text{power output}}{\text{power input}}$$


$$\dot{H} = (\text{mech. power output}) + (\text{maint. heat rate}) + (\text{contract. heat rate})$$

Muscle properties


... that need to be reproduced in a macroscopic model




Force-length
relationship



Force-velocity
relation



Mechanical
efficiency





Enthalpy rate

$$\mu = \frac{\text{power output}}{\text{power input}}$$


$$\dot{H} = (\text{mech. power output}) + (\text{maint. heat rate}) + (\text{contract. heat rate})$$

Force-length relationship




Besides the muscle fibers also the surrounding soft tissue develops force leading to the difference in the force-length relationship of a sarcomere and a muscle.

Hill-type muscle model




Hill-type muscle model

Muscle-tendon unit (MTU)



Martynenko et al. [15] and Haeufle et al. [16]

Macroscopic modelling



Force of muscle-tendon unit F_{MTU}


1. Force equilibrium:

$$F_{CE} + F_{PEE} = F_{SEE} + F_{SDE}$$

2. Define equations for F_{CE} , F_{PEE} , F_{SEE} and F_{SDE} and rearrange to obtain equation for \dot{l}_{CE} .

3. Use \dot{l}_{CE} to calculate forces

4. $F_{MTU} = F_{SEE} + F_{SDE}$




Martynenko et al. [15] and Haeufle et al. [16]

Force of the contractile element F_{CE}

$$F_{CE} = A \cdot F_l \cdot F_v \cdot f_{max}$$


product approach

a
→ activity [%]



activation dynamics

Hill type
muscle model
Hill 1938



contraction dynamics
material laws

Force of the contractile element F_{CE}

Equations in Haeufle et al. [16]

$$F_{CE,c}(\dot{l}_{CE} \leq 0) = F_{\max} \left(\frac{qF_{isom} + A_{rel}}{1 - \frac{\dot{l}_{CE}}{B_{rel} l_{CE,opt}}} - A_{rel} \right)$$



with

$$F_{isom}(l_{CE}) = \exp \left\{ - \left| \frac{\frac{l_{CE}}{l_{CE,opt}} - 1}{\Delta W_{limb}} \right|^{v_{CE,limb}} \right\}$$

with activation q , maximum isometric force F_{\max} , F_{isom} representing the force-length relation and A_{rel}, B_{rel} being the normalized Hill parameters.

Benefits of biological muscles




Increased stability



Haeufle et al. [17], Haeufle, Grimmer, and Seyfarth [18], and Kalveram et al. [19]


Benefits of biological muscles

Reduced control effort





Benefits of biological muscles

Improved learning



Wochner et al. [21]

Summary + Discussion



Summary + Discussion

Do we miss/neglect any typical characteristics of a biological muscle?

Age

Reservoir chemical substrates

Fatigue

History effects

Training

???

Outlook

For the next exercise: Download the Matlab implementation of the muscle model of Häufle et al. (2014)



Journal of Biomechanics 47 (2014) 1531–1536

Contents lists available at ScienceDirect

Journal of Biomechanics

journal homepage: www.elsevier.com/locate/jbiomech
www.JBiomech.com



Short communication

Hill-type muscle model with serial damping and eccentric force–velocity relation



D.F.B. Häufle ^{a,*}, M. Günther ^{a,b}, A. Bayer ^a, S. Schmitt ^{a,c}

^a Universität Stuttgart, Institut für Sport- und Bewegungswissenschaft, Allmandring 28, D-70569 Stuttgart, Germany

^b Friedrich Schiller Universität, Institut für Sportwissenschaft, Seidelstrasse 20, D-07743 Jena, Germany

^c Universität Stuttgart, Stuttgart Research Centre for Simulation Technology (SRC SimTech), Pfaffenwaldring 5a, D-70569 Stuttgart, Germany

References

- [1] Syn Schmitt et al. "The dynamics of the skeletal muscle: A systems biophysics perspective on muscle modeling with the focus on Hill-type muscle models." 2019.
- [2] Armando Hasudungan. *Myology - Skeletal Muscle Contraction*. 2012.
- [3] Andrew F Huxley and Rolf Niedergerke. "Structural changes in muscle during contraction: interference microscopy of living muscle fibres." 1954.
- [4] H Hatze. "A myocybernetic control model of skeletal muscle." 1977.
- [5] Robert Rockenfeller and Michael Günther. "Inter-filament spacing mediates calcium binding to troponin: a simple geometric-mechanistic model explains the shift of force-length maxima with muscle activation" 2018.
- [6] Archibald Vivian Hill. *First and last experiments in muscle mechanics*. 1970.
- [7] Archibald Vivian Hill. "The heat of shortening and the dynamic constants of muscle." 1938.
- [8] OpenStax. *Types of Contraction*. 2016.
- [9] Pete Burridge. *The force-velocity relationship*.
- [10] Glenn K Klute et al. "Artificial muscles: Actuators for biorobotic systems." 2002.
- [11] Dilson E Rassier et al. "Length dependence of active force production in skeletal muscle." 1999.
- [12] AM Gordon et al. "The variation in isometric tension with sarcomere length in vertebrate muscle fibres." 1966.
- [13] Antony Crofts. *Muscle and motility*. 1996.
- [14] Michael Günther and Syn Schmitt. "A macroscopic ansatz to deduce the Hill relation." 2010.
- [15] Oleksandr V Martynenko et al. "Development and verification of a physiologically motivated internal controller for the open-source extended Hill-type muscle model in LS-DYNA." 2023.
- [16] DFB Haeufle et al. "Hill-type muscle model with serial damping and eccentric force–velocity relation." 2014.
- [17] DFB Haeufle et al. "Integration of intrinsic muscle properties, feed-forward and feedback signals for generating and stabilizing hopping." 2012.
- [18] DFB Haeufle et al. "The role of intrinsic muscle properties for stable hopping—stability is achieved by the force–velocity relation." 2010.
- [19] KarlTheodor Kalveram et al. "Energy management that generates terrain following versus apex-preserving hopping in man and machine." 2012.
- [20] Daniel FB Haeufle et al. "Muscles reduce neuronal information load: quantification of control effort in biological vs. robotic pointing and walking." 2020.
- [21] Isabell Wochner et al. "Learning with Muscles: Benefits for Data-Efficiency and Robustness in Anthropomorphic Tasks." 2023.
- [22] Isabell Wochner. "The benefit of muscle-actuated systems: internal mechanics, optimization and learning." 2023.