


PDE-Constrained Optimization for Multiscale Particle Dynamics

With Industrial Applications

Jonna Roden

University of Edinburgh, MIGSAA

Supervision by Dr Ben Goddard and Dr John Pearson



Introduction

There are many industrial processes that can be described by the integro-PDEs for multiscale particle dynamics and optimized using PDE-constrained optimization techniques. An example of such an industrial problem is minimizing the sedimentation time of yeast particles in beer to optimize the brewing process. The industrial partners of the PhD project are WEST Beer and uFraction8. Solving PDE-constrained optimization problems involving integro-PDE constraints requires the development of new theoretical and numerical methods, which is addressed in this work.

Multiscale Particle Dynamics

The dynamics of the particle density y at position x and time t can be described by the PDE, see [1]:

$$\begin{aligned} \partial_t y(x, t) &= \nabla^2 y(x, t) + u(x, t) + \alpha \nabla \cdot \int_{\Omega} y(x, t) y(x', t) \nabla V_2(|x - x'|) dx', & \text{in } \Omega \times [0, T], \\ \frac{\partial y(x, t)}{\partial n} + \int_{\Omega} y(x, t) y(x', t) \frac{\partial V_2(x)}{\partial n} dx' &= 0, & \text{on } \partial\Omega \times [0, T], \\ y(x, 0) &= y_0(x). \end{aligned}$$

The natural boundary condition for this problem is no-flux. However, other boundary conditions can be imposed, such as Dirichlet boundary conditions (used in Figure 2).

Optimal Control Problem

One PDE-constrained optimization problem involving the above particle dynamics is:

$$\min_{y, u} \frac{1}{2} \|y(x, t) - \hat{y}(x, t)\|_{L_2(\Omega \times [0, T])}^2 + \frac{\beta}{2} \|u(x, t)\|_{L_2(\Omega \times [0, T])}^2$$

subject to:

$$\begin{aligned} \partial_t y(x, t) &= \nabla^2 y(x, t) + u(x, t) + \alpha \nabla \cdot \int_{\Omega} y(x, t) y(x', t) \nabla V_2(|x - x'|) dx' \\ &+ \text{BCs} + \text{ICs for } y. \end{aligned}$$

In this example, the forcing term u in the PDE is the control variable. However, the control can be enforced in other ways, e.g. through boundary conditions.

First Order Optimality System

To find an optimal solution (\bar{y}, \bar{u}) to the above optimal control problem, first order optimality conditions are derived. The resulting system can be reduced to two coupled PDEs, that can be solved numerically:

Forward Problem

$$\partial_t y(x, t) = \Delta y(x, t) + \frac{1}{\beta} p(x, t) + \alpha \int_{\Omega} y(x, t) y(x', t) \nabla_x V_2(|x - x'|) dx',$$

Adjoint Equation

$$\begin{aligned} \partial_t p(x, \tau) &= \Delta p(x, \tau) - y(x, \tau) + \hat{y}(x, \tau) \\ &+ \alpha \int_{\Omega} \left(\nabla_x p(x, \tau) + \nabla_{x'} p(x', \tau) \right) y(x', \tau) \nabla_x V_2(|x - x'|) dx', \\ &+ \text{BCs} + \text{ICs for } y(x, t) \text{ and } p(x, \tau). \end{aligned}$$

The Adjoint Equation is running backward in time, with $\tau = T + t_0 - t$, for numerical stability.

Numerical Methods

New approaches to solving the optimality system are needed because of the non-linear, non-local nature of the particle interaction term in the PDE-constraint. Additionally, the boundary conditions are non-local and problem specific. Moreover, due to the final time condition in the adjoint equation, the problem is a boundary value problem in time as well as in space. For these reasons, standard methods are no longer sufficient to solve this type of problem.

Pseudospectral Methods

Pseudospectral methods on non-periodic domains are based on polynomial interpolation on non-equispaced points.

- Collocation points: Chebyshev points $\{x_j\}$ on $[-1, 1]$, defined as [2]:

$$x_j = \cos\left(\frac{j\pi}{N}\right), \quad j = 0, 1, \dots, N.$$

- Interpolation: Barycentric Lagrange interpolation, derived in [3].
- Differentiation: Chebyshev differentiation matrices, see [2].
- Integration: Clenshaw–Curtis quadrature, derived in [4].
- Advantages of pseudospectral methods: Can apply non-local boundary conditions easily, exponential convergence for smooth functions, see [5]:

$$\text{Pseudospectral Error} \approx O\left[\left(\frac{1}{N}\right)^N\right].$$

- Disadvantages of pseudospectral methods: complex domains are challenging to deal with, method involves (small) dense matrices.

Multiple Shooting Method [6]

- Discretize the time interval into subintervals $[t_i, t_{i+1}]$.
- Create initial guesses for y and p on each t_i .
- Solve the PDEs on the subintervals.
- Interpolate y and p between t_i and t_{i+1} when necessary.
- Match the PDE solution to the initial guess for y and p at each t_i by optimizing the initial guess.




Figure 1: Multiple Shooting for y and p .

Results

The optimality system can be solved using the multiple shooting approach and pseudospectral methods in time and space. The evolution of y and p are displayed in Figure 2, where each red dashed line represents a time step t_i , decreasing from the initial distribution at time t_0 to the target at time T . When $\alpha > 0$, the particles are interacting repulsively and the particles spread more than when $\alpha = 0$ and the particles are non-interacting. Note that in this example Dirichlet boundary conditions are imposed.




Figure 2: Solution to the optimality system without ($\alpha = 0$) and with ($\alpha > 0$) particle interaction term. Each red dashed line indicates a time step moving from the initial distribution at time t_0 to the target distribution at time T .

Forthcoming Research

In the future, the existing code library [7] will be extended to compute a wide range of PDE-constrained optimization problems for multiscale particle dynamics involving non-linear, non-local terms in one and two dimensions. The numerical method will be applied to industrial problems that aim at optimizing processes involving particle dynamics, such as nano filtration and sedimentation of yeast in beer. For more information on the applications of this method, please see the poster by Mildred Aduamoah.

References

- [1] M. Rex and H. Löwen. Dynamical density functional theory for colloidal dispersions including hydrodynamic interactions. *The European Physical Journal. E, Soft Matter*, 28:139–46, 2008.
- [2] L. N. Trefethen. *Spectral Methods in Matlab*. SIAM, 2000.
- [3] J. P. Berrut and L. N. Trefethen. Barycentric Lagrange interpolation. *SIAM Review*, 46(3):501–517, 2004.
- [4] C. W. Clenshaw and A. R. Curtis. A method for numerical integration on an automatic computer. *Numerische Mathematik*, 2(1):197–205, 1960.
- [5] J. P. Boyd. *Chebyshev and Fourier Spectral Methods*. Dover Publications, Inc, 2000.
- [6] T. Carraro, M. Geiger, and R. Rannacher. Indirect multiple shooting for nonlinear parabolic optimal control problems with control constraints. *SIAM Journal on Scientific Computing*, 36, 01 2014.
- [7] A. Nold, B. D. Goddard, P. Yatsyshin, N. Savva, and S. Kalliadas. Pseudospectral methods for density functional theory in bounded and unbounded domains. *Journal of Computational Physics*, 334:639–664, 2017.