

## Optimality Conditions for Two Species

We have the following set of forward equations:

$$\begin{aligned}\frac{\partial \rho_a}{\partial t} &= D_a \nabla^2 \rho_a - D_a \nabla \cdot (\rho_a F_a(\mathbf{w})) + D_a \nabla \cdot (\rho_a \nabla V_{ext,a}) + D_a \kappa \nabla \cdot \int_{\Omega} \rho_a(r) \rho_a(r') \mathbf{K}_{aa}(r, r') dr' \\ &\quad + D_a \tilde{\kappa} \nabla \cdot \int_{\Omega} \rho_a(r) \rho_b(r') \mathbf{K}_{ab}(r, r') dr' \\ \frac{\partial \rho_b}{\partial t} &= D_b \nabla^2 \rho_b - D_b \nabla \cdot (\rho_b F_b(\mathbf{w})) + D_b \nabla \cdot (\rho_b \nabla V_{ext,b}) + D_b \kappa \nabla \cdot \int_{\Omega} \rho_b(r) \rho_b(r') \mathbf{K}_{bb}(r, r') dr' \\ &\quad + D_b \tilde{\kappa} \nabla \cdot \int_{\Omega} \rho_b(r) \rho_a(r') \mathbf{K}_{ba}(r, r') dr',\end{aligned}$$

where  $D = \frac{1}{\gamma m}$ . No flux boundary conditions are:

$$\begin{aligned}&\left( D_a \nabla \rho_a - D_a \rho_a F_a(\mathbf{w}) + D_a \rho_a \nabla V_{ext,a} + D_a \kappa \int_{\Omega} \rho_a(r) \rho_a(r') \mathbf{K}_{aa}(r, r') dr' \right. \\ &\quad \left. + D_a \tilde{\kappa} \int_{\Omega} \rho_a(r) \rho_b(r') \mathbf{K}_{ab}(r, r') dr' \right) \cdot \mathbf{n} = 0 \\ &\left( D_b \nabla \rho_b - D_b \rho_b F_b(\mathbf{w}) + D_b \rho_b \nabla V_{ext,b} + D_b \kappa \int_{\Omega} \rho_b(r) \rho_b(r') \mathbf{K}_{bb}(r, r') dr' \right. \\ &\quad \left. + D_b \tilde{\kappa} \int_{\Omega} \rho_b(r) \rho_a(r') \mathbf{K}_{ba}(r, r') dr' \right) \cdot \mathbf{n} = 0\end{aligned}$$

The cost functional is:

$$J(\rho_a, \rho_b, \mathbf{w}) := \frac{1}{2} \|\rho_a - \hat{\rho}_a\|_{L_2(\Sigma)}^2 + \frac{\alpha}{2} \|\rho_b - \hat{\rho}_b\|_{L_2(\Sigma)}^2 + \frac{\beta}{2} \|\mathbf{w}\|_{L_2(\Sigma)}^2.$$

The Lagrangian is then:

$$\begin{aligned}
\mathcal{L}(\rho_a, \rho_b, \mathbf{w}, q_a, q_b) = & \frac{1}{2} \int_0^T \int_{\Omega} (\rho_a - \widehat{\rho}_a)^2 dr dt + \frac{\alpha}{2} \int_0^T \int_{\Omega} (\rho_b - \widehat{\rho}_b)^2 dr dt + \frac{\beta}{2} \int_0^T \int_{\Omega} \mathbf{w}^2 dr dt \\
& - \int_0^T \int_{\Omega} \left( \frac{\partial \rho_a}{\partial t} - D_a \nabla^2 \rho_a + D_a \nabla \cdot (\rho_a F_a(\mathbf{w})) - D_a \nabla \cdot (\rho_a \nabla V_{ext,a}) \right. \\
& \left. - D_a \kappa \nabla \cdot \int_{\Omega} \rho_a(r) \rho_a(r') \mathbf{K}_{aa}(r, r') dr' - D_a \tilde{\kappa} \nabla \cdot \int_{\Omega} \rho_a(r) \rho_b(r') \mathbf{K}_{ab}(r, r') dr \right) q_a dr dt \\
& - \int_0^T \int_{\Omega} \left( \frac{\partial \rho_b}{\partial t} - D_b \nabla^2 \rho_b + D_b \nabla \cdot (\rho_b F_b(\mathbf{w})) - D_b \nabla \cdot (\rho_b \nabla V_{ext,b}) \right. \\
& \left. - D_b \kappa \nabla \cdot \int_{\Omega} \rho_b(r) \rho_b(r') \mathbf{K}_{bb}(r, r') dr' - D_b \tilde{\kappa} \nabla \cdot \int_{\Omega} \rho_b(r) \rho_a(r') \mathbf{K}_{ba}(r, r') dr' \right) q_b dr dt \\
& - \int_0^T \int_{\partial\Omega} \left( D_a \nabla \rho_a - D_a \rho_a F_a(\mathbf{w}) + D_a \rho_a \nabla V_{ext,a} + D_a \kappa \int_{\Omega} \rho_a(r) \rho_a(r') \mathbf{K}_{aa}(r, r') dr' \right. \\
& \left. + D_a \tilde{\kappa} \int_{\Omega} \rho_a(r) \rho_b(r') \mathbf{K}_{ab}(r, r') dr' \right) \cdot \mathbf{n} q_a dr dt \\
& - \int_0^T \int_{\partial\Omega} \left( D_b \nabla \rho_b - D_b \rho_b F_b(\mathbf{w}) + D_b \rho_b \nabla V_{ext,b} + D_b \kappa \int_{\Omega} \rho_b(r) \rho_b(r') \mathbf{K}_{bb}(r, r') dr' \right. \\
& \left. + D_b \tilde{\kappa} \int_{\Omega} \rho_b(r) \rho_a(r') \mathbf{K}_{ba}(r, r') dr' \right) \cdot \mathbf{n} q_b dr dt
\end{aligned}$$

## 1 Adjoint 1

Taking the derivative with respect to  $\rho_a$  gives

$$\begin{aligned}
\mathcal{L}_{\rho_a}(\rho_a, \rho_b, \mathbf{w}, q_a, q_b)h = & \int_0^T \int_{\Omega} (\rho_a - \widehat{\rho}_a) h dr dt + \int_0^T \int_{\Omega} \left( - \frac{\partial h}{\partial t} q_a + D_a \nabla^2 h q_a - D_a \nabla \cdot (h F_a(\mathbf{w})) q_a \right. \\
& + D_a \nabla \cdot (h \nabla V_{ext,a}) q_a + D_a \kappa q_a \nabla \cdot \int_{\Omega} h(r) \rho_a(r') \mathbf{K}_{aa}(r, r') dr' \\
& + D_a \kappa q_a \nabla \cdot \int_{\Omega} \rho_a(r) h(r') \mathbf{K}_{aa}(r, r') dr' + D_a \tilde{\kappa} q_a \nabla \cdot \int_{\Omega} h(r) \rho_b(r') \mathbf{K}_{ab}(r, r') dr \\
& \left. + D_b \tilde{\kappa} q_b \nabla \cdot \int_{\Omega} \rho_b(r) h(r') \mathbf{K}_{ba}(r, r') dr' \right) dr dt \\
& - \int_0^T \int_{\partial\Omega} \left( D_a \nabla h - D_a h F_a(\mathbf{w}) + D_a h \nabla V_{ext,a} + D_a \kappa \int_{\Omega} h(r) \rho_a(r') \mathbf{K}_{aa}(r, r') dr' \right. \\
& + D_a \kappa \int_{\Omega} \rho_a(r) h(r') \mathbf{K}_{aa}(r, r') dr' + D_a \tilde{\kappa} \int_{\Omega} h(r) \rho_b(r') \mathbf{K}_{ab}(r, r') dr' \left. \right) \cdot \mathbf{n} q_a dr dt \\
& - \int_0^T \int_{\partial\Omega} D_b \tilde{\kappa} \int_{\Omega} \rho_b(r) h(r') \mathbf{K}_{ba}(r, r') dr' \cdot \mathbf{n} q_b dr dt
\end{aligned}$$

And so:

$$\begin{aligned}
\mathcal{L}_{\rho_a}(\rho_a, \rho_b, \mathbf{w}, q_a, q_b)h = & \int_0^T \int_{\Omega} (\rho_a - \widehat{\rho}_a) h dr dt + \int_0^T \int_{\Omega} \left( \frac{\partial q_a}{\partial t} h + D_a \nabla^2 q_a h + D_a \nabla q_a \cdot (h F_a(\mathbf{w})) \right. \\
& - D_a \nabla q_a \cdot (h \nabla V_{ext,a}) - D_a \kappa \nabla q_a(r) h(r) \int_{\Omega} \rho_a(r') \mathbf{K}_{aa}(r, r') dr' \\
& - D_a \kappa h(r) \int_{\Omega} \nabla q_a(r') \rho_a(r') \mathbf{K}_{aa}(r', r) dr' - D_a \tilde{\kappa} \nabla q_a h(r) \int_{\Omega} \rho_b(r') \mathbf{K}_{ab}(r, r') dr \\
& \left. - D_b \tilde{\kappa} h(r) \int_{\Omega} \nabla q_b(r') \rho_b(r') \mathbf{K}_{ba}(r', r) dr' \right) dr dt \\
& + \int_{\Omega} q_a(T) h(T) - q_a(0) h(0) dr \\
& + \int_0^T \int_{\Omega} \left( D_a \kappa h(r) \int_{\partial\Omega} q_a(r') \rho_a(r') \mathbf{K}_{aa}(r', r) dr' \cdot \mathbf{n} \right. \\
& \left. + D_b \tilde{\kappa} h(r) \int_{\partial\Omega} q_b(r') \rho_b(r') \mathbf{K}_{ba}(r', r) dr' \cdot \mathbf{n} \right) dr dt \\
& + \int_0^T \int_{\partial\Omega} D_a \frac{\partial h}{\partial n} q_a - D_a \frac{\partial q_a}{\partial n} h - D_a F_a(\mathbf{w}) h q_a \cdot \mathbf{n} + D_a \nabla V_{ext,a} h q_a \cdot \mathbf{n} dr dt \\
& + \int_0^T \int_{\partial\Omega} \left( D_a \kappa h(r) q_a(r) \int_{\Omega} \rho_a(r') \mathbf{K}_{aa}(r, r') \cdot \mathbf{n} dr' \right. \\
& \left. + D_a \tilde{\kappa} q_a(r) h(r) \int_{\Omega} \rho_b(r') \mathbf{K}_{ab}(r, r') \cdot \mathbf{n} dr' \right) dr dt \\
& - \int_0^T \int_{\partial\Omega} \left( D_a \nabla h q_{a,\partial\Omega} - D_a h F_a(\mathbf{w}) q_{a,\partial\Omega} + D_a h \nabla V_{ext,a} q_{a,\partial\Omega} \right. \\
& \left. + D_a \kappa q_{a,\partial\Omega}(r) h(r) \int_{\Omega} \rho_a(r') \mathbf{K}_{aa}(r, r') dr' + D_a \tilde{\kappa} q_{a,\partial\Omega} h(r) \int_{\Omega} \rho_b(r') \mathbf{K}_{ab}(r, r') dr' \right) \cdot \mathbf{n} dr dt \\
& - \int_0^T \int_{\Omega} \left( D_a \kappa h(r) \int_{\partial\Omega} q_{a,\partial\Omega}(r') \rho_a(r') \mathbf{K}_{aa}(r', r) dr' \right. \\
& \left. + D_b \tilde{\kappa} h(r) \int_{\Omega} q_{b,\partial\Omega}(r') \rho_b(r') \mathbf{K}_{ba}(r', r) dr' \right) \cdot \mathbf{n} dr dt
\end{aligned}$$

Then for  $\frac{\partial h}{\partial n} \neq 0$  we get;

$$\begin{aligned}
(D_a q_a - D_a q_{a,\partial\Omega}) \mathbf{n} &= \mathbf{0} \\
q_a &= q_{a,\partial\Omega}
\end{aligned}$$

And all boundary terms cancel so that we get:

$$\frac{\partial q_a}{\partial n} = 0 \quad \text{on} \quad \partial\Omega.$$

And we also get  $q_a(T) = 0$ .

We get:

$$\begin{aligned}\frac{\partial q_a}{\partial t} = & -D_a \nabla^2 q_a - \rho_a + \widehat{\rho}_a - D_a \nabla q_a \cdot F_a(\mathbf{w}) + D_a \nabla q_a \cdot \nabla V_{ext,a} \\ & + D_a \kappa \nabla q_a(r) \int_{\Omega} \rho_a(r') \mathbf{K}_{aa}(r, r') dr' + D_a \kappa \int_{\Omega} \nabla q_a(r') \rho_a(r') \mathbf{K}_{aa}(r', r) dr' \\ & + D_a \tilde{\kappa} \nabla q_a(r) \int_{\Omega} \rho_b(r') \mathbf{K}_{ab}(r, r') dr' + D_b \tilde{\kappa} \int_{\Omega} \nabla q_b(r') \rho_b(r') \mathbf{K}_{ba}(r', r) dr' .\end{aligned}$$

## 2 Adjoint 2

The second adjoint equation is almost equivalent to the first:

$$\begin{aligned}\frac{\partial q_b}{\partial t} = & -D_b \nabla^2 q_b - \alpha \rho_b + \alpha \widehat{\rho}_b - D_b \nabla q_b \cdot F_b(\mathbf{w}) + D_b \nabla q_b \cdot \nabla V_{ext,b} \\ & + D_b \kappa \nabla q_b(r) \int_{\Omega} \rho_b(r') \mathbf{K}_{bb}(r, r') dr' + D_b \kappa \int_{\Omega} \nabla q_b(r') \rho_b(r') \mathbf{K}_{bb}(r', r) dr' \\ & + D_b \tilde{\kappa} \nabla q_b \int_{\Omega} \rho_a(r') \mathbf{K}_{ba}(r, r') dr' + D_a \tilde{\kappa} \int_{\Omega} \nabla q_a(r') \rho_a(r') \mathbf{K}_{ab}(r', r) dr' .\end{aligned}$$

And the boundary condition is:

$$\frac{\partial q_b}{\partial n} = 0 \quad \text{on} \quad \partial\Omega.$$

And we also get  $q_b(T) = 0$ .

## 3 Gradient Equation

We consider the derivative of the Lagrangian with respect to  $\mathbf{w}$ . However, we will need to consider the Frechét derivative of terms involving  $F(\mathbf{w})$  first. From the definition from the Frechét derivative, we know that we have to consider the first order term of the Taylor expansion, so that we have:

$$F(\mathbf{w} + \mathbf{h}) - F(\mathbf{w}) = \nabla F(\mathbf{w}) \cdot \mathbf{h}.$$

However, this doesn't seem quite right. Consider from year end review:  $F(\rho) = \nabla \ln(\rho)$ . The resulting Frechét derivative was  $\nabla(\frac{h}{\rho})$ , which doesn't match with the above definition.