

PDE-Constrained Optimization for Multiscale Particle Dynamics

Jonna Roden


University of Edinburgh/MIGSAA

Joint work with Ben Goddard and John Pearson

April 21, 2021

Structure of the Talk

- ▶ PDE-Constrained Optimization
- ▶ Optimization for DDFT
- ▶ Numerical Methods
- ▶ Results



PDE-Constrained Optimization

A simple model

$$\min_{\rho, f} \quad \frac{1}{2} \|\rho - \hat{\rho}\|_{L_2(\Sigma)}^2 + \frac{\beta}{2} \|f\|_{L_2(\Sigma)}^2$$

subject to:

$$\partial_t \rho = \nabla^2 \rho + f \quad \text{in } \Sigma$$

BC and IC:

$$\frac{\partial \rho}{\partial n} = 0 \quad \text{on } \partial \Sigma$$

$$\rho(0, \vec{x}) = \rho_0(\vec{x})$$

PDE-Constrained Optimization

Deriving (first-order) optimality conditions

Define the Lagrangian \mathcal{L} :

$$\mathcal{L}(\rho, f, q) = \frac{1}{2} \|\rho - \hat{\rho}\|_{L_2(\Sigma)}^2 + \frac{\beta}{2} \|f\|_{L_2(\Sigma)}^2 - \int_{\Sigma} q \left(\partial_t \rho - \nabla^2 \rho - f \right) d\vec{x} dt - \int_{\partial \Sigma} q \frac{\partial \rho}{\partial n} d\vec{x} dt$$

Compute the directional derivatives:

$$\mathcal{L}_q(\rho^*, f^*, q)h = 0, \quad \mathcal{L}_{\rho}(\rho^*, f^*, q)h = 0, \quad \mathcal{L}_f(\rho^*, f^*, q)h = 0.$$

PDE-Constrained Optimization

Deriving (first-order) optimality conditions

Computing $\mathcal{L}_\rho(\rho^*, f^*, q)h$:

$$\begin{aligned}\mathcal{L}(\rho, f, q) = & \frac{1}{2} \int_{\Sigma} (\rho - \widehat{\rho})^2 d\vec{x} dt + \frac{\beta}{2} \int_{\Sigma} f^2 d\vec{x} dt - \int_{\Sigma} q \left(\partial_t \rho - \nabla^2 \rho - f \right) d\vec{x} dt \\ & - \int_{\partial \Sigma} q \frac{\partial \rho}{\partial n} d\vec{x} dt\end{aligned}$$

$$\begin{aligned}\mathcal{L}_\rho(\rho^*, f^*, q)h = & \int_{\Omega} (q(T)h(T) - q(0)h(0)) d\vec{x} - \int_{\Sigma} \left(h(-\rho + \widehat{\rho}) - h\partial_t q - h\nabla^2 q \right) d\vec{x} dt \\ & - \int_{\partial \Sigma} q \frac{\partial h}{\partial n} - q \frac{\partial h}{\partial n} + h \frac{\partial q}{\partial n} d\vec{x} dt\end{aligned}$$

PDE-Constrained Optimization

Deriving (first-order) optimality conditions

Computing $\mathcal{L}_\rho(\rho^*, f^*, q)h = 0$:

$$\mathcal{L}_\rho(\rho^*, f^*, q)h = \int_{\Omega} q(T)h(T)d\vec{x} - \int_{\Sigma} h \left(-\rho + \hat{\rho} - \partial_t q - \nabla^2 q \right) d\vec{x}dt - \int_{\partial\Sigma} h \frac{\partial q}{\partial n} d\vec{x}dt = 0$$

Adjoint equation:

$$\begin{aligned} \partial_t q &= -\nabla^2 q - \rho + \hat{\rho} && \text{in } \Sigma \\ \frac{\partial q}{\partial n} &= 0 && \text{on } \partial\Sigma \\ q(T) &= 0 \end{aligned}$$

PDE-Constrained Optimization

Deriving (first-order) optimality conditions

Computing $\mathcal{L}_f(\rho^*, f^*, q)h = 0$:

$$\begin{aligned}\mathcal{L}(\rho, f, q) = & \frac{1}{2} \int_{\Sigma} (\rho - \widehat{\rho})^2 d\vec{x} dt + \frac{\beta}{2} \int_{\Sigma} f^2 d\vec{x} dt - \int_{\Sigma} q \left(\partial_t \rho - \nabla^2 \rho - f \right) d\vec{x} dt \\ & - \int_{\partial \Sigma} q \frac{\partial \rho}{\partial n} d\vec{x} dt\end{aligned}$$

$$\mathcal{L}_f(\rho^*, f^*, q)h = \int_{\Sigma} h(\beta f + q) d\vec{x} dt = 0$$

Gradient equation:

$$f = -\frac{1}{\beta} q$$

PDE-Constrained Optimization

The (first-order) optimality system

$$\partial_t \rho = \nabla^2 \rho + f$$

$$\partial_t q = -\nabla^2 q - \rho + \hat{\rho}$$

$$f = -\frac{1}{\beta} q$$

$$\rho(0, \vec{x}) = \rho_0(\vec{x}), \quad q(T, \vec{x}) = 0, \quad +\text{BCs}$$

PDE-Constrained Optimization

A simple model

$$\min_{\rho, f} \quad \frac{1}{2} \|\rho - \hat{\rho}\|_{L_2(\Sigma)}^2 + \frac{\beta}{2} \|f\|_{L_2(\Sigma)}^2$$

subject to:

$$\partial_t \rho = \nabla^2 \rho + f \quad \text{in } \Sigma$$

BC and IC:

$$\frac{\partial \rho}{\partial n} = 0 \quad \text{on } \partial \Sigma$$

$$\rho(0, \vec{x}) = \rho_0(\vec{x})$$

Optimization for DDFT

A (simple) DDFT model

$$\min_{\rho, \vec{w}} \quad \frac{1}{2} \|\rho - \hat{\rho}\|_{L_2(\Sigma)}^2 + \frac{\beta}{2} \|\vec{w}\|_{L_2(\Sigma)}^2$$

subject to:

$$\partial_t \rho = \nabla^2 \rho - \nabla \cdot (\rho \vec{w}) \quad \text{in } \Sigma$$

BC and IC:

$$\frac{\partial \rho}{\partial n} - \rho \vec{w} \cdot \vec{n} = 0 \quad \text{on } \partial \Sigma$$

$$\rho(0, \vec{x}) = \rho_0(\vec{x})$$

Optimization for DDFT

A (simple) DDFT model

$$\min_{\rho, \vec{w}} \quad \frac{1}{2} \|\rho - \hat{\rho}\|_{L_2(\Sigma)}^2 + \frac{\beta}{2} \|\vec{w}\|_{L_2(\Sigma)}^2$$

subject to:

$$\partial_t \rho = \nabla^2 \rho - \nabla \cdot (\rho \vec{w}) + \nabla \cdot \int_{\Omega} \rho(\vec{x}) \rho(\vec{x}') \nabla V_2(|\vec{x} - \vec{x}'|) d\vec{x}' \quad \text{in } \Sigma$$

BC and IC:

$$\frac{\partial \rho}{\partial n} - \rho \vec{w} \cdot \vec{n} + \int_{\Omega} \rho(\vec{x}) \rho(\vec{x}') \frac{\partial V_2}{\partial n}(|\vec{x} - \vec{x}'|) d\vec{x}' = 0 \quad \text{on } \partial \Sigma$$
$$\rho(0, \vec{x}) = \rho_0(\vec{x})$$

Optimization for DDFT

The (first-order) optimality system

$$\begin{aligned}\partial_t \rho &= \nabla^2 \rho - \nabla \cdot (\rho \vec{w}) + \nabla \cdot \int_{\Omega} \rho(\vec{x}) \rho(\vec{x}') \nabla V_2(|\vec{x} - \vec{x}'|) d\vec{x}' \\ \partial_t q &= -\nabla^2 q - \nabla q \cdot \vec{w} - \rho + \hat{\rho} \\ &\quad + \int_{\Omega} \rho(\vec{x}') \left(\nabla_{\vec{x}} q(\vec{x}) - \nabla_{\vec{x}'} q(\vec{x}') \right) \cdot \nabla V_2(|\vec{x} - \vec{x}'|) d\vec{x}' \\ \vec{w} &= -\frac{1}{\beta} \rho \nabla q\end{aligned}$$

$$\rho(0, \vec{x}) = \rho_0(\vec{x}), \quad q(T, \vec{x}) = 0, \quad + \text{BCs}$$

Optimization for DDFT

Problem: Negative diffusion term in q causes numerical instability.

Solution: Change of time variable for this PDE: $\tau = T - t$.

$$\partial_t \rho(t, \vec{x}) = \nabla^2 \rho(t, \vec{x}) - \nabla \cdot (\rho(t, \vec{x}) \vec{w}(t, \vec{x})) + \nabla \cdot \int_{\Omega} \rho(t, \vec{x}) \rho(t, \vec{x}') \nabla V_2(|\vec{x} - \vec{x}'|) d\vec{x}'$$

$$\begin{aligned} \partial_{\tau} q(\tau, \vec{x}) = & \nabla^2 q(\tau, \vec{x}) + \nabla q(\tau, \vec{x}) \cdot \vec{w}(\tau, \vec{x}) + \rho(\tau, \vec{x}) - \hat{\rho}(\tau, \vec{x}) \\ & - \int_{\Omega} \rho(\tau, \vec{x}') \left(\nabla_{\vec{x}} q(\tau, \vec{x}) - \nabla_{\vec{x}'} q(\tau, \vec{x}') \right) \cdot \nabla V_2(|\vec{x} - \vec{x}'|) d\vec{x}' \end{aligned}$$

$$\vec{w}(t, \vec{x}) = -\frac{1}{\beta} \rho(t, \vec{x}) \nabla q(t, \vec{x})$$

$$\rho(0, \vec{x}) = \rho_0(\vec{x}), \quad q(0, \vec{x}) = 0, \quad + \text{BCs}$$

Numerical Methods

- ▶ Challenge 1: Particle interaction term is nonlinear and nonlocal (+ nonlocal BCs). How to avoid shortcomings of standard methods (FEM/FDM)?

⇒ **Pseudospectral methods**

- ▶ Challenge 2: One PDE is forward in time, the other backward. How to do time stepping?

⇒ **Fixed point algorithm**

Numerical Methods

Pseudospectral Methods

- ▶ Reduce both PDEs to systems of ODEs.
- ▶ Discretize time (accurate interpolation).
- ▶ Equations can now be solved using a DAE solver (when given all necessary inputs).

Numerical Methods

Fixed point algorithm

Initialize with guess $\vec{w}^{(0)}$.

1. Solve $\partial_t \rho = \nabla^2 \rho - \nabla \cdot (\rho \vec{w}^{(i)}) + \nabla \cdot \int_{\Omega} \rho(\vec{x}) \rho(\vec{x}') \nabla V_2(|\vec{x} - \vec{x}'|) d\vec{x}'$.

2. Solve $\partial_{\tau} q = \nabla^2 q + \nabla q \cdot \vec{w}^{(i)} + \rho^{(i)} - \hat{\rho} - \int_{\Omega} \rho^{(i)}(\vec{x}') \left(\nabla q(\vec{x}) - \nabla q(\vec{x}') \right) \cdot \nabla V_2(|\vec{x} - \vec{x}'|) d\vec{x}'$.

3. Solve $\vec{w}_g^{(i)} = -\frac{1}{\beta} \rho^{(i)} \nabla q^{(i)}$.

4. Measure the error: $\mathcal{E} = \|\vec{w}^{(i)} - \vec{w}_g^{(i)}\|$.

5. Update control, with $\lambda \in [0, 1]$: $\vec{w}^{(i+1)} = (1 - \lambda) \vec{w}^{(i)} + \lambda \vec{w}_g^{(i)}$.

Iterate until $\mathcal{E} < TOL$.

Optimization for DDFT

A more general DDFT model

$$\min_{\rho, \vec{w}} \quad \frac{1}{2} \|\rho - \hat{\rho}\|_{L_2(\Sigma)}^2 + \frac{\beta}{2} \|\vec{w}\|_{L_2(\Sigma)}^2$$

subject to:

$$\partial_t \rho = \nabla \cdot \left(\rho \nabla \frac{\delta \mathcal{F}[\rho]}{\delta \rho} - \rho \vec{w} \right) := \nabla \cdot \vec{j} \quad \text{in } \Sigma$$

BC and IC:

$$\vec{j} \cdot \vec{n} = 0 \quad \text{on } \partial \Sigma$$

$$\rho(0, \vec{x}) = \rho_0(\vec{x})$$

Optimization for DDFT

Reminder: (Simple) DDFT model

$$\min_{\rho, \vec{w}} \quad \frac{1}{2} \|\rho - \hat{\rho}\|_{L_2(\Sigma)}^2 + \frac{\beta}{2} \|\vec{w}\|_{L_2(\Sigma)}^2$$

subject to:




$$\partial_t \rho = \nabla^2 \rho - \nabla \cdot (\rho \vec{w}) + \nabla \cdot \int_{\Omega} \rho(\vec{x}) \rho(\vec{x}') \nabla V_2(|\vec{x} - \vec{x}'|) d\vec{x}' \quad \text{in } \Sigma$$

BC and IC:

$$\frac{\partial \rho}{\partial n} - \rho \vec{w} \cdot \vec{n} + \int_{\Omega} \rho(\vec{x}) \rho(\vec{x}') \frac{\partial V_2}{\partial n}(|\vec{x} - \vec{x}'|) d\vec{x}' = 0 \quad \text{on } \partial \Sigma$$
$$\rho(0, \vec{x}) = \rho_0(\vec{x})$$


Results

Overall Cost: $\mathcal{J} = \frac{1}{2} \|\rho - \hat{\rho}\|_{L_2(\Sigma)}^2 + \frac{\beta}{2} \|\vec{w}\|_{L_2(\Sigma)}^2$, $\mathcal{J}_{\vec{w}=\vec{0}} = 0.0130$.



Results

Overall Cost: $\mathcal{J} = \frac{1}{2}\|\rho - \hat{\rho}\|_{L_2(\Sigma)}^2 + \frac{\beta}{2}\|\vec{w}\|_{L_2(\Sigma)}^2$, $\mathcal{J}_{\vec{w}=\vec{0}} = 0.0130$, $\mathcal{J}_{opt} = 7.2994 \times 10^{-4}$.




Current work



Next steps

Industrial partners of the PhD



Summary

Up to now:

- ▶ Deriving PDE-constrained optimization models.
- ▶ Developing a suitable numerical method to solve them.




Current:

- ▶ Complex domains.
- ▶ Extended models (e.g. sedimentation, multiple species).
- ▶ Different boundary conditions.


Up next:

- ▶ Application of the method to other extended models.
- ▶ Application of the numerical framework to industrial processes.

References

-  M. Aduamoah, B. D. Goddard, J. W. Pearson and J. C. Roden.
PDE-constrained optimization models and pseudospectral methods for multiscale particle dynamics.
Preprint, 2020.
-  M. Burger, M. Di Francesco, P.A. Markowich and M.-T. Wolfram.
Mean field games with nonlinear mobilities in pedestrian dynamics.
Discrete and Continuous Dynamical Systems - Series B, 19(5), 1311-1333, 2014.
-  A. Nold, B.D. Goddard, P. Yatsyshin, N. Savva and S. Kalliadasis.
Pseudospectral methods for density functional theory in bounded and unbounded domains.
Journal of Computational Physics, 334, 639-664, 2017.
<https://datashare.is.ed.ac.uk/handle/10283/2647> (2DChebClass)

References: Figures

-  Bacteria. Digital Image.
USCNews. 12 February 2008, <https://news.usc.edu/135660/how-bacteria-adapt-to-hostile-environments/>
-  Red and White Bloodcells. Digital Image.
The Franklin Institute. <https://www.fi.edu/heart/white-blood-cells>
-  ufraction8 Logo. Digital Image.
www.ufraction8. ufraction8.com
-  WEST Logo. Digital Image.
WEST Brewery www.westbeer.com