# FLS 6441 - Methods III: Explanation and Causation

Week 9 - Controlling for Confounding

Jonathan Phillips

May 2019

# Classification of Research Designs

		Independence of Treatment Assignment	Researcher Controls Treatment Assignment?
Controlled Experiments	Field Experiments	✓	✓
	Survey and Lab Experiments	√	√
Natural Experiments	Natural Experiments	√	
	Instrumental Variables	√	
	Discontinuities	√	
Observational Studies	Difference-in-Differences		
	Controlling for Confounding		
	Matching		
	Comparative Cases and Process Tracing		

# Section 1

► Adding control variables changes the comparison we are making: Treatment is associated with higher values of the Outcome...holding constant the values of X

- ► Adding control variables changes the comparison we are making: Treatment is associated with higher values of the Outcome...holding constant the values of X
- ► But when does controlling allow us to move from "is associated with" to causes?

► Up until now causal estimates have required treatment assignment to be **independent of potential outcomes** 

- Up until now causal estimates have required treatment assignment to be independent of potential outcomes
- But it's also acceptable if treatment assignment is conditionally independent of potential outcomes

- Up until now causal estimates have required treatment assignment to be independent of potential outcomes
- But it's also acceptable if treatment assignment is conditionally independent of potential outcomes

$$(y_0,y_1)\perp D|X$$

- Up until now causal estimates have required treatment assignment to be independent of potential outcomes
- But it's also acceptable if treatment assignment is conditionally independent of potential outcomes

$$(y_0,y_1)\perp D|X$$

► After controlling for *X*, treatment is independent of potential outcomes: No confounding

- Up until now causal estimates have required treatment assignment to be independent of potential outcomes
- But it's also acceptable if treatment assignment is conditionally independent of potential outcomes

$$(y_0, y_1) \perp D|X$$

- ► After controlling for *X*, treatment is independent of potential outcomes: No confounding
- 'No unmeasured confounders'

- Up until now causal estimates have required treatment assignment to be independent of potential outcomes
- But it's also acceptable if treatment assignment is conditionally independent of potential outcomes

$$(y_0, y_1) \perp D|X$$

- ► After controlling for X, treatment is independent of potential outcomes: No confounding
- 'No unmeasured confounders'
- ► This is an assumption

- ▶ Up until now causal estimates have required treatment assignment to be independent of potential outcomes
- But it's also acceptable if treatment assignment is conditionally independent of potential outcomes

$$(y_0,y_1)\perp D|X$$

- ► After controlling for *X*, treatment is independent of potential outcomes: No confounding
- 'No unmeasured confounders'
- ► This is an assumption
  - We cannot directly test it

- Up until now causal estimates have required treatment assignment to be independent of potential outcomes
- But it's also acceptable if treatment assignment is conditionally independent of potential outcomes

$$(y_0, y_1) \perp D|X$$

- ► After controlling for X, treatment is independent of potential outcomes: No confounding
- 'No unmeasured confounders'
- ► This is an assumption
  - We cannot directly test it
  - We have to make an argument and provide supporting evidence

▶ If we exclude a covariate, we bias our regression estimate:

- ▶ If we exclude a covariate, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

- ▶ If we exclude a covariate, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► If we try this 'wrong' regression with a missing confounder:

- ▶ If we exclude a covariate, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

If we try this 'wrong' regression with a missing confounder:

$$Y_i = \alpha + \beta D_i + \epsilon_i$$

- ▶ If we exclude a covariate, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► If we try this 'wrong' regression with a missing confounder:

$$Y_i = \alpha + \beta D_i + \epsilon_i$$

- ▶ If we exclude a covariate, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► If we try this 'wrong' regression with a missing confounder:

$$Y_i = \alpha + \beta D_i + \epsilon_i$$

$$X_i = \psi + \delta D_i + \epsilon_i$$

- ▶ If we exclude a covariate, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► If we try this 'wrong' regression with a missing confounder:

$$Y_i = \alpha + \beta D_i + \epsilon_i$$

$$X_i = \psi + \delta D_i + \epsilon_i$$

$$Y_i = \alpha + \beta D_i + \gamma (\psi + \delta D_i + \epsilon_i) + \epsilon_i$$

- ▶ If we exclude a covariate, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► If we try this 'wrong' regression with a missing confounder:

$$Y_i = \alpha + \beta D_i + \epsilon_i$$

$$X_i = \psi + \delta D_i + \epsilon_i$$

$$Y_i = \alpha + \beta D_i + \gamma (\psi + \delta D_i + \epsilon_i) + \epsilon_i$$

$$Y_i = \alpha + D_i(\beta + \gamma \delta) + \gamma(\psi + \epsilon_i) + \epsilon_i$$

- ▶ If we exclude a covariate, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► If we try this 'wrong' regression with a missing confounder:  $Y_i = \alpha + \beta D_i + \epsilon_i$ 

► What happens to our coefficient estimate?  $X_i = \psi + \delta D_i + \epsilon_i$ 

$$Y_i = \alpha + \beta D_i + \gamma (\psi + \delta D_i + \epsilon_i) + \epsilon_i$$

$$Y_i = \alpha + D_i(\beta + \gamma \delta) + \gamma(\psi + \epsilon_i) + \epsilon_i$$

► So the coefficient we estimate is wrong by this amount:

- ▶ If we exclude a covariate, we bias our regression estimate:
- ► 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► If we try this 'wrong' regression with a missing confounder:  $Y_i = \alpha + \beta D_i + \epsilon_i$ 

► What happens to our coefficient estimate?  $X_i = \psi + \delta D_i + \epsilon_i$ 

$$Y_{i} = \alpha + \beta D_{i} + \gamma (\psi + \delta D_{i} + \epsilon_{i}) + \epsilon_{i}$$

$$Y_i = \alpha + D_i(\beta + \gamma \delta) + \gamma(\psi + \epsilon_i) + \epsilon_i$$

► So the coefficient we estimate is wrong by this amount:

► What does controlling **do**?

- ► What does controlling **do**?
  - ► It means **removing the variation** in the data due to the confounder

- ▶ What does controlling do?
  - ► It means **removing the variation** in the data due to the confounder
  - ► Equivalently, it means separating our data for each value of the confounder: **Subclassification**

- ▶ What does controlling do?
  - It means removing the variation in the data due to the confounder
  - Equivalently, it means separating our data for each value of the confounder: Subclassification
  - ► Then, within each group, the confounder is **constant** and can't affect the relationship between *D* and *Y*.

- ▶ What does controlling do?
  - It means removing the variation in the data due to the confounder
  - Equivalently, it means separating our data for each value of the confounder: Subclassification
  - ► Then, within each group, the confounder is **constant** and can't affect the relationship between *D* and *Y*.
  - We have created balance between the treated and control groups on all other characteristics

- ► We receive a dataset with twenty variables: *D*, *Y* and 18 more.
  - ► Which variables should we include as controls?

- ► We receive a dataset with twenty variables: *D*, *Y* and 18 more.
  - ► Which variables should we include as controls?
- ► We usually do **NOT** need to include all of them

- ▶ We receive a dataset with twenty variables: D, Y and 18 more.
  - ► Which variables should we include as controls?
- ► We usually do **NOT** need to include all of them
  - ► Only the variables **necessary** to stop confounding

- ▶ We receive a dataset with twenty variables: D, Y and 18 more.
  - ► Which variables should we include as controls?
- ► We usually do **NOT** need to include all of them
  - ► Only the variables **necessary** to stop confounding
  - ► Including unnecessary variables also produces bias

- ▶ We receive a dataset with twenty variables: D, Y and 18 more.
  - ► Which variables should we include as controls?
- ► We usually do **NOT** need to include all of them
  - ► Only the variables **necessary** to stop confounding
  - Including unnecessary variables also produces bias
  - ► We lose power (degrees of freedom) for every control we add

- ▶ We receive a dataset with twenty variables: D, Y and 18 more.
  - ► Which variables should we include as controls?
- ▶ We usually do NOT need to include all of them
  - ► Only the variables **necessary** to stop confounding
  - Including unnecessary variables also produces bias
  - ► We lose power (degrees of freedom) for every control we add
  - And additional variables reduce overlap (increase model-dependence)

- ▶ We receive a dataset with twenty variables: D, Y and 18 more.
  - ► Which variables should we include as controls?
- ▶ We usually do NOT need to include all of them
  - ► Only the variables **necessary** to stop confounding
  - Including unnecessary variables also produces bias
  - ► We lose power (degrees of freedom) for every control we add
  - And additional variables reduce overlap (increase model-dependence)

## Section 2

Which Variables to Control For

➤ To know which variables to control for, it helps to draw a causal diagram

- ➤ To know which variables to control for, it helps to draw a causal diagram
  - ► A **Directed Acyclical Graph** (DAG)

- ➤ To know which variables to control for, it helps to draw a causal diagram
  - ► A **Directed Acyclical Graph** (DAG)
  - Arrows only in one direction

- ► To know which variables to control for, it helps to draw a causal diagram
  - ► A **Directed Acyclical Graph** (DAG)
  - Arrows only in one direction
  - No circular loops!

- ► Causation is like **Water**, flowing along the graph
  - We want to focus on one 'flow' of causation from treatment to outcomes

- ► Causation is like **Water**, flowing along the graph
  - We want to focus on one 'flow' of causation from treatment to outcomes
  - Avoiding mixing with the other flows of causation in the network

- ► Three Rules to achieve Conditional Independence:
  - Include as controls enough variables to block all back-door paths from treatment to the outcome

- ► Three Rules to achieve Conditional Independence:
  - Include as controls enough variables to block all back-door paths from treatment to the outcome
  - 2. Exclude any variables that are **post-treatment**

- ► Three Rules to achieve Conditional Independence:
  - Include as controls enough variables to block all back-door paths from treatment to the outcome
  - 2. Exclude any variables that are **post-treatment**
  - 3. Exclude any variables that are colliders

► To identify back-door paths:

- ► To identify back-door paths:
  - Start with an arrow pointing at treatment

- ► To identify back-door paths:
  - Start with an arrow pointing at treatment
  - ► Trace the path 'backwards' (the direction of the arrows doesn't matter) until you reach treatment


- ► To identify back-door paths:
  - Start with an arrow pointing at treatment
  - ► Trace the path 'backwards' (the direction of the arrows doesn't matter) until you reach treatment
  - ► Repeat for every possible path from treatment to outcome


- ► To identify back-door paths:
  - Start with an arrow pointing at treatment
  - ► Trace the path 'backwards' (the direction of the arrows doesn't matter) until you reach treatment
  - ► Repeat for every possible path from treatment to outcome
- Block back-door paths by controlling for any variable along the path


- ► To identify back-door paths:
  - Start with an arrow pointing at treatment
  - ► Trace the path 'backwards' (the direction of the arrows doesn't matter) until you reach treatment
  - ► Repeat for every possible path from treatment to outcome
- Block back-door paths by controlling for any variable along the path
- Identify the Minimum set of controls to achieve conditional independence


- ► To identify back-door paths:
  - Start with an arrow pointing at treatment
  - ► Trace the path 'backwards' (the direction of the arrows doesn't matter) until you reach treatment
  - Repeat for every possible path from treatment to outcome
- Block back-door paths by controlling for any variable along the path
- ► Identify the **Minimum set of controls** to achieve conditional independence
  - ► Any set of variables which blocks All back-door paths

- ► To identify back-door paths:
  - Start with an arrow pointing at treatment
  - ► Trace the path 'backwards' (the direction of the arrows doesn't matter) until you reach treatment
  - Repeat for every possible path from treatment to outcome
- Block back-door paths by controlling for any variable along the path
- ► Identify the **Minimum set of controls** to achieve conditional independence
  - ► Any set of variables which blocks All back-door paths
  - ► Include these as control variables in our regression











► Including **post-treatment** variables will introduce bias

- Including post-treatment variables will introduce bias
  - Because variables measured 'after' treatment can also be affected by treatment

- Including post-treatment variables will introduce bias
  - Because variables measured 'after' treatment can also be affected by treatment
  - ► They're not confounders, but **mediating variables**

- Including post-treatment variables will introduce bias
  - Because variables measured 'after' treatment can also be affected by treatment
  - ► They're not confounders, but **mediating variables**
  - Controlling for them changes the definition of the causal effect we are estimating





► Colliders are variables on back-door paths which have arrows pointing both into them and out of them

- ► Colliders are variables on back-door paths which have arrows pointing both into them and out of them
- ► The water 'collides' in both directions and 'cancels out'

- ► Colliders are variables on back-door paths which have arrows pointing both into them and out of them
- ► The water 'collides' in both directions and 'cancels out'
- ▶ But if we do 'control' for a collider we *introduce* a bias in the relationship between *D* and *Y*

- ► Colliders are variables on back-door paths which have arrows pointing both into them and out of them
- ► The water 'collides' in both directions and 'cancels out'
- ► But if we do 'control' for a collider we *introduce* a bias in the relationship between *D* and *Y*
- ► So we must avoid controlling for them

- ► Colliders are variables on back-door paths which have arrows pointing both into them and out of them
- ▶ The water 'collides' in both directions and 'cancels out'
- ► But if we do 'control' for a collider we *introduce* a bias in the relationship between *D* and *Y*
- ► So we must avoid controlling for them
- ► Hard!





Figure 3.4: Simulation of conditional dependence within values of a collider variable.



### Example adapted from MW, p.72

- 1. List all of the **back-door paths** from *D* to *Y*
- 2. Identify any **post-treatment** variables: Do NOT include as controls
- Identify any back-door paths with collider variables: Mark these as already blocked
- 4. Find a minimum set of variables that blocks all remaining back-door paths
- 5. Double-check your minimum set of control variables does not contain any post-treatment or collider variables

