FLS 6441 - Methods III: Explanation and Causation

Week 9 - Controlling for Confounding

Jonathan Phillips

April 2020

Classification of Research Designs

		Independence of Treatment Assignment	Researcher Controls Treatment Assignment?
Controlled Experiments	Field Experiments	✓	✓
	Survey and Lab Experiments	√	√
Natural Experiments	Natural Experiments	√	
	Instrumental Variables	√	
	Discontinuities	√	
Observational Studies	Difference-in-Differences		
	Controlling for Confounding		
	Matching		
	Comparative Cases and Process Tracing		

Section 1

► What if we don't have repeated observations over time for the same units?

- ► What if we don't have repeated observations over time for the same units?
- ▶ Or what if *everyone* is treated at the same point in time?

- ► What if we don't have repeated observations over time for the same units?
- Or what if everyone is treated at the same point in time?
- ► We cannot use Difference-in-Differences

- What if we don't have repeated observations over time for the same units?
- Or what if everyone is treated at the same point in time?
- ► We cannot use Difference-in-Differences
- ► For cross-sectional observational studies, the next-best alternative is...

- What if we don't have repeated observations over time for the same units?
- Or what if everyone is treated at the same point in time?
- We cannot use Difference-in-Differences
- ► For cross-sectional observational studies, the next-best alternative is...
- ▶ Controls!

- ► What we know: Adding control variable X changes the comparison we are making:
 - ► Treatment is associated with higher values of the Outcome...for units with the same values of X

- ► What we know: Adding control variable X changes the comparison we are making:
 - ► Treatment is associated with higher values of the Outcome...for units with the same values of X
- What we don't yet know: When does controlling allow us to say:
 - ► Treatment *causes* higher values of the Outcome?

▶ Up until now causal estimates have required treatment assignment to be independent of potential outcomes

 Up until now causal estimates have required treatment assignment to be independent of potential outcomes

$$(y_0,y_1)\perp D$$

Up until now causal estimates have required treatment assignment to be independent of potential outcomes

$$(y_0,y_1)\perp D$$

 Up until now causal estimates have required treatment assignment to be independent of potential outcomes

$$(y_0,y_1)\perp D$$

$$(y_0, y_1) \perp D|X$$

Up until now causal estimates have required treatment assignment to be independent of potential outcomes

$$(y_0,y_1)\perp D$$

But it's also acceptable if treatment assignment is conditionally independent of potential outcomes

$$(y_0, y_1) \perp D|X$$

► After controlling for X, treatment is independent of potential outcomes: 'No unmeasured confounders'

 Up until now causal estimates have required treatment assignment to be independent of potential outcomes

$$(y_0,y_1)\perp D$$

$$(y_0, y_1) \perp D|X$$

- ► After controlling for X, treatment is independent of potential outcomes: 'No unmeasured confounders'
- ► This is an assumption

Up until now causal estimates have required treatment assignment to be independent of potential outcomes

$$(y_0,y_1)\perp D$$

$$(y_0, y_1) \perp D|X$$

- ► After controlling for X, treatment is independent of potential outcomes: 'No unmeasured confounders'
- ► This is an assumption
 - We cannot directly test it

 Up until now causal estimates have required treatment assignment to be independent of potential outcomes

$$(y_0,y_1)\perp D$$

$$(y_0, y_1) \perp D|X$$

- ► After controlling for X, treatment is independent of potential outcomes: 'No unmeasured confounders'
- ► This is an assumption
 - We cannot directly test it
 - We have to make an argument and provide supporting evidence

► Why does controlling for confounders help provide conditional independence?


- Why does controlling for confounders help provide conditional independence?
- We need to know what problem what bias confounders create:


- Why does controlling for confounders help provide conditional independence?
- We need to know what problem what bias confounders create:
 - ► The problem is of 'fake correlations' D and Y look like they're related, even though treatment does not affect the outcome.
- ► Controlling removes these fake correlations by only comparing D and Y for units with the same value of X


Causal Diagrams (DAGs)













Income


gender





gender





▶ If we omit a confounding variable, we bias our regression estimate:

- ▶ If we omit a confounding variable, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

- ▶ If we omit a confounding variable, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► The 'wrong' regression with a missing confounder:

- ► If we omit a confounding variable, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► The 'wrong' regression with a missing confounder:

$$Y_i = \alpha + \beta D_i + \epsilon_i$$

- ▶ If we omit a confounding variable, we bias our regression estimate:
- ► 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► The 'wrong' regression with a missing confounder:

$$Y_i = \alpha + \beta D_i + \epsilon_i$$

What happens to our coefficient estimate?

- ▶ If we omit a confounding variable, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► The 'wrong' regression with a missing confounder:

$$Y_i = \alpha + \beta D_i + \epsilon_i$$

▶ What happens to our coefficient estimate?

$$X_i = \psi + \delta D_i + \epsilon_i$$

- ► If we omit a confounding variable, we bias our regression estimate:
- 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► The 'wrong' regression with a missing confounder:

$$Y_i = \alpha + \beta D_i + \epsilon_i$$

▶ What happens to our coefficient estimate?

$$X_i = \psi + \delta D_i + \epsilon_i$$

$$Y_i = \alpha + \beta D_i + \gamma (\psi + \delta D_i + \epsilon_i) + \epsilon_i$$

- ► If we omit a confounding variable, we bias our regression estimate:
- ► 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► The 'wrong' regression with a missing confounder:

$$Y_i = \alpha + \beta D_i + \epsilon_i$$

What happens to our coefficient estimate?

$$X_i = \psi + \delta D_i + \epsilon_i$$

$$Y_i = \alpha + \beta D_i + \gamma (\psi + \delta D_i + \epsilon_i) + \epsilon_i$$

$$Y_i = \alpha + D_i(\beta + \gamma \delta) + \gamma(\psi + \epsilon_i) + \epsilon_i$$

- ► If we omit a confounding variable, we bias our regression estimate:
- ► 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► The 'wrong' regression with a missing confounder:

$$Y_i = \alpha + \beta D_i + \epsilon_i$$

What happens to our coefficient estimate?

$$X_i = \psi + \delta D_i + \epsilon_i$$

$$Y_i = \alpha + \beta D_i + \gamma (\psi + \delta D_i + \epsilon_i) + \epsilon_i$$

$$Y_i = \alpha + D_i(\beta + \gamma \delta) + \gamma(\psi + \epsilon_i) + \epsilon_i$$

► So the coefficient we estimate is wrong by this amount:

- ▶ If we omit a confounding variable, we bias our regression estimate:
- ► 'True' regression with all confounders:

$$Y_i = \alpha + \beta D_i + \gamma X_i + \epsilon_i$$

► The 'wrong' regression with a missing confounder:

$$Y_i = \alpha + \beta D_i + \epsilon_i$$

► What happens to our coefficient estimate?

$$X_i = \psi + \delta D_i + \epsilon_i$$

$$Y_i = \alpha + \beta D_i + \gamma (\psi + \delta D_i + \epsilon_i) + \epsilon_i$$

$$Y_i = \alpha + D_i(\beta + \gamma \delta) + \gamma(\psi + \epsilon_i) + \epsilon_i$$

► So the coefficient we estimate is wrong by this amount:

$$\beta_{wrong} = \beta_{true} + \gamma \delta$$

► What does controlling **do**?

- ► What does controlling **do**?
 - ► It means **removing the variation** in the data due to the confounder

- ▶ What does controlling do?
 - ► It means **removing the variation** in the data due to the confounder
 - ► Equivalently, it means separating our data for each value of the confounder: **Subclassification**

- ▶ What does controlling do?
 - It means removing the variation in the data due to the confounder
 - Equivalently, it means separating our data for each value of the confounder: Subclassification
 - ► Then, within each group, the confounder is **constant** and can't affect the relationship between *D* and *Y*.

- ▶ What does controlling do?
 - It means removing the variation in the data due to the confounder
 - Equivalently, it means separating our data for each value of the confounder: Subclassification
 - ► Then, within each group, the confounder is **constant** and can't affect the relationship between *D* and *Y*.
 - We have created balance between the treated and control groups on the confounder

- ▶ We receive a dataset with twenty variables: D, Y and 18 more.
 - ► Which variables should we include as controls?

- ► We receive a dataset with twenty variables: *D*, *Y* and 18 more.
 - ► Which variables should we include as controls?
- ► We usually should **NOT** include all of them

- We receive a dataset with twenty variables: D, Y and 18 more.
 - Which variables should we include as controls?
- ► We usually should **NOT** include all of them
 - Only the variables necessary to stop confounding

- ▶ We receive a dataset with twenty variables: D, Y and 18 more.
 - Which variables should we include as controls?
- We usually should NOT include all of them
 - Only the variables necessary to stop confounding
 - ► Including unnecessary variables can produce bias
 - "Bad controls"/"Post-treatment Bias"

- ▶ We receive a dataset with twenty variables: D, Y and 18 more.
 - Which variables should we include as controls?
- We usually should NOT include all of them
 - Only the variables necessary to stop confounding
 - Including unnecessary variables can produce bias
 - ► "Bad controls"/"Post-treatment Bias"
 - ► We lose power (degrees of freedom) for every control we add

- ▶ We receive a dataset with twenty variables: D, Y and 18 more.
 - ► Which variables should we include as controls?
- We usually should NOT include all of them
 - Only the variables necessary to stop confounding
 - Including unnecessary variables can produce bias
 - ► "Bad controls"/"Post-treatment Bias"
 - ► We lose power (degrees of freedom) for every control we add
 - ► And additional variables reduce overlap (increase model-dependence)

- ▶ We receive a dataset with twenty variables: D, Y and 18 more.
 - ▶ Which variables should we include as controls?
- We usually should NOT include all of them
 - Only the variables necessary to stop confounding
 - Including unnecessary variables can produce bias
 - ► "Bad controls"/"Post-treatment Bias"
 - ► We lose power (degrees of freedom) for every control we add
 - And additional variables reduce overlap (increase model-dependence)

Section 2

Which Variables to Control For

➤ To know which variables to control for, it helps to draw a causal diagram

- ➤ To know which variables to control for, it helps to draw a causal diagram
 - ► A **Directed Acyclical Graph** (DAG)

- ► To know which variables to control for, it helps to draw a causal diagram
 - ► A Directed Acyclical Graph (DAG)
 - Arrows only in one direction

- ► To know which variables to control for, it helps to draw a causal diagram
 - ► A Directed Acyclical Graph (DAG)
 - Arrows only in one direction
 - No circular loops!

Treatment — Outcome

- ► Causation is like **Water**, flowing along the graph
 - We want to focus on one 'flow' of causation from treatment to outcomes

- ► Causation is like **Water**, flowing along the graph
 - We want to focus on one 'flow' of causation from treatment to outcomes
 - Avoiding mixing with the other flows of causation in the network



- ► Three Rules to achieve Conditional Independence:
 - Include as controls enough variables to block all back-door paths from treatment to the outcome

- ► Three Rules to achieve Conditional Independence:
 - Include as controls enough variables to block all back-door paths from treatment to the outcome
 - 2. Exclude any variables that are **post-treatment**

- ► Three Rules to achieve Conditional Independence:
 - Include as controls enough variables to block all back-door paths from treatment to the outcome
 - 2. Exclude any variables that are post-treatment
 - 3. Exclude any variables that are colliders

► To identify back-door paths:

- ► To identify back-door paths:
 - ► Start with an arrow pointing at treatment

- ▶ To identify back-door paths:
 - Start with an arrow pointing at treatment
 - ► Trace the path 'backwards' (the direction of the arrows doesn't matter) until you reach the outcome

- ▶ To identify back-door paths:
 - Start with an arrow pointing at treatment
 - ► Trace the path 'backwards' (the direction of the arrows doesn't matter) until you reach the outcome
 - ► Repeat for every possible path from treatment to outcome


- ▶ To identify back-door paths:
 - Start with an arrow pointing at treatment
 - ► Trace the path 'backwards' (the direction of the arrows doesn't matter) until you reach the outcome
 - Repeat for every possible path from treatment to outcome
- ▶ Block back-door paths by controlling for any variable along the path


- ► To identify back-door paths:
 - Start with an arrow pointing at treatment
 - ► Trace the path 'backwards' (the direction of the arrows doesn't matter) until you reach the outcome
 - ► Repeat for every possible path from treatment to outcome
- ▶ Block back-door paths by controlling for any variable along the path
- ► Identify the **Minimum set of controls** that blocks *All* back-door paths

- ▶ To identify back-door paths:
 - Start with an arrow pointing at treatment
 - ► Trace the path 'backwards' (the direction of the arrows doesn't matter) until you reach the outcome
 - ► Repeat for every possible path from treatment to outcome
- ▶ Block back-door paths by controlling for any variable along the path
- ► Identify the Minimum set of controls that blocks All back-door paths
 - ► This achieves conditional independence of treatment from potential outcomes!


- To identify back-door paths:
 - Start with an arrow pointing at treatment
 - ► Trace the path 'backwards' (the direction of the arrows doesn't matter) until you reach the outcome
 - ► Repeat for every possible path from treatment to outcome
- ▶ Block back-door paths by controlling for any variable along the path
- ► Identify the Minimum set of controls that blocks All back-door paths
 - ► This achieves conditional independence of treatment from potential outcomes!
 - ► Include these as control variables in our regression







1. Back-door Paths




► Including **post-treatment** variables will introduce bias

- Including post-treatment variables will introduce bias
 - Because variables measured 'after' treatment can also be affected by treatment

- Including post-treatment variables will introduce bias
 - Because variables measured 'after' treatment can also be affected by treatment
 - They're not confounders, but mechanisms/mediating variables

- Including post-treatment variables will introduce bias
 - Because variables measured 'after' treatment can also be affected by treatment
 - ► They're not confounders, but mechanisms/mediating variables
 - ► Controlling for them changes the definition of the causal effect we are estimating




► Colliders are variables on back-door paths which have arrows pointing both into them *and* out of them

- ► Colliders are variables on back-door paths which have arrows pointing both into them *and* out of them
- ► The water 'collides' in both directions so the source variables are not correlated, and produce no bias

- Colliders are variables on back-door paths which have arrows pointing both into them and out of them
- ► The water 'collides' in both directions so the source variables are not correlated, and produce no bias
- ▶ But if we do accidentally 'control' for a collider we introduce a bias in the relationship between D and Y

- Colliders are variables on back-door paths which have arrows pointing both into them and out of them
- ► The water 'collides' in both directions so the source variables are not correlated, and produce no bias
- ► But if we do accidentally 'control' for a collider we *introduce* a bias in the relationship between *D* and *Y*
- So we must avoid controlling for colliders

- ► Colliders are variables on back-door paths which have arrows pointing both into them *and* out of them
- ► The water 'collides' in both directions so the source variables are not correlated, and produce no bias
- ► But if we do accidentally 'control' for a collider we *introduce* a bias in the relationship between *D* and *Y*
- So we must avoid controlling for colliders
- ► Hard!







Figure 3.4: Simulation of conditional dependence within values of a collider variable.




1. List all of the **back-door paths** from *D* to *Y*

- 1. List all of the **back-door paths** from D to Y
- 2. Identify any post-treatment variables: Do NOT include as controls

- 1. List all of the **back-door paths** from *D* to *Y*
- 2. Identify any **post-treatment** variables: Do NOT include as controls
- Identify any back-door paths with collider variables: Mark these as already blocked

- 1. List all of the **back-door paths** from *D* to *Y*
- 2. Identify any **post-treatment** variables: Do NOT include as controls
- 3. Identify any back-door paths with **collider** variables: Mark these as already blocked
- 4. Find a minimum set of variables that blocks all remaining back-door paths

- 1. List all of the **back-door paths** from *D* to *Y*
- 2. Identify any **post-treatment** variables: Do NOT include as controls
- Identify any back-door paths with collider variables: Mark these as already blocked
- 4. Find a minimum set of variables that blocks all remaining back-door paths
- 5. Double-check your minimum set of control variables does not contain any post-treatment or collider variables



- ► Three Rules to achieve Conditional Independence:
 - 1. Include as controls enough variables to **block all back-door paths** from treatment to the outcome

- ► Three Rules to achieve Conditional Independence:
 - Include as controls enough variables to block all back-door paths from treatment to the outcome
 - ► In practice, variables which theory and past studies identify as potential confounders

- ► Three Rules to achieve Conditional Independence:
 - Include as controls enough variables to block all back-door paths from treatment to the outcome
 - In practice, variables which theory and past studies identify as potential confounders
 - 2. Exclude any variables that are **post-treatment**

- ► Three Rules to achieve Conditional Independence:
 - Include as controls enough variables to block all back-door paths from treatment to the outcome
 - In practice, variables which theory and past studies identify as potential confounders
 - 2. Exclude any variables that are **post-treatment**
 - ► In practice, know when your variables were measured

- ► Three Rules to achieve Conditional Independence:
 - Include as controls enough variables to block all back-door paths from treatment to the outcome
 - In practice, variables which theory and past studies identify as potential confounders
 - 2. Exclude any variables that are **post-treatment**
 - ► In practice, know when your variables were measured
 - 3. Exclude any variables that are **colliders**

- ► Three Rules to achieve Conditional Independence:
 - Include as controls enough variables to block all back-door paths from treatment to the outcome
 - In practice, variables which theory and past studies identify as potential confounders
 - 2. Exclude any variables that are **post-treatment**
 - ► In practice, know when your variables were measured
 - 3. Exclude any variables that are colliders
 - ► In practice, don't include unnecessary controls