ESTIMATING NEUTRINO FLUX FROM LOW MASS STELLAR MODEL USING MACHINE LEARNING TECHNIQUE

IRFAN MUQRI BIN ABD RAHMAN

FACULTY OF SCIENCE UNIVERSITI MALAYA KUALA LUMPUR

2024

ESTIMATING NEUTRINO FLUX FROM LOW MASS STELLAR MODEL USING MACHINE LEARNING TECHNIQUE

IRFAN MUQRI BIN ABD RAHMAN

FINAL YEAR PROJECT REPORT SUBMITTED IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE OF BACHELOR OF SCIENCE IN PHYSICS

DEPARTMENT OF PHYSICS FACULTY OF SCIENCE UNIVERSITI MALAYA KUALA LUMPUR

2024

UNIVERSITI MALAYA ORIGINAL LITERARY WORK DECLARATION

Name of Candidate: IRFAN MUQRI BIN ABD RAHMAN (021214-01-0451)

Matric No: U2103688

Name of Degree: BACHELOR OF SCIENCE IN PHYSICS

Title of Project Paper ("this Work"):

ESTIMATING NEUTRINO FLUX FROM LOW MASS STELLAR MODEL USING

MACHINE LEARNING TECHNIQUE

Field of Study: Experimental Physics

I do solemnly and sincerely declare that:

- (1) I am the sole author of this Work;
- (2) This Work is original;
- (3) Any use of any work in which copyright exists was done by way of fair dealing and for permitted purposes and any excerpt or extract from, or reference to or reproduction of any copyright work has been disclosed expressly and sufficiently and the title of the Work and its authorship have been acknowledged in this Work;
- (4) I do not have any actual knowledge nor do I ought reasonably to know that the making of this work constitutes an infringement of any copyright work;
- (5) I hereby assign all and every rights in the copyright to this Work to the University of Malaya ("UM"), who henceforth shall be owner of the copyright in this Work and that any reproduction or use in any form or by any means whatsoever is prohibited without the written consent of UM having been first had and obtained;
- (6) I am fully aware that if in the course of making this Work I have infringed any copyright whether intentionally or otherwise, I may be subject to legal action or any other action as may be determined by UM.

Candidate's Signature Date: January 6, 2025

Subscribed and solemnly declared before,

Witness's Signature Date: January 6, 2025

Name:

Designation:

ESTIMATING NEUTRINO FLUX FROM LOW MASS STELLAR MODEL USING MACHINE LEARNING TECHNIQUE

ABSTRACT

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

MENGANGGARKAN FLUKS NEUTRINO DARIPADA MODEL BINTANG JISIM RENDAH MENGGUNAKAN TEKNIK PEMBELAJARAN MESIN

ABSTRAK

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

ACKNOWLEDGEMENTS

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

TABLE OF CONTENTS

Abstrac	t
Abstrak	iv
Acknow	rledgements
Table of	Contentsvi
List of I	Figures
List of 7	Tables ix
List of S	Symbols and Abbreviations
List of A	Appendices
СНАРТ	TER 1: INTRODUCTION
1.1	Problem Statement
1.2	Research Questions
1.3	Research Objectives
1.4	Relevance of Research
СНАРТ	TER 2: LITERATURE REVIEW 3
2.1	Stellar Model
2.2	Neutrino
2.3	Neutrino Sources
2.4	Solar Neutrino
2.5	Theoretical vs Experimental
2.6	Current Works
2.7	Machine Learning
СНАРТ	TER 3: METHODOLOGY 7
3.1	Data
3.2	Machine Learning

CHAPTER 4: RESULTS	8
CHAPTER 5: DISCUSSION	9
CHAPTER 6: CONCLUSION	10
Deferences	11

LIST OF FIGURES

Figure 2.1: Neutrino sources	4
------------------------------	---

LIST OF TABLES

Table 2.1: Most common reaction that produce ν	4
--	---

LIST OF SYMBOLS AND ABBREVIATIONS

EA : Example Abbreviation

LIST OF APPENDICES

																																				_
Appendix A: Example																																			- 1	3
appendix is Example	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	1.	_

CHAPTER 1: INTRODUCTION

Neutrino is one of the elementary particles in the fermion group and were theoretically invented in 1930 by Pauli to preserve energy–momentum conservation. It is a neutral particle which only has a very small mass and is only affected by gravitation and weak interaction. Neutrinos are known to be the most abundant particles in the universe after photon with a density of approximately 330 cm⁻³ pan universe (Athar & Singh, 2020). All these neutrinos come from a lot of sources, they are produced inside the sun, the earth, the entire atmosphere, during the birth, collision, and death of stars and huge flux of neutrinos is emitted during supernovae explosions. In stars like our sun the amount of neutrinos produce is very abundant as it is part of the fission process through beta decay and electron capture, for example in the proton-proton chain reaction where 2 ¹H fuses into one ²H converting one of the protons into neutron (Iliadis, 2007). And due to it having small collision cross-section most of these neutrinos go out of the sun without being blocked going out to every direction, a lot of them pass through us. Neutrino flux is the measure of the number of neutrinos flowing through, on earth we use detectors such as the Super Kamiokande.

Machine learning is a technique that improves system performance by learning from experience via computational means (Zhou, 2021). In physics, especially through recent decades, machine learning has taken its roots, especially in statistical analysis with a large amount of data. For example, it has been used in processing satellite data in atmospheric physics, in weather forecasts, predicting the behavior of systems of many particles, discovering functional materials and generating new organic molecules (Rodrigues, 2023).

We do have data for solar neutrinos from the sun which is a low mass stellar model from various of detectors from earth. However, for stars far away we must estimate the neutrino flux since there is no method to directly detect the neutrino. There are some theoretical calculations that can be done to estimate the neutrino flux but, in this research, we plan to use machine learning to do the estimation.

1.1 Problem Statement

Neutrinos are present in huge quantities owing to various astrophysical phenomena such as those in a stellar medium like the Sun. Although solar neutrino flux is able to be measured on the Earth by using a detector like Super Kamiokande, neutrinos emitted from distant stars and the physics of their detection still presents a problem because of the difficulty in interaction and the detection of such distant and sparse interactions by current technology. There is no proper means of detecting neutrino radiation from stellar sources at the present time, though theorists continue to make predictions based on elaborate computations.

To address these issues, machine learning approaches, which are particularly efficient when it comes to complex data processing tasks and prediction problems located in various physics, would be one of the available options. This study aims to solve the problem by estimating the neutrino flux from a low mass stellar model through the use of a machine learning approach. Here we will try to construct and train a supervised machine learning model aimed at predicting neutrino flux from stars which are not accessible by direct observation, utilizing available solar neutrino flux data and theoretical models.

1.2 Research Questions

The research aims to answer the following questions:

- 1. What is the neutrino flux production beyond the Standard Solar Model in low mass stars?
- 2. Is machine learning able to give the best estimate in the current solar neutrino productions and what is the limit when extended to beyond the Standard Solar Model?

1.3 Research Objectives

The research aims to achieve the following objectives:

- 1. To train using machine learning techniques to estimate neutrino flux that can mimic real flux measured
- 2. To test the accuracy of linear regression from the machine learning in estimating neutrino flux

1.4 Relevance of Research

- Provide new understanding in the neutrino flux or production beyond Standard Solar Model
- 2. To test the robustness of the linear regression in handling the neutrino flux calculation from low mass stars grid.

CHAPTER 2: LITERATURE REVIEW

2.1 Stellar Model

A stellar model is a mathematical model that describes a star structure and evolution. Stellar structure models describe the internal structure of a star in detail and make predictions about the luminosity, the color and the future evolution of the star. Different classes and ages of stars have different internal structures, reflecting their elemental makeup and energy transport mechanisms. Stellar evolution is the process by which a star changes over the course of its lifetime. A stellar evolutionary model is a mathematical model that can be used to compute the evolutionary phases of a star from its formation until it becomes a remnant.

Example of a stellar model is the Standard Solar Model (SSM) which is a mathematical model that describes the Sun as a spherical ball of gas, and it's used to test the validity of stellar evolution theory. The SSM is used to test the validity of stellar evolution theory. In fact, the only way to determine the two free parameters of the stellar evolution model, the helium abundance and the mixing length parameter (used to model convection in the Sun), are to adjust the SSM to fit the observed Sun.

2.2 Neutrino

An elementary particle first proposed by Pauli in 1931 and named by Fermi; the neutrino was to account for the missing energy released in β decay. Due to conservation of electric charge, it must be neutral, and conservation of angular momentum requires it to have a spin of $\frac{1}{2}$. There are two types of neutrinos emitted in β decay neutrino, e and antineutrino, e and they are emitted in β + and β - decays respectively. The neutrino has a very small cross section, σ of 1.2×10^{-43} cm² (Krane, 1991). Due to this, neutrinos can escape from the sun in about 2 seconds which is very fast compared to the 100,000 years for a photon (Rosso et al., 2018). The first neutrino was detected experimentally in the 1950s by Reines and Cowan using a nuclear reactor as the source where β - decays occur producing e.

2.3 Neutrino Sources

There are multiple sources of neutrino both on earth and extraterrestrial, for example the Cosmic Neutrino Background (CNB), Supernovae, Solar and Thermal neutrinos from the sun, Geoneutrinos and many more as shown in Figure 2.1 (Athar & Singh, 2020).




Figure 17.1 Different sources of neutrinos

Figure 2.1: Neutrino sources.

Most of them produce neutrinos through β decays and electron(positron) capture like shown in Table 2.1 (Rosso et al., 2018):

Table 2.1: Most common reaction that produce ν .

Name	Reaction
β^+ decays	$n \rightarrow p + e^- + \nu_e$
β^- decays	$p \to n + e^+ + \bar{\nu}_e$
e^- capture	$e^- + p \rightarrow n + \nu_e$
e^+ capture	$e^+ + n \rightarrow p + \bar{\nu}_e$

that produce ν .

And unlike most other particles which are charged neutrinos can be assumed to come directly from their source as they are not deflected by magnetic field and unlike photons, neutrinos weakly interact making them able to pass through matter easily, the interaction length of a 1-TeV neutrino is about 2.5 million kilometers of water, or 250 kt cm⁻², whereas high-energy photons are blocked by a few hundred g cm⁻² (Barenboim & Quigg, 2003).

Graph 1 (Vitagliano et al., 2020) shows the fluxes from different neutrino sources. The range of energy for different sources is quite huge, spanning over 24 orders of magnitude.

2.4 Solar Neutrino

The sun through the thermonuclear fusion reaction produces a huge amount of e with energy the order of 1MeV. Since neutrino interaction with matter is weak, almost all the neutrino produced in the core goes out from the sun. The earth receives about

 6×10^{10} cm⁻² s flux of solar neutrinos. For the sun 2.3% of its nuclear energy production is in the form of MeV range electron neutrinos (Giunti & Kim, 2007). This comes from the effective nuclear reaction

$$4p + 2e \rightarrow 2^4 He + 2v_e + 26.73 MeV$$
 (2.1)

Which comes from several reaction chains and cycles (Vitagliano et al., 2020).

Table 2 (Vitagliano et al., 2020) shows the different reactions that produce neutrinos that can be detected on earth that the sun produces in its thermonuclear process. While table 3 (Rosso et al., 2018) shows the expectation of the solar neutrino flux from each branch coming to earth. Referring to table Graph 1, we can see the shape of the flux in a solid blue line labelled "Solar (nuclear)".

2.5 Theoretical vs Experimental

2.5.1 Theoretical

For the theoretical side of stellar physics, most of them are done in doing stellar modelling to predict structures and evolution of certain mass stars. Example of theoretical work done in neutrino is Stellar Neutrino Emission across the Mass–Metallicity Plane done by (Farag et al., 2020) where they explore neutrino emission from nonrotating, single-star models across six initial metallicities and 70 initial masses from the zero-age main sequence to the final fate. The data produced in the model are used in this research in order to train the model.

Since the time and computational power needed to run even one model is quite arduous, by using machine learning, the gap between the modelled grid can be predicted using machine learning.

2.5.2 Experimental

Due to neutrinos having an almost negligible mass and interact weakly with matter, the experiments that are required to detect them are usually done on a large scale in the range of km3 which can be setback by technological challenges (Addazi et al., 2022). Notable experiments include Super Kamiokande in Japan, the Sudbury Neutrino Observatory (SNO) in Canada, and the Jiangmen Underground Neutrino Observatory (JUNO) in China. Data from these experiments can validate the prediction obtained from a machine learning model.

2.6 Current Works

2.7 Machine Learning

Machine learning has been on the rise rapidly and can potentially revolutionize how we do sciences, including physics. According to Rodrigues (2023) even though machine learning can help solve fundamental problems in physics, many physicists still do not recognise the importance of these techniques. In physics, machine learning is often used in settings with complex problems and lots of data like in the large hadron collider (LHC) experiments where the data can reach up to ten of thousands of petabytes per year.

Figure 2: Supervise machine learning pipeline In Figure 2 (Rodrigues, 2023), usually in supervised learning 80% of data is used to train the model while 20% is for testing the model. In research done by (Niu et al., 2019), by using Bayesian neural network (BNN) which is a machine learning technique, they manage to get results that well reproduce the experimental data with very high accuracy and provide reasonable uncertainty in half-life prediction as shown in Figure 3 compared to prediction done using calculation.

Figure 3: β decay half-life of Ni isotopes Where the experimental values in NUBASE2016 are denoted by spheres. The half-life predictions with calculation are shown by the dotted line, and their counterparts improved by BNN-I2 and BNN-I4 approaches, and their uncertainties are shown by vertical line hatched region and green hatched region, respectively. The mean predicted half-lives of BNN-I4 approach are marked by the dashed line.

CHAPTER 3: METHODOLOGY

3.1 Data


$$Flux = \frac{L}{4\pi r^2}$$
 (3.2)

3.2 Machine Learning

$$\begin{bmatrix} m_1 \\ m_2 \\ m_3 \end{bmatrix} J = \begin{bmatrix} L_{11} & A_{11} & L_{12} & A_{12} & L_{13} & A_{13} \\ L_{21} & A_{21} & L_{22} & A_{22} & L_{23} & A_{23} \\ L_{31} & A_{31} & L_{32} & A_{32} & L_{33} & A_{33} \end{bmatrix}$$
(3.3)

where m is mass, L is luminosity, A is star age and J is

Then using SciKit Learn



CHAPTER 4: RESULTS

CHAPTER 5: DISCUSSION

CHAPTER 6: CONCLUSION

REFERENCES

- Addazi, A., Alvarez-Muniz, J., Alves Batista, R., Amelino-Camelia, G., Antonelli, V., Arzano, M., ... Zornoza, J. (2022). Quantum gravity phenomenology at the dawn of the multi-messenger era—a review. *Progress in Particle and Nuclear Physics*, *125*, 103948. Retrieved from https://www.sciencedirect.com/science/article/pii/S0146641022000096 doi: https://doi.org/10.1016/j.ppnp.2022.103948
- Athar, M. S., & Singh, S. K. (2020). *The Physics of Neutrino Interactions*. Cambridge University Press.
- Barenboim, G., & Quigg, C. (2003, 4). Neutrino observatories can characterize cosmic sources and neutrino properties. *Physical review. D. Particles, fields, gravitation, and cosmology/Physical review. D. Particles and fields*, 67(7). Retrieved from https://doi.org/10.1103/physrevd.67.073024 doi: 10.1103/physrevd.67.073024
- Farag, E., Timmes, F. X., Taylor, M., Patton, K. M., & Farmer, R. (2020, apr). On stellar evolution in a neutrino hertzsprung–russell diagram. *The Astrophysical Journal*, 893(2), 133. Retrieved from https://dx.doi.org/10.3847/1538-4357/ab7f2c doi: 10.3847/1538-4357/ab7f2c
- Giunti, C., & Kim, C. W. (2007). Fundamentals of Neutrino Physics and Astrophysics. Oxford University Press.
- Iliadis, C. (2007). Nuclear Physics of Stars. John Wiley & Sons.
- Krane, K. S. (1991). Introductory Nuclear Physics. John Wiley & Sons.
- Niu, Z. M., Liang, H. Z., Sun, B. H., Long, W. H., & Niu, Y. F. (2019, 6). Predictions of nuclear β-decay half-lives with machine learning and their impact on r-process nucleosynthesis. *Physical review. C*, 99(6). Retrieved from https://doi.org/10.1103/physrevc.99.064307 doi: 10.1103/physrevc.99.064307
- Rodrigues, F. A. (2023, 10). Machine learning in physics: A short guide. *EPL (Europhysics Letters)*, 144(2), 22001. Retrieved from https://doi.org/10.1209/0295-5075/ad0575 doi: 10.1209/0295-5075/ad0575
- Rosso, A. G., Mascaretti, C., Palladino, A., & Vissani, F. (2018, 7). Introduction to neutrino astronomy*. *The European Physical Journal Plus*, *133*(7). Retrieved from https://doi.org/10.1140/epjp/i2018-12143-6 doi: 10.1140/epjp/i2018-12143-6
- Vitagliano, E., Tamborra, I., & Raffelt, G. (2020, 12). Grand unified neutrino spectrum at Earth: Sources and spectral components. *Reviews of Modern Physics*, 92(4). Retrieved from https://doi.org/10.1103/revmodphys.92.045006 doi: 10.1103/revmodphys.92.045006
- Zhou, Z.-H. (2021). *Machine learning*. Retrieved from https://doi.org/10.1007/

978-981-15-1967-3 doi: 10.1007/978-981-15-1967-3

APPENDIX A: EXAMPLE