

Project-plan

Background

Trauma is a wide term including various physical injuries to the human body. It is one of the leading causes of mortality and morbidity in the world, representing about 9 % of annual global deaths. Over the last decade almost 50 million people worldwide have died from trauma.(1)

Not only does trauma represent a large share of the global mortality rate, but studies have also shown a significant difference in outcome depending on where patients are treated. It has for example been shown that trauma patients in Sweden who were treated at a trauma center rather than a non-trauma center have a 41 % lower 30-day adjusted mortality rate. (2)

To further stress the need for more knowledge and research about Trauma care, some studies indicate that the number of trauma-related deaths that potentially could have been prevented is as high as 20 to over 50 %. (3–5) Preventable deaths varies not only between trauma centers and non-trauma centers. There is a significant difference between high income counties and low and middle income countries (LMICs). It is estimated that 2 million lives could be saved annually if LMICs and high income countries had equal fatality numbers for severe injuries.(6)

Knowledge gap

Today It is poorly understood whether different subgroups have greater opportunities for improvement (OFI) than others. The most common cause of death after injury is hemorrhage. Among these patients, problem in decision making, technical skills and making decisions between surgery and radiology were some of the areas where OFIs were found.(7)

Since trauma patients is a very heterogeneous group, it is important to have a sufficient understanding of OFIs for different trauma subgroups.(8)

Aim

In this study we aim to assess the frequency of opportunity for improvement in the following important clinical subgroups. Men and women, blunt and penetrating injuries, minor and major trauma ,and across body regions injured.

Trauma Registries

In sweden there is a national trauma register (SweTrau) holdng records of trauma cases from 48 out of 49 hospitals who recieves major trauma. To meet the inclusions criterias for SweTrau you must be over 15 years of age, been exposed to a traumatic event serious enough to activate the Trauma code in the hospital or have a NISS over 15 without trauma code activation. (9)

The exclusion criterias for SweTrau are as following: Patients where the only traumatic injury is a chronic subdural hematoma and patients where a trauma alarm is triggered without an underlying traumatic event.(9)

The KUH trauma register is a part of the Swedish Trauma Register, with the same inclusion and exclusion criteria. (källa jonatan, VAR FINNS INFO OM KAROLINSKAS REGISTER?)

The Trauma Quality Database (TCQD) holds all all severe trauma cases treated at KUH Solna. VAR FINNS MER INFO OM TCQD?

Advanced Trauma Life Support and Primary Trauma Care

Several different systems are being used in Trauma care. Such as Advanced Trauma Life Support (ATLS) and Primary Trauma Care (PTC), where ATLS is the more established system. The purpose of these systems is to secure a time-efficient, standardized and structured way of treating trauma patients.(10,11)

ATLS is practiced in over 80 countries and 1 million doctors have gone thru this training.(10)

PTC is also used in over 80 countries, however more frequently in low and middle-income countries. One reason for this could be that the PTC program is free while ATLS is not.(11)

Tycker ni att jag kan beskriva de olika systemen mer i detalj här för utfyllnad?

Oppotrunity for improvement

Whether there is opportunity for improvement for a specific case or not is decided by a group of experts during a conference where every trauma case is discussed. OFI is defined as when the trauma care for a patient does not match the best practice guidelines in at least one aspect.

At KUH all trauma patient end up in a data base. Audit filters then point out certain trauma cases where things might have differed from the golden standard treatment based on different criterias. Such criterias are GCS 8 or less where the patient was not intubated, time to CT, time to Surgery and so on. Then a manual selection is done by a nurse, where some cases are removed from the group of potential OFI cases because obvious reasons for the treatment can be found. The patients who are then left are discussed at a conference where doctors from several specialties participate. At this conference every case is gone through thoroughly. Then those patients where OFI is found are marked with "YES" in the OFI column in the KUH Trauma register and those patient where no OFI is found are mared with "NO."

(BEskrivet utifrån vad Jonatan berättade, finns det någon källa man kan hänvisa till. Tex det dokument som jonatan visade när vi träffades.)

New Injury Severity Score

The NISS is an anatomic severity scale developed to assess multiple-injured patients. The NISS score a patient get is based on the Abbreviated Injury Scale (AIS) points from the injured body regions. The higher NISS score the more serious injury the patients suffers from.(12)

NISS takes injuries in 6 different body regions into account. Those are head/neck, face, chest, abdomen, extremiteis and external. The NISS is the sum of the squares of the highest AIS codes in these regions, meaning that all three injuries can be in the same region or in separate regions. AIS can vary from 1-5, and the NISS from 1-75.(12)

Initial Trauma care

Trauma patients are divided into priority one and two by the paramedics using certain criteria, such as trauma mechanism, GCS points and blood pressure. To Karolinska Solna only those who are classified as a priority one by the pre hospital professionals are admitted. (13)

A Trauma priority one is considered directly life threatening. Therefore when arriving to Karolinska Solna every one of these patients are taken care of by a full trauma team. This team consists of a trauma leader wo is a general surgeon or a resident in general surgery and an anesthetist with a nurse specialized in anesthesiology. The team also has an rthopedic surgeon, radiologist, radiology nurse, emergency medicine nurse, surgical nurse and assistant nurses.(13)

Methods

Study design

This is a registry based cohort study that uses data from two different Swedish trauma registers. The first one is the Trauma registry at the Karolinska University Hospital in Solna, which includes about 21000 patients between the years 2012 and 2021. The second register is the Trauma quality database (TCQD) which is a subset of the trauma registry and includes about 2200 patients selected for review between 2014 and 2021.

By linking these databases together the opportunity for improvement in the trauma subgroups mentioned in the introduction will be assessed.

Setting

The Karolinska University Hospital in Solna, Stockholm, is the leading trauma center in Sweden, and ranked as one of the best hospitals in the world. It is also considered as a level 1 trauma center.⁽¹⁴⁾ The Trauma department at KUH Solna handles about 1800 trauma cases every year, out of which about 200 are children. KUH is the primary trauma center for a population of 2,8 million people.⁽¹⁵⁾

Participants

All patients from KUH trauma register and from the TCQD.

Variables and data sources/measurements

Ska jag här bara beskriva igen att det mått som vi använder är OFI?

Bias

To prevent research bias all data will be scrambled while used to write the analysis model.

Study size

Quantitative variables

Statistical methods

Ethical considerations

All patients have given consent before added to Swetrau and TCQD. As this is a registry based study no interventions were made that could effect the patients. The only risk with this type of studies is that personal information could leak. In these registers however, all personal data was scrambled to minimize that risk.

Ethical permission has been granted for this project. Dnr number for this permission is 2021-02541 and 2021-0253.

DET STÅR ATT ALLA GETT MEDGIVANDE FÖR ATT LÄGGAS TILL I SWETRAU PÅ DERAS HEMSIDA. MEN HUR FUNKAR DET MED DE SOM DOG INNAN SAMTYCKE GAVS? Jonatan pratade om att man antog samtycke för de patienterna. Men hittar ingen info om detta.

En skriftlig redogörelse av projektets utveckling

All data är redan insamlad från start och finns tillgängliga i de register som beskrivs i projektredogörelsen. Datan kommer delas in efter olika trauma-subgrupper. De variabler som kommer användas är "Ja" där det finns "Opportunity for improvement" samt "nej" där det inte finns "opportunity for improvement."

Hittills har jag - Läst in mig på ämnet - Skrivit Introduktion - Påbörjat materiella och metod - Börjat sätta mig in i R som är det program som kommer användas för att sammanställa data.

- Hur data planeras presenteras (t ex tabeller och hur de kan struktureras, diagramtyper, vävnadssnitt, intervjuer etc)

Datan kommer presenteras i en tabell där de aktuella subgrupperna kommer presenteras samt huruvida det finns OFI för dessa grupper. Det kommer även redogöras för i vilken utsträckning varje subgrupp har OFI.

- Hur data planeras analyseras; vilka statistiska metoder för jämförelser mellan en eller flera grupper, andra matematiska metoder, kvalitativ analys etc → Detta skulle jag behöva diskutera med er innan jag kan svara.

Ingen reservplan behövs då all data finns insamlad och risken för att projektet inte går att genomföra somplanerat bedöms minimal.

References

1. GBD 2017 Causes of Death Collaborators. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980-2017: A systematic analysis for the global burden of disease study 2017. *Lancet*. 2018 Nov;392(10159):1736–88.
2. Candefjord S, Asker L, Caragounis E-C. Mortality of trauma patients treated at trauma centers compared to non-trauma centers in Sweden: A retrospective study. *Eur J Trauma Emerg Surg*. 2022 Feb;48(1):525–36.
3. Drake SA, Holcomb JB, Yang Y, Thetford C, Myers L, Brock M, et al. Establishing a regional trauma preventable/potentially preventable death rate. *Ann Surg*. 2020 Feb;271(2):375–82.
4. Ray JJ, Meizoso JP, Satahoo SS, Davis JS, Van Haren RM, Dermer H, et al. Potentially preventable prehospital deaths from motor vehicle collisions. *Traffic Inj Prev*. 2016 Oct;17(7):676–80.
5. Ghorbani P, Strømmer L. Analysis of preventable deaths and errors in trauma care in a Scandinavian trauma level-I centre. *Acta Anaesthesiol Scand*. 2018 Sep;62(8):1146–53.
6. Mock C, Joshipura M, Arreola-Risa C, Quansah R. An estimate of the number of lives that could be saved through improvements in trauma care globally. *World J Surg*. 2012 May;36(5):959–63.
7. O'Reilly D, Mahendran K, West A, Shirley P, Walsh M, Tai N. Opportunities for improvement in the management of patients who die from haemorrhage after trauma. *British Journal of Surgery* [Internet]. 2013 Mar;100(6):749–55. Available from: <http://dx.doi.org/10.1002/bjs.9096>
8. Coats T. Blunt major trauma: Now two different diseases? *Br J Hosp Med (Lond)*. 2020 Apr;81(4):1–3.
9. <https://rcsyd.se/swetrau/om-swetrau/about-swetrau-in-english/swetrau-the-swedish-trauma-registry>.
10. <https://www.facs.org/quality-programs/trauma/education/advanced-trauma-life-support/about/>.
11. Kadhum M, Sinclair P, Lavy C. Are primary trauma care (PTC) courses beneficial in low- and middle-income countries - a systematic review. *Injury*. 2020 Feb;51(2):136–41.
12. Li H, Ma Y-F. New injury severity score (NISS) outperforms injury severity score (ISS) in the evaluation of severe blunt trauma patients. *Chin J Traumatol*. 2021 Sep;24(5):261–5.
13. Granström A, Wihlke G, Brattström O, Ostlund A. [Activation of the trauma team is related to injury severity. Triage stringency can yield optimal use of resources]. *Lakartidningen*. 2012;109(4):154–7.
14. <https://news.cision.com/karolinska-universitetssjukhuset/r/karolinska-university-hospital-is-the-eighth-best-hospital-in-the-world,c3517653>.
15. <https://www.karolinska.se/for-varldgivarer/tema-akut-och-reparativ-medicin/trauma-akutkirurgi-och-ortopedi/traumacentrum/>.