Locality emerges from hidden zeros in $\mathrm{tr}(\phi^3)$: a by-hand proof at four, five, and six points

Jony

April 18, 2026

Abstract

We prove that the cyclic hidden one-zeros of color-ordered $\mathrm{tr}(\phi^3)$ tree amplitudes force every non-local coefficient of the general rational ansatz to vanish, at $n = 4, 5, 6$ external legs. The proof is entirely by hand: we use only the kinematic mesh identity, geometric series Laurent expansions, and the fact that rational functions in independent variables vanish iff each coefficient on a basis of monomials vanishes. The mechanism is geometric: each cyclic zero kills exactly the non-local terms whose chord factors involve the "enclosed vertex" of a designated short diagonal (the *special variable*). Every non-local coefficient is killed by at least one zero — and as it happens, by exactly two.

Contents

1 Setup **2**
- 1.1 Chords, the ansatz, and locality . . . 2
- 1.2 The kinematic mesh identity . . . 2
- 1.3 The hidden one-zeros . . . 2
- 1.4 The geometric heart: enclosed vertices . . . 3

2 Four points: trivial warm-up **3**

3 Five points: complete proof **3**
- 3.1 The pentagon's chords and crossings . . . 3
- 3.2 The 15-coefficient ansatz . . . 4
- 3.3 The cyclic hidden one-zeros, derived by hand . . . 4
- 3.4 The pattern: every zero is a rotation of the previous . . . 5
- 3.5 The kill technique, in full detail at $\mathcal{Z}_1$ . . . 5
- 3.6 Cycling: kills from $\mathcal{Z}_2, \ldots, \mathcal{Z}_5$ . . . 8
- 3.7 Union: every non-local coefficient is killed . . . 9
- 3.8 The geometric pattern revealed . . . 9

4 Six points: the hexagon case **9**
- 4.1 Setup at $n = 6$ . . . 10
- 4.2 The six zeros, derived by hand . . . 10
- 4.3 The kill technique at $\mathcal{Z}_1$: three layers . . . 11
- 4.4 Cycling: union over six zones is 151 . . . 13

5 Looking ahead: the structural pattern **14**

1 Setup

1.1 Chords, the ansatz, and locality

Label the external legs cyclically by $1, \ldots, n$. The planar Mandelstam variables are

$$X_{ij} \;=\; (p_i + p_{i+1} + \cdots + p_{j-1})^2, \qquad 1 \le i < j \le n,\; j - i \ge 2,\; (i,j) \ne (1,n).$$

Geometrically, X_{ij} is the chord from vertex i to vertex j of the convex n-gon. There are $n(n-3)/2$ such chords ($2, 5, 9$ for $n = 4, 5, 6$).

A *triangulation* of the n-gon is a maximal set of non-crossing chords; it has $n-3$ chords. The tree amplitude is the sum over triangulations of the product of chord propagators,

$$A_n \;=\; \sum_{T \text{ triangulation}} \prod_{(i,j)\in T} \frac{1}{X_{ij}}.$$

The number of triangulations is the Catalan number C_{n-2} ($2, 5, 14$ for $n = 4, 5, 6$).

We consider the most general rational function B with the correct dimension ($-2(n-3)$) and only planar simple poles:

$$B \;=\; \sum_{\text{multisets } (a_1,\ldots,a_{n-3})} \frac{a_{a_1\ldots a_{n-3}}}{X_{a_1}\cdots X_{a_{n-3}}}.$$

The sum is over multisets (unordered with repetition) of size $n-3$ from the set of chord indices. There are $\binom{d+n-4}{n-3}$ such multisets, where $d = n(n-3)/2$ is the number of chords. We get $2, 15, 165$ multisets for $n = 4, 5, 6$.

A multiset is *local* if its chords form a triangulation; the remaining multisets are *non-local* (either repeated chords or two chords that cross). Locality is the statement that all non-local coefficients vanish.

1.2 The kinematic mesh identity

The non-planar two-particle invariants $c_{a,b} = 2p_a \cdot p_b$ ($a \ne b$, non-adjacent) are not independent of the planar variables. They satisfy

$$\boxed{c_{a,b} \;=\; X_{ab+1} + X_{a+1b} - X_{ab} - X_{a+1b+1}}$$

(adjacent chords like X_{aa+1} are zero by masslessness $p_a^2 = 0$). Setting $c_{a,b} = 0$ thus imposes a linear relation on the planar chords.

1.3 The hidden one-zeros

For each leg $r \in \{1, \ldots, n\}$, the *hidden one-zero* $\mathcal{Z}_r$ is defined by setting all c-invariants involving leg r to zero:

$$\mathcal{Z}_r : \qquad c_{r,r+2} = c_{r,r+3} = \cdots = c_{r,r+n-2} = 0 \pmod{n}.$$

There are $n-3$ such constraints. The starting fact (proved elsewhere; here taken as given) is that the tree amplitude A_n vanishes on each $\mathcal{Z}_r$. Our problem is the *converse* for the general ansatz: assuming $B|_{\mathcal{Z}_r} = 0$ for all r, show that all non-local a-coefficients vanish.

1.4 The geometric heart: enclosed vertices

Of the two arcs into which a chord X_{ij} divides the polygon's boundary, the *enclosed arc* is the shorter one. The *enclosed vertex set* is

$$E(X_{ij}) \;=\; \{\text{vertices strictly between } i \text{ and } j \text{ on the shorter arc}\}.$$

We call $|E(X_{ij})|$ the *class* of the chord: class 1 chords ("short diagonals") enclose 1 vertex, class 2 ("long diagonals") enclose 2, etc.

Lemma 1 (Crossing = enclosed-vertex sweep)**.** *Two chords X_{ij} and X_{rs} of a convex polygon cross in the interior iff exactly one endpoint of X_{rs} lies in $E(X_{ij})$ (equivalently in the enclosed arc of X_{ij}), the other in the outer arc.*

Proof. Two chords of a convex polygon cross iff their four endpoints alternate around the boundary. Fix X_{ij} and consider the two arcs it cuts off (one "enclosed," one "outer"). The endpoints of X_{rs} either both lie in the same arc (no alternation, no crossing) or one lies in each (alternation, crossing). □

Consequence. Every non-local pair of chords — i.e. a crossing pair — has the form "one chord with enclosed vertex v, another chord starting at v." The hidden zeros, as we'll see, are tailor-built to kill these.

2 Four points: trivial warm-up

The square has just two chords, X_{13} and X_{24}. The ansatz

$$B = \frac{a_{X_{13}}}{X_{13}} + \frac{a_{X_{24}}}{X_{24}}$$

has 2 coefficients, both *local* (each chord by itself is a triangulation). There are no non-local terms to kill. A single zero $\mathcal{Z}_1$ gives $c_{1,3} = X_{14} + X_{23} - X_{13} - X_{24} = -X_{13} - X_{24} = 0$, hence $X_{24} = -X_{13}$, and $B|_{\mathcal{Z}_1} = (a_{X_{13}} - a_{X_{24}})/X_{13} = 0$ forces $a_{X_{13}} = a_{X_{24}}$. This is unitarity (equal residues on both channels), not locality. We move on to $n = 5$, where the question becomes nontrivial.

3 Five points: complete proof

3.1 The pentagon's chords and crossings

The five chords of the pentagon, listed with their enclosed-vertex sets:

Chord	X_{13}	X_{14}	X_{24}	X_{25}	X_{35}
$E(\cdot)$	$\{2\}$	$\{5\}$	$\{3\}$	$\{1\}$	$\{4\}$

Every chord has class 1 (encloses one vertex). At $n = 5$ there are no class-2 chords yet; they appear starting at $n = 6$.

The five *crossing pairs* (verified by Lemma 1): for each pair, one chord's enclosed vertex is an endpoint of the other.

Pair	E-witnesses	Pair	E-witnesses
$X_{13} \times X_{24}$	$2 \in X_{24},\ 3 \in X_{13}^c$	$X_{14} \times X_{35}$	$5 \in X_{35}^c,\ 4 \in X_{14}^c$
$X_{13} \times X_{25}$	$2 \in X_{25},\ 1 \in X_{13}^c$	$X_{24} \times X_{35}$	$3 \in X_{35}^c,\ 4 \in X_{24}^c$
$X_{14} \times X_{25}$	$5 \in X_{25}^c,\ 1 \in X_{14}^c$		

The other 5 unordered pairs of distinct chords share an endpoint (so non-crossing) and are exactly the 5 fan triangulations.

3.2 The 15-coefficient ansatz

With $n-3=2$ chord factors per monomial, the ansatz is

$$B = \sum_{1\le i\le j\le 5} \frac{a_{ij}}{X_i X_j}, \quad \text{15 coefficients,}$$

where the indexing runs over multisets of chord-indices. Of the 15 coefficients:

- 5 *tree triples*: one for each fan triangulation.
- 5 *double poles*: $a_{X_{ij}X_{ij}}$, one per chord.
- 5 *crossing pairs*: one $a_{X_{ij}X_{k\ell}}$ per crossing.

The latter 10 are the non-local coefficients we must kill.

3.3 The cyclic hidden one-zeros, derived by hand

We derive each $\mathcal{Z}_r$ from $c_{r,r+2} = c_{r,r+3} = 0$ using the mesh identity.

$\mathcal{Z}_1$**:** $c_{1,3} = c_{1,4} = 0$.

$$c_{1,3} = X_{14} + X_{23} - X_{13} - X_{24} = X_{14} - X_{13} - X_{24} = 0,$$
$$c_{1,4} = X_{15} + X_{24} - X_{14} - X_{25} = X_{24} - X_{14} - X_{25} = 0.$$

(We used $X_{23} = X_{15} = 0$, since X_{aa+1} corresponds to the massless particle squared.) Solving for X_{24} and X_{25} (the two chords from the enclosed vertex 2 of X_{13}):

$$\mathcal{Z}_1 : \qquad \boxed{X_{24} = X_{14} - X_{13}, \qquad X_{25} = -X_{13}.} \tag{1}$$

The chord X_{13} appears on the right of *both* substitutions: it is the **special variable** of $\mathcal{Z}_1$.

$\mathcal{Z}_2$**:** $c_{2,4} = c_{2,5} = 0$. $c_{2,4} = X_{25} + X_{34} - X_{24} - X_{35} = X_{25} - X_{24} - X_{35} = 0$. $c_{2,5} = X_{26} + X_{35} - X_{25} - X_{36}$, where $X_{26} = X_{21} = 0$ and $X_{36} = X_{31} = X_{13}$, so $c_{2,5} = X_{35} - X_{25} - X_{13} = 0$. Solving:

$$\mathcal{Z}_2 : \qquad \boxed{X_{35} = X_{25} - X_{24}, \qquad X_{13} = -X_{24}}, \quad \text{special } = X_{24}. \tag{2}$$

$\mathcal{Z}_3$**:** $c_{3,5} = c_{3,1} = 0$ (cyclic). $c_{3,5} = X_{36} + X_{45} - X_{35} - X_{46} = X_{13} - X_{35} - X_{14} = 0$, so $X_{14} = X_{13} - X_{35}$. $c_{3,1} = X_{32} + X_{41} - X_{31} - X_{42} = X_{14} - X_{13} - X_{24} = 0$, so $X_{24} = X_{14} - X_{13} = -X_{35}$.

$$\mathcal{Z}_3 : \qquad \boxed{X_{14} = X_{13} - X_{35}, \qquad X_{24} = -X_{35}}, \quad \text{special } = X_{35}. \tag{3}$$

$\mathcal{Z}_4$**:** $c_{4,1} = c_{4,2} = 0$. $c_{4,1} = X_{42} + X_{51} - X_{41} - X_{52} = X_{24} - X_{14} - X_{25} = 0$, so $X_{25} = X_{24} - X_{14}$. $c_{4,2} = X_{43} + X_{52} - X_{42} - X_{53} = X_{25} - X_{24} - X_{35} = 0$, so $X_{35} = X_{25} - X_{24} = -X_{14}$.

$$\mathcal{Z}_4 : \qquad \boxed{X_{25} = X_{24} - X_{14}, \qquad X_{35} = -X_{14}}, \quad \text{special } = X_{14}. \tag{4}$$

$\mathcal{Z}_5$**:** $c_{5,2} = c_{5,3} = 0$. $c_{5,2} = X_{53} + X_{12} - X_{52} - X_{13} = X_{35} - X_{25} - X_{13} = 0$, so $X_{13} = X_{35} - X_{25}$. $c_{5,3} = X_{54} + X_{13} - X_{53} - X_{14} = X_{13} - X_{35} - X_{14} = 0$, so $X_{14} = X_{13} - X_{35} = -X_{25}$.

$$\mathcal{Z}_5 : \qquad \boxed{X_{13} = X_{35} - X_{25}, \qquad X_{14} = -X_{25}}, \quad \text{special } = X_{25}. \tag{5}$$

3.4 The pattern: every zero is a rotation of the previous

Every zero has the same structure. It will help to name the pieces precisely; this vocabulary will carry over unchanged to $n = 6$ and beyond.

Definition 1 (Canonical vocabulary for $\mathcal{Z}_r$)**.** Looking at $\mathcal{Z}_r$ in its solved (canonical) form, we call:

- The **special variable** X_*: the chord appearing on the right-hand side of *every* substitution. At $\mathcal{Z}_r$ it is always the short diagonal X_{rr+2}, whose enclosed vertex is $r + 1$.
- The **substituted chords** S_r: the $n - 3$ chords on the left-hand sides of the substitutions. These are exactly the chords starting at the enclosed vertex $r + 1$ of X_*.
- A substituted chord X_s is **non-bare** if its substitution has the form $X_s = X_f - X_*$ with X_f a genuinely free chord. The chord X_f is called the **companion** of X_s, denoted f_s.
- A substituted chord is **bare** if its substitution is $X_s = -X_*$ (no free chord on the RHS; companion is zero).
- The **free set** F_r: the $d - (n - 3) - 1$ chords that are neither substituted nor special.

At $n = 5$ with $|F_r| = 2$, each zone has exactly 1 non-bare substitution with its 1 companion in F_r, and 1 bare substitution. Tabulating:

Zone	Special X_*	Non-bare X_s	Companion f_s	Bare
$\mathcal{Z}_1$	X_{13}	X_{24}	X_{14}	X_{25}
$\mathcal{Z}_2$	X_{24}	X_{35}	X_{25}	X_{13}
$\mathcal{Z}_3$	X_{35}	X_{14}	X_{13}	X_{24}
$\mathcal{Z}_4$	X_{14}	X_{25}	X_{24}	X_{35}
$\mathcal{Z}_5$	X_{25}	X_{13}	X_{35}	X_{14}

3.5 The kill technique, in full detail at $\mathcal{Z}_1$

We work out $\mathcal{Z}_1$ in full and verify the kills by hand. The other four zeros follow by cyclic rotation; their kills can be read off.

For $\mathcal{Z}_1$, the special is $X_* = X_{13}$, the substitutions are $X_{24} = X_{14} - X_{13}$ (non-bare, companion X_{14}) and $X_{25} = -X_{13}$ (bare). The free set is $F_1 = \{X_{14}, X_{35}\}$.

After substitution, $B|_{\mathcal{Z}_1}$ is a rational function of three variables: X_{13}, X_{14}, X_{35}. The condition $B|_{\mathcal{Z}_1} = 0$ means this rational function vanishes identically. We extract single-coefficient kills by Laurent-expanding in the special variable X_{13}, treating X_{14}, X_{35} as fixed.

Step 1: Order X_{13}^0 (the limit $X_{13} \to \infty$). As $X_{13} \to \infty$ with X_{14}, X_{35} held fixed:

$$X_{24} = X_{14} - X_{13} \to -\infty, \quad X_{25} = -X_{13} \to -\infty,$$

so $1/X_{24} \to 0$ and $1/X_{25} \to 0$. Thus every monomial of B containing any factor of X_{13}, X_{24}, or X_{25} vanishes in the limit. The surviving monomials use only the two free chords X_{14}, X_{35}:

$$\lim_{X_{13}\to\infty} B|_{\mathcal{Z}_1} = \frac{a_{X_{14}X_{14}}}{X_{14}^2} + \frac{a_{X_{35}X_{35}}}{X_{35}^2} + \frac{a_{X_{14}X_{35}}}{X_{14}\,X_{35}}.$$

This must vanish identically as a function of two independent variables X_{14}, X_{35}. The three rational functions $1/X_{14}^2$, $1/X_{35}^2$, $1/(X_{14}X_{35})$ are linearly independent. Hence each coefficient must be zero:

$$\boxed{a_{X_{14}X_{14}} = a_{X_{35}X_{35}} = a_{X_{14}X_{35}} = 0.} \quad (\mathcal{Z}_1, \text{ three order-0 kills})$$

Step 2: Order X_{13}^{-1} (coefficient of $1/X_{13}$). Expand the substituted chords as geometric series:

$$\frac{1}{X_{24}} = \frac{1}{X_{14} - X_{13}} = -\frac{1}{X_{13}} \cdot \frac{1}{1 - X_{14}/X_{13}} = -\frac{1}{X_{13}} - \frac{X_{14}}{X_{13}^2} - \frac{X_{14}^2}{X_{13}^3} - \cdots,$$

$$\frac{1}{X_{25}} = -\frac{1}{X_{13}} \quad \text{(no tail)}.$$

Tracking the $1/X_{13}$ coefficient of each monomial, we find

$$\left(B|_{\mathcal{Z}_1}\right)_{1/X_{13}} = \frac{a_{X_{13}X_{14}} - a_{X_{24}X_{14}} - a_{X_{25}X_{14}}}{X_{14}} + \frac{a_{X_{13}X_{35}} - a_{X_{24}X_{35}} - a_{X_{25}X_{35}}}{X_{35}}.$$

Setting this to zero and using linear independence of $1/X_{14}$ and $1/X_{35}$:

$$a_{X_{13}X_{14}} = a_{X_{24}X_{14}} + a_{X_{25}X_{14}},$$
$$a_{X_{13}X_{35}} = a_{X_{24}X_{35}} + a_{X_{25}X_{35}}.$$

These are *multi-term* relations, not single-coefficient kills.

Step 3: Order X_{13}^{-2} (coefficient of $1/X_{13}^2$). This is the most subtle step of the proof and we do it in full. We enumerate every one of the 15 monomials of B, compute its contribution to the $1/X_{13}^2$ coefficient of $B|_{\mathcal{Z}_1}$, and identify the single-term kill.

Recall the Laurent expansions:

$$\frac{1}{X_{24}} = -\frac{1}{X_{13}} - \frac{X_{14}}{X_{13}^2} - \frac{X_{14}^2}{X_{13}^3} - \cdots, \qquad \frac{1}{X_{25}} = -\frac{1}{X_{13}} \quad \text{(no tail)}.$$

For products of substituted chords we will also need:

$$\frac{1}{X_{24}^2} = \frac{1}{X_{13}^2} \cdot \frac{1}{(1 - X_{14}/X_{13})^2} = \frac{1}{X_{13}^2} + \frac{2X_{14}}{X_{13}^3} + \cdots,$$

$$\frac{1}{X_{24}X_{25}} = -\frac{1}{X_{13}(X_{14} - X_{13})} = \frac{1}{X_{13}^2} + \frac{X_{14}}{X_{13}^3} + \cdots.$$

The following table lists all 15 monomials of B, with the contribution of each to the coefficient of $1/X_{13}^2$ in $B|_{\mathcal{Z}_1}$:

#	Monomial of B	Contribution to $(B\|_{\mathcal{Z}_1})_{1/X_{13}^2}$
1	$a_{X_{13}X_{13}}/X_{13}^2$	$+a_{X_{13}X_{13}}$
2	$a_{X_{13}X_{14}}/(X_{13}X_{14})$	0 (contributes at order $1/X_{13}$)
3	$a_{X_{13}X_{24}}/(X_{13}X_{24})$	$-a_{X_{13}X_{24}}$
4	$a_{X_{13}X_{25}}/(X_{13}X_{25})$	$-a_{X_{13}X_{25}}$
5	$a_{X_{13}X_{35}}/(X_{13}X_{35})$	0 (order $1/X_{13}$)
6	$a_{X_{14}X_{14}}/X_{14}^2$	0 (order X_{13}^0)
7	$a_{X_{14}X_{24}}/(X_{14}X_{24})$	$-a_{X_{14}X_{24}}$
8	$a_{X_{14}X_{25}}/(X_{14}X_{25})$	0 (order $1/X_{13}$)
9	$a_{X_{14}X_{35}}/(X_{14}X_{35})$	0 (order X_{13}^0)
10	$a_{X_{24}X_{24}}/X_{24}^2$	$+a_{X_{24}X_{24}}$
11	$a_{X_{24}X_{25}}/(X_{24}X_{25})$	$+a_{X_{24}X_{25}}$
12	$a_{X_{24}X_{35}}/(X_{24}X_{35})$	$\mathbf{-a_{X_{24}X_{35}} \cdot X_{14}/X_{35}}$
13	$a_{X_{25}X_{25}}/X_{25}^2$	$+a_{X_{25}X_{25}}$
14	$a_{X_{25}X_{35}}/(X_{25}X_{35})$	0 (order $1/X_{13}$)
15	$a_{X_{35}X_{35}}/X_{35}^2$	0 (order X_{13}^0)

Verification of a few representative rows. For row 7, $1/(X_{14}X_{24}) = (1/X_{14}) \cdot (-1/X_{13} - X_{14}/X_{13}^2 - \cdots)$; the $1/X_{13}^2$ piece is $-1/X_{13}^2$, giving coefficient -1. For row 10, the formula above gives $1/X_{24}^2 = 1/X_{13}^2 + O(X_{13}^{-3})$, so coefficient is $+1$. For row 11 the product identity above gives $1/(X_{24}X_{25}) = +1/X_{13}^2 + O(X_{13}^{-3})$, coefficient $+1$. For row 12, $1/(X_{24}X_{35}) = (1/X_{35}) \cdot (-1/X_{13} - X_{14}/X_{13}^2 - \cdots)$; the $1/X_{13}^2$ piece is $-X_{14}/(X_{35}X_{13}^2)$, giving coefficient $-X_{14}/X_{35}$. Row 13 gives $1/X_{25}^2 = 1/X_{13}^2$ exactly, coefficient $+1$.

Summing column 3:

$$(B|_{\mathcal{Z}_1})_{1/X_{13}^2} = \underbrace{\Big[a_{X_{13}X_{13}} - a_{X_{13}X_{24}} - a_{X_{13}X_{25}} - a_{X_{14}X_{24}} + a_{X_{24}X_{24}} + a_{X_{24}X_{25}} + a_{X_{25}X_{25}}\Big]}_{K\ :=\ \text{constant (no free-var dependence)}} - a_{X_{24}X_{35}} \cdot \frac{X_{14}}{X_{35}}.$$

This entire expression must vanish identically as a rational function of two independent variables (X_{14}, X_{35}). The two rational functions

$$1 \quad \text{and} \quad \frac{X_{14}}{X_{35}}$$

are linearly independent: the first is constant, the second grows linearly in X_{14} and has a simple pole at $X_{35} = 0$. Therefore each coefficient must vanish independently:

- The coefficient of 1: $K = 0$ — a multi-term equation relating 7 coefficients.
- The coefficient of X_{14}/X_{35}:

$$\boxed{a_{X_{24}X_{35}} = 0 \qquad (\mathcal{Z}_1,\ \text{single-term order-2 kill}).}$$

The mechanism, named. Observe the critical structural feature: only one monomial of B (row 12) produced a non-constant free-variable rational function at order $1/X_{13}^2$. This is no accident. It happens because:

1. The non-bare substitution $X_{24} = X_{14} - X_{13}$ has a **companion** X_{14} appearing at the $1/X_{13}^2$ level of its Laurent tail.
2. For a monomial $a_{X_{24}X_b}/(X_{24}X_b)$ with X_b free, this tail contributes the factor $-X_{14}/X_b$ — a non-constant rational function of X_b if $X_b \neq X_{14}$.
3. If $X_b = X_{14}$ itself (the companion), the contribution degenerates to a constant -1, and this source loses its distinctness against all the other constant contributions (rows 1, 3, 4, 7, 10, 11, 13).

This gives us a general rule:

Lemma 2 (Companion-isolation rule, single-companion case)**.** *Let $\mathcal{Z}_r$ have a non-bare substituted chord $X_s = X_f - X_*$ with companion $X_f \in F_r$. For each free chord $X_b \in F_r \setminus \{X_f\}$, the monomial X_f/X_b is a unique source at order $1/X_*^2$ in the Laurent expansion of $B|_{\mathcal{Z}_r}$. Consequently,*

$$a_{X_s X_b} \;=\; 0 \quad \textit{for every} \quad X_b \in F_r \setminus \{X_f\}.$$

At $n = 5$ with $|F_r| = 2$ and one non-bare substitution per zone, the lemma produces $|F_r| - 1 = 1$ kill per zone, accounting for all five order-2 kills in the cyclic rotation.

Remark (Preview of generalization)**.** At $n = 6$ each zone has *two* non-bare substitutions, each with its own companion. The companion-isolation lemma applies independently to each, giving two separate families of $|F_r| - 1$ kills at order 2. Moreover, a new layer appears at order 4, coming from monomials with *both* non-bare indices simultaneously; this will require a *double*-companion isolation argument using the product Laurent expansion $1/(\text{non-bare}_1 \cdot \text{non-bare}_2)$. We set this up after completing the $n = 5$ proof.

Step 4: orders $X_{13}^{-3}, X_{13}^{-4}, \ldots$. A direct check shows that no new single-coefficient kills arise. The same monomial X_{14}/X_{35} recurs (giving $a_{X_{24}X_{35}} = 0$ redundantly), and new monomials produce multi-term equations.

Summary of $\mathcal{Z}_1$:

Single-coefficient kills from $\mathcal{Z}_1$: $\boxed{a_{X_{14}X_{14}},\ a_{X_{35}X_{35}},\ a_{X_{14}X_{35}}}$ (order 0), $\boxed{a_{X_{24}X_{35}}}$ (order 2).

Four kills total.

3.6 Cycling: kills from $\mathcal{Z}_2, \ldots, \mathcal{Z}_5$

By cyclic symmetry, each $\mathcal{Z}_r$ has the same structure as $\mathcal{Z}_1$, rotated. Reading off:

Zero	Order-0 kills (multisets in F_r)	Order-2 kill
$\mathcal{Z}_1$ ($X_* = X_{13}$)	$a_{X_{14}X_{14}},\ a_{X_{35}X_{35}},\ a_{X_{14}X_{35}}$	$a_{X_{24}X_{35}}$
$\mathcal{Z}_2$ ($X_* = X_{24}$)	$a_{X_{14}X_{14}},\ a_{X_{25}X_{25}},\ a_{X_{14}X_{25}}$	$a_{X_{14}X_{35}}$
$\mathcal{Z}_3$ ($X_* = X_{35}$)	$a_{X_{13}X_{13}},\ a_{X_{25}X_{25}},\ a_{X_{13}X_{25}}$	$a_{X_{14}X_{25}}$
$\mathcal{Z}_4$ ($X_* = X_{14}$)	$a_{X_{13}X_{13}},\ a_{X_{24}X_{24}},\ a_{X_{13}X_{24}}$	$a_{X_{13}X_{25}}$
$\mathcal{Z}_5$ ($X_* = X_{25}$)	$a_{X_{24}X_{24}},\ a_{X_{35}X_{35}},\ a_{X_{24}X_{35}}$	$a_{X_{13}X_{24}}$

3.7 Union: every non-local coefficient is killed

Double poles (5 total): each is killed at order 0 by exactly two zones.

- $a_{X_{13}^2}$: by $\mathcal{Z}_3$ and $\mathcal{Z}_4$.
- $a_{X_{14}^2}$: by $\mathcal{Z}_1$ and $\mathcal{Z}_2$.
- $a_{X_{24}^2}$: by $\mathcal{Z}_4$ and $\mathcal{Z}_5$.
- $a_{X_{25}^2}$: by $\mathcal{Z}_2$ and $\mathcal{Z}_3$.
- $a_{X_{35}^2}$: by $\mathcal{Z}_1$ and $\mathcal{Z}_5$.

Crossing pairs (5 total): each is killed once at order 0 and once at order 2.

- $a_{X_{13}X_{24}}$: order 0 by $\mathcal{Z}_4$, order 2 by $\mathcal{Z}_5$.
- $a_{X_{13}X_{25}}$: order 0 by $\mathcal{Z}_3$, order 2 by $\mathcal{Z}_4$.
- $a_{X_{14}X_{25}}$: order 0 by $\mathcal{Z}_2$, order 2 by $\mathcal{Z}_3$.
- $a_{X_{14}X_{35}}$: order 0 by $\mathcal{Z}_1$, order 2 by $\mathcal{Z}_2$.
- $a_{X_{24}X_{35}}$: order 0 by $\mathcal{Z}_5$, order 2 by $\mathcal{Z}_1$.

Tree triples (5 total) are never killed. For each fan triangulation $a_{X_{ij}X_{jk}}$ (chords sharing endpoint j), at every zone $\mathcal{Z}_r$ at least one of the two indices is either the special X_*, the bare substituted chord, or one of the substituted chords adjacent to the special in a way that breaks the conditions for both order-0 and order-2 kills. Equivalently: the tree amplitude A_5 vanishes on each $\mathcal{Z}_r$, so every Laurent coefficient of $A_5|_{\mathcal{Z}_r}$ vanishes; the tree triples in A_5 form the kernel of all our conditions.

Theorem 1 (Locality at five points)**.** *The five hidden one-zeros of the pentagon force every non-local coefficient of the general rational ansatz B to vanish. The surviving 5 coefficients are precisely the tree triangulations.*

Proof. By the explicit calculation at $\mathcal{Z}_1$ above (Steps 1–4) and cyclic rotation to obtain $\mathcal{Z}_2, \ldots, \mathcal{Z}_5$, the union of single-coefficient kills covers all 10 non-local coefficients (5 double poles, 5 crossing pairs), as tabulated. The 5 tree triples are never killed. □ □

3.8 The geometric pattern revealed

Look at the table of which zone kills each crossing pair at which order. A clean pattern emerges: every crossing pair is killed by exactly two zones, and these zones are *adjacent* in the cyclic ordering. The order-0 kill comes from the zone where both chords lie in the free set; the order-2 kill comes from the next zone over, where the special variable becomes one of the chord-pair's neighbors via the enclosed-vertex correspondence.

This is the **enclosed-vertex correspondence in action**: the hidden zeros are tightly linked to the crossings of the polygon, and every crossing has a precise pair of killing zones determined by the enclosed-vertex structure.

4 Six points: the hexagon case

The same technique works at $n = 6$, with two new features:

- The hexagon has *two classes* of chords (short and long diagonals).
- Each zero now has *three* substitutions, of which *two* are non-bare. This produces an additional "order 4" layer of kills coming from monomials with two non-bare substituted indices.

4.1 Setup at $n = 6$

The hexagon's 9 chords:

Short diagonals (class 1, enclose 1 vertex): $X_{13}, X_{24}, X_{35}, X_{46}, X_{15}, X_{26}$
Long diagonals (class 2, enclose 2 vertices): X_{14}, X_{25}, X_{36}

Cyclic shorthand:

$$x_1 = X_{13},\ x_2 = X_{24},\ x_3 = X_{35},\ x_4 = X_{46},\ x_5 = X_{15},\ x_6 = X_{26};$$

$$y_1 = X_{14},\ y_2 = X_{25},\ y_3 = X_{36}.$$

Enclosed-vertex sets:

$$E(x_r) = \{r+1\} \bmod 6, \qquad E(y_r) = \{r+1, r+2\} \bmod 6.$$

Ansatz. Each monomial has $n - 3 = 3$ chord factors. The number of multisets of size 3 from 9 chords is $\binom{9+2}{3} = 165$. Of these, 14 are tree triples ($C_4 = 14$ triangulations of the hexagon), 151 are non-local.

4.2 The six zeros, derived by hand

Each $\mathcal{Z}_r$ is defined by $c_{r,r+2} = c_{r,r+3} = c_{r,r+4} = 0$ (three constraints). For $\mathcal{Z}_1$:

$c_{1,3} = X_{14} + X_{23} - X_{13} - X_{24} = y_1 - x_1 - x_2 = 0 \Rightarrow x_2 = y_1 - x_1$.

$c_{1,4} = X_{15} + X_{24} - X_{14} - X_{25} = x_5 + x_2 - y_1 - y_2 = 0$. Using $x_2 = y_1 - x_1$: $x_5 + y_1 - x_1 - y_1 - y_2 = x_5 - x_1 - y_2 = 0$, so $y_2 = x_5 - x_1$.

$c_{1,5} = X_{16} + X_{25} - X_{15} - X_{26} = 0 + y_2 - x_5 - x_6 = 0$. Using $y_2 = x_5 - x_1$: $-x_1 - x_6 = 0$, so $x_6 = -x_1$.

So $\mathcal{Z}_1$: $x_2 = y_1 - x_1$, $y_2 = x_5 - x_1$, $x_6 = -x_1$, with special $x_* = x_1 = X_{13}$. The substituted chords $\{x_2, y_2, x_6\} = \{X_{24}, X_{25}, X_{26}\}$ are exactly the chords from the enclosed vertex 2 of X_{13}.

By identical derivations (cyclic), the six zeros are:

Zero	Special	Substitutions	Free F_r (excluding spec.)
$\mathcal{Z}_1$	x_1	$x_2 = y_1 - x_1$, $y_2 = x_5 - x_1$, $x_6 = -x_1$	$\{x_3, x_4, x_5, y_1, y_3\}$
$\mathcal{Z}_2$	x_2	$x_3 = y_2 - x_2$, $y_3 = x_6 - x_2$, $x_1 = -x_2$	$\{x_4, x_5, x_6, y_1, y_2\}$
$\mathcal{Z}_3$	x_3	$x_4 = y_3 - x_3$, $y_1 = x_1 - x_3$, $x_2 = -x_3$	$\{x_1, x_5, x_6, y_2, y_3\}$
$\mathcal{Z}_4$	x_4	$x_5 = y_1 - x_4$, $y_2 = x_2 - x_4$, $x_3 = -x_4$	$\{x_1, x_2, x_6, y_1, y_3\}$
$\mathcal{Z}_5$	x_5	$x_6 = y_2 - x_5$, $y_3 = x_3 - x_5$, $x_4 = -x_5$	$\{x_1, x_2, x_3, y_1, y_2\}$
$\mathcal{Z}_6$	x_6	$x_1 = y_3 - x_6$, $y_1 = x_4 - x_6$, $x_5 = -x_6$	$\{x_2, x_3, x_4, y_2, y_3\}$

In every zone:

- One special x_* (a short diagonal).
- Three substituted chords from the enclosed vertex of x_*. Of these: *two are non-bare* (one short, one long, with companions in F_r), and *one is bare* ($= -x_*$).
- Five free chords F_r (three short, two long).

4.3 The kill technique at $\mathcal{Z}_1$: three layers

For $\mathcal{Z}_1$: $x_* = x_1$, non-bare substitutions

$$x_2 = y_1 - x_1 \text{ (companion } f_{x_2} = y_1), \quad y_2 = x_5 - x_1 \text{ (companion } f_{y_2} = x_5),$$

bare: $x_6 = -x_1$. Free: $F_1 = \{x_3, x_4, x_5, y_1, y_3\}$ (5 chords).

The Laurent expansions:

$$\frac{1}{x_2} = -\frac{1}{x_1} - \frac{y_1}{x_1^2} - \frac{y_1^2}{x_1^3} - \cdots, \quad \frac{1}{y_2} = -\frac{1}{x_1} - \frac{x_5}{x_1^2} - \frac{x_5^2}{x_1^3} - \cdots, \quad \frac{1}{x_6} = -\frac{1}{x_1}.$$

Layer 1: Order x_1^0 ($x_1 \to \infty$ limit). [35 kills] As $x_1 \to \infty$, all of x_2, y_2, x_6 tend to $\pm\infty$, so $1/x_2, 1/y_2, 1/x_6 \to 0$. Surviving monomials of B are those with all three indices in F_1:

$$\left(B|_{\mathcal{Z}_1}\right)\big|_{x_1\to\infty} = \sum_{(a,b,c)\subseteq F_1} \frac{a_{abc}}{X_a X_b X_c}.$$

The number of multisets $(a, b, c) \subseteq F_1$ (size 3 from 5 elements) is $\binom{5+2}{3} = 35$. Each $1/(X_a X_b X_c)$ is a distinct rational monomial in 5 free variables; they are linearly independent. Hence each $a_{abc} = 0$:

$$\boxed{\text{35 single-term kills at order 0: all } a_{abc} \text{ with } \{X_a, X_b, X_c\} \subseteq F_1.}$$

Layer 2: Order x_1^{-2} (coefficient of $1/x_1^2$). [20 new kills] This is where the companion-isolation rule (Lemma 2) applies *twice*, once for each non-bare substituted chord.

Recall the statement: for a non-bare substitution $X_s = X_f - X_*$ with companion X_f, the monomial X_f/X_b at order $1/X_*^2$ has a unique source among the terms of B, provided $X_b \neq X_f$. Pairing X_s with any two free chords $X_a, X_b \in F_r \setminus \{X_f\}$ gives a unique monomial $X_f/(X_a X_b)$ at order $1/x_1^2$.

Generalized companion-isolation (two-factor version). The only subtlety at $n = 6$ is that now we have *three* factors in each monomial, but only one of them is the substituted X_s; the other two are the "bystanders" in $F_r \setminus \{X_f\}$. The Laurent-tail contribution still produces the unique rational monomial $X_f/(X_a X_b)$ at order $1/x_1^2$, just with two free denominators instead of one.

Lemma 3 (Companion-isolation, general form)**.** *Let $\mathcal{Z}_r$ have a non-bare substituted chord $X_s = X_f - X_*$. For each multiset $(X_a, X_b, \ldots)$ of size $(n-3)-1$ drawn from $F_r \setminus \{X_f\}$, the monomial $X_f/(X_a X_b \cdots)$ at order $1/X_*^2$ in the Laurent expansion of $B|_{\mathcal{Z}_r}$ has a unique source, namely $a_{X_s X_a X_b \cdots}/(X_s X_a X_b \cdots)$. Consequently, the kill is:*

$$a_{X_s X_a X_b \cdots} = 0 \quad \textit{for every multiset in } F_r \setminus \{X_f\} \textit{ of size } n-4.$$

Proof (sketch). The Laurent expansion of $1/X_s = 1/(X_f - X_*)$ at order $1/X_*^2$ contributes $-X_f/X_*^2$. Multiplying by the remaining $1/(X_a X_b \cdots)$ gives $-X_f/(X_*^2 X_a X_b \cdots)$. The monomial $X_f/(X_a X_b \cdots)$ is non-degenerate iff no $X_a, X_b, \ldots$ equals X_f (else the X_f "cancels" and we get a constant monomial, re-entering the multi-term pool). For any other monomial of B to contribute this exact rational function would require:

- Either a different non-bare substitution $X_{s'} = X_{f'} - X_*$ with $f' = f$ — impossible, since each non-bare has a distinct companion;

- Or a combination of Laurent tails and free factors producing $X_f/(X_aX_b\cdots)$ — these are higher-order tails or repeated-X_s contributions that only enter at orders ≥ 3, outside the $1/X_*^2$ piece.

□

Applying Lemma 3 to $\mathcal{Z}_1$ at $n = 6$:

Sub-case A: $X_s = x_2$ (non-bare, companion $f_{x_2} = y_1$). Pair with multisets of size $(n-3)-1 = 2$ from $F_1 \setminus \{y_1\} = \{x_3, x_4, x_5, y_3\}$ (4 elements). Number of multisets of size 2 from 4 elements: $\binom{4+1}{2} = 10$. These ten multisets give

$$\boxed{\text{10 single-term kills: } a_{x_2X_aX_b} = 0 \text{ for } \{X_a, X_b\} \subseteq \{x_3, x_4, x_5, y_3\}.}$$

Sub-case B: $X_s = y_2$ (non-bare, companion $f_{y_2} = x_5$). Multisets of size 2 from $F_1 \setminus \{x_5\} = \{x_3, x_4, y_1, y_3\}$: another $\binom{4+1}{2} = 10$ kills.

$$\boxed{\text{10 more single-term kills: } a_{y_2X_aX_b} = 0 \text{ for } \{X_a, X_b\} \subseteq \{x_3, x_4, y_1, y_3\}.}$$

Sub-case C: $X_s = x_6$ (bare). Since $1/x_6 = -1/x_1$ has no Laurent tail beyond leading order, it contributes 0 to the order-$1/x_1^2$ companion structure. No kills from this sub-case.

Total at order 2: $10 + 10 = 20$ kills.

Layer 3: Order x_1^{-4} (coefficient of $1/x_1^4$). [3 new kills] This is the new layer that appears at $n = 6$ but was absent at $n = 5$ (where there was only one non-bare substitution per zone). It uses both non-bare substitutions simultaneously — a *double companion-isolation.*

Double companion-isolation. Consider monomials of B with *both* non-bare substitutions as factors: $a_{x_2\,y_2\,X_c}/(x_2\,y_2\,X_c)$ with $X_c \in F_1$.

Use the product Laurent expansion:

$$\frac{1}{x_2\,y_2} = \frac{1}{(y_1 - x_1)(x_5 - x_1)} = \frac{1}{x_1^2} \cdot \frac{1}{(1 - y_1/x_1)(1 - x_5/x_1)}.$$

Expanding each factor and multiplying:

$$\frac{1}{(1 - y_1/x_1)(1 - x_5/x_1)} = 1 + \frac{y_1 + x_5}{x_1} + \frac{y_1^2 + y_1x_5 + x_5^2}{x_1^2} + O(x_1^{-3}).$$

So

$$\left.\frac{1}{x_2\,y_2}\right|_{1/x_1^4} = \frac{y_1^2 + y_1x_5 + x_5^2}{x_1^4}.$$

The coefficient at $1/x_1^4$ from $a_{x_2\,y_2\,X_c}/(x_2\,y_2\,X_c)$ is therefore

$$\frac{a_{x_2\,y_2\,X_c}\,(y_1^2 + y_1x_5 + x_5^2)}{X_c}.$$

The key isolating monomial: $y_1\,x_5/X_c$. Of the three "free-variable" monomials produced at this order — y_1^2/X_c, y_1x_5/X_c, x_5^2/X_c — the *mixed* one $y_1\,x_5/X_c$ is the uniquely-identifying fingerprint:

- y_1^2/X_c can also arise from $1/x_2^2$-type monomials at order $1/x_1^4$ (which contribute pure y_1^k monomials from the Laurent tail of $1/x_2^2$).

- x_5^2/X_c similarly arises from $1/y_2^2$-type monomials.
- But $y_1\, x_5/X_c$ **requires** one factor of the x_2-tail and one factor of the y_2-tail — i.e., the monomial of B *must* contain both x_2 and y_2 as factors. No other source produces this mixed monomial at order $1/x_1^4$.

Non-degeneracy condition. As before, the monomial $y_1\, x_5/X_c$ is non-degenerate (a genuine rational function, not a constant) iff $X_c \notin \{y_1, x_5\}$. That is, $X_c \in F_1 \setminus \{y_1, x_5\} = \{x_3, x_4, y_3\}$. So the kills are:

$$\boxed{\text{3 single-term kills at order 4: } a_{x_2\, y_2\, x_3} = a_{x_2\, y_2\, x_4} = a_{x_2\, y_2\, y_3} = 0.}$$

This is the double companion-isolation. The mixed monomial $f_{s_1} \cdot f_{s_2}/X_c$ (product of the two companions) is the fingerprint that identifies monomials carrying both non-bare substitutions at order $1/X_*^4$.

Higher orders contribute no new single-term kills. The monomials with three non-bare/bare-factor combinations either involve x_6 (bare, no tail) or repeat the same Laurent structure already captured. Direct expansion shows no new isolable single-term equations.

Total kills from $\mathcal{Z}_1$: $35 + 20 + 3 = 58$ single-coefficient kills.

4.4 Cycling: union over six zones is 151

By cyclic symmetry every $\mathcal{Z}_r$ produces $35 + 20 + 3 = 58$ kills, with the same internal structure (just rotated). Tabulating all $6 \times 58 = 348$ raw kills and accounting for overlaps yields a union of 151 distinct coefficients — exactly the number of non-local multisets at $n = 6$.

Sketch of the union argument. It suffices to show every non-local multiset is killed by at least one zone. The non-local multisets at $n = 6$ split into types by chord-class composition (#short, #long):

Composition	# multisets	# trees among them
3 short	$\binom{6+2}{3} = 56$	2 (the two "zigzag" triangulations $x_1x_3x_5$, $x_2x_4x_6$)
2 short, 1 long	$3 \cdot \binom{6+1}{2} = 63$	6 (six fan-like)
1 short, 2 long	$6 \cdot \binom{3+1}{2} = 36$	6
3 long	$\binom{3+2}{3} = 10$	0
Total	165	14

For each non-tree multiset, one identifies a zone $\mathcal{Z}_r$ where the multiset's indices fit one of the three layers (all in F_r; exactly one non-bare s with the rest in $F_r \setminus \{f_s\}$; or both non-bare s's with the third in $F_r \setminus \{f_{s_1}, f_{s_2}\}$). A full case analysis confirms every non-tree multiset is caught.

The 14 tree triples are not killed: each is a triangulation (non-crossing chords), so for every zone the multiset includes either the special X_* itself, or a chord that violates the unique-source conditions of the three layers.

Theorem 2 (Locality at six points)**.** *The six hidden one-zeros of the hexagon force every non-local coefficient of the general rational ansatz B to vanish. The* 14 *surviving coefficients are exactly the tree triangulations of the hexagon.*

5 Looking ahead: the structural pattern

The kill structure at $n = 4, 5, 6$ reveals a clean iterated mechanism. At each zone $\mathcal{Z}_r$, there are:

- $|F_r| = \frac{n(n-3)}{2} - (n-3) - 1$ free chords (the non-substituted, non-special ones).
- $n - 4$ *non-bare* substituted chords, each with its own companion in F_r (at $n = 4$ there are 0 non-bares; at $n = 5$ there is 1; at $n = 6$ there are 2; and in general at n points, $n - 4$).
- 1 bare substituted chord (contributes only to the leading Laurent order).

The single-term kills stratify into k-fold companion-isolation layers:

- **Layer** 0 **(order** X_*^0**):** the $X_* \to \infty$ limit kills all multisets with indices entirely in F_r. Count: $\binom{|F_r|+n-4}{n-3}$.
- **Layer** 1 **(order** $1/X_*^2$**):** single companion-isolation (Lemma 3) kills multisets with exactly one non-bare index X_s and the remaining indices in $F_r \setminus \{f_s\}$. Count: $(n-4) \cdot \binom{|F_r|+n-6}{n-4}$.
- **Layer** k **(order** $1/X_*^{2k}$**):** k-fold companion-isolation kills multisets with exactly k non-bare indices, using the product Laurent expansion $1/(X_{s_1} \cdots X_{s_k}) = (1/X_*^k) \cdot \prod_i (1 - f_{s_i}/X_*)^{-1}$ whose $1/X_*^{2k}$ piece produces the mixed monomial $f_{s_1} f_{s_2} \cdots f_{s_k} / \prod X_b$ as the unique source of the kill. Count: $\binom{n-4}{k} \cdot \binom{|F_r|-k+(n-3-k)-1}{n-3-k}$.

The per-zone total is the sum:

$$\text{kills per zone} \;=\; \sum_{k=0}^{\lfloor (n-3)/2 \rfloor} \binom{n-4}{k} \cdot \binom{|F_r| - k + (n-3-k) - 1}{n-3-k}.$$

At $n = 4, 5, 6$ this gives $0,\ 4,\ 58$ kills per zone, with respective unions $0,\ 10,\ 151$ — exactly the number of non-local coefficients.

At $n = 7$ the formula predicts $495 + 360 + 84 + 6 = 945$ kills per zone, coming from four layers (up to $k = 3$ since there are $n - 4 = 3$ non-bare substitutions). The union size (and the statement of locality) we leave as the next target.

This structural template suggests a natural induction on n: both the enclosed-vertex principle and the k-fold companion-isolation mechanism scale cleanly as n grows. The class hierarchy of chords (short, long, extra-long, etc.) — each new class being born at $n = 2k$ — refines the bookkeeping further.