

## 2. Probability Distribution

## 2.0. Introduction

- 1) Chapter Scope
  - A. Examples of probability distributions
  - B. Their properties
- 2) Purpose of Introducing Distributions
  - A. a building blocks for more complex models
  - B. a recipe to discuss some essential statistical concept, e.g., Bayesian inference
  - C. to model the probability distribution  $p(\mathbf{x})$ , i.e., density estimation
    - \* Model Selection becomes an issue since density estimation is fundamentally ill-posed problem in that infinitely many distributions can fit the observed data set.
- 3) Parametric distribution vs. Non-Parametric distribution
  - A. Parametric distribution
    - i. binomial distribution, multinomial distribution, Gaussian distribution (continuous R.V.)
    - ii. For density estimation, the parameters shall be determined with an observed data set.
      1. Frequentist: specific values for parameters (earned by optimizing some criterion, e.g., likelihood function)
      2. Bayesian: estimate posterior distribution with introduced prior distributions over the parameters as well as the observed data
    - iii. Conjugate Priors: To simplify the Bayesian analysis, use conjugate prior which let posterior distribution be in the same form of prior distribution.
      1. Exponential family of distributions is presented as it possesses a number of important properties.
  - B. Non-Parametric distribution
    - i. Distribution form is not forced by a user but typically depends on the size of the data set
    - ii. Still has the parameters but they do not determine the distribution form but the complexity
    - iii. Histogram, nearest-neighbors, kernels

**Table. 1 Conjugate prior with posterior distribution in exponential family**

Conjugate Prior	Posterior Distribution
Dirichlet distribution	Multinomial distribution
Gaussian distribution	Gaussian distribution

## 2.1. Binary Variables

### 2.1.1. Bernoulli distribution

#### 2.1.1.1. Definition

$$\text{Bern}(x|\mu) = \mu^x(1-\mu)^{1-x}, \text{ where } 0 \leq \mu < 1 \text{ and } x \in \{0, 1\} \quad (2.1)$$

#### 2.1.1.2. Properties

$$\mathbb{E}[x] = \mu \quad (2.2)$$

$$\text{var}[x] = \mu(1-\mu) \quad (2.3)$$

#### 2.1.1.3. Density estimation

$$\mathcal{D} = \{x_1, \dots, x_N\}$$

##### 2.1.1.3.1. Frequentist

- 1) Estimate  $\mu$  by maximizing the likelihood function, i.e., maximize the log of likelihood

$$\ln(p(\mathcal{D}|\mu)) = \sum_{n=1}^N \ln p(x_n|\mu) = \sum_{n=1}^N \{x_n \ln \mu + (1-x_n) \ln(1-\mu)\} \quad (2.4)$$

- 2) The above log likelihood function depends on the N observations only through their sum, i.e., *sufficient statistics*:  $\sum_n x_n$ .

- 3)  $\mu_{ML} = \frac{m}{N} = \text{sample mean}$

##### 2.1.1.3.2. Bayesian

Flip a coin 3 times resulting all heads  $\rightarrow$  what is the reasonable prediction? (overfitting)

##### 2.1.1.3.2.1. Binomial distribution

###### 1) Definition

$$\text{Bin}(m|N, \mu) = \binom{N}{m} \mu^m (1-\mu)^{N-m} \quad (2.5)$$

###### 2) Properties

For independent events, 1) the means of the sum is the sum of the mean and 2) the variance of the sum is the sum of the variance

$$\mathbb{E}[m] = N\mu \quad (2.6)$$

$$\text{var}[m] = N\mu(1-\mu) \quad (2.7)$$

### 2.1.1.3.2.2. The beta distribution (conjugate prior for the binomial distribution)

#### 1) Motivation for the conjugate prior distribution

- A. Prior distribution is required in order to develop a Bayesian treatment.
- B. Make posterior distribution have the same functional form as the prior (conjugacy).

#### 2) Definition

$\text{Beta}(\mu|a, b) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \mu^{a-1}(1-\mu)^{b-1}$ , gamma coefficient for the normalization purpose

$$\mathbb{E}[\mu] = \frac{a}{a+b} \quad (2.8)$$

$$\text{var}[\mu] = \frac{ab}{(a+b)^2(a+b+1)} \quad (2.9)$$

#### 3) Gamma function

$$\Gamma(x) \equiv \int_0^\infty u^{x-1} e^{-u} du \quad (2.10)$$

$$\Gamma(x+1) = x\Gamma(x), \Gamma(1) = 1, \Gamma(x+1) = x! \quad (2.11)$$

a and b controls the distribution of the parameter  $\mu$ , and thus called *hyperparameters*

#### 4) Posterior distribution

The posterior distribution of  $\mu$ : prior distribution(beta)  $\times$  likelihood function(binomial)

→ Normalization

$$p(\mu|m, l, a, b) = \text{Beta}(\mu|a, b) \times \text{Bin}(m|N, \mu) = \frac{\Gamma(m+a+l+b)}{\Gamma(m+a)\Gamma(l+b)} \mu^{m+a-1}(1-\mu)^{l+b-1} \quad (2.12)$$

$$l = N - m = \# \text{ of tails}$$

$$m = \# \text{ of heads}$$

##### A. Sequential nature

- i. a and b (in the prior): *effective number of observations* of  $x = 1$  and  $x = 0$ , respectively
- ii. The posterior can act as the prior if subsequent observation is followed. If following observation is  $x=1(x=0)$ ,  $a(b)$  will increase by 1.
- iii. Bayesian viewpoint raises such sequential approach of learning

B. Prediction of the future outcome

$$p(x = 1|\mathcal{D}) = \int_0^1 p(x = 1|\mu)p(\mu|\mathcal{D})d\mu = \int_0^1 \mu p(\mu|\mathcal{D})d\mu = \mathbb{E}[\mu|\mathcal{D}] \quad (2.13)$$

$$p(x = 1|\mathcal{D}) = \frac{m + a}{m + a + l + b} \quad (2.14)$$

→ Total fraction of an effective number for  $x = 1$  (including both real&fictitious)

### C. Big Data

i. Bayesian result converges to ML (general phenomenon):

$$p(x = 1|\mathcal{D}) = \mathbb{E}[\mu|\mathcal{D}] = \mu_{ML} = \text{sample mean} = \frac{m}{N} \quad (2.15)$$

ii.  $\text{var}[\mu|\mathcal{D}]$  is approaching zero (Eq 2.9):

In general, the posterior variance is smaller than the prior variance on average (not for every observation)

$$\mathbb{E}_{\mathcal{D}}[\text{var}_{\theta}[\theta|\mathcal{D}]] = \text{var}_{\theta}[\theta] - \text{var}_{\mathcal{D}}[\mathbb{E}_{\theta}[\theta|\mathcal{D}]] \quad (2.16)$$

## 2.2. Multinomial Variables

### 2.2.1. Multinomial Distribution

#### 2.1.1.4. Definition

$$\text{Mult}(m_1, m_2, \dots, m_K | \boldsymbol{\mu}, N) = \binom{N}{m_1, m_2, \dots, m_K} \prod_{k=1}^K \mu_k^{m_k} \quad (2.17)$$

$$m_k = \sum_n x_{nk}$$

It is the generalization of the Bernoulli distribution to more than two outcomes(states).

#### 2.1.1.5. Properties

$$\mathbb{E}[\mathbf{x}|\boldsymbol{\mu}] = \sum_{\mathbf{x}} p(\mathbf{x}|\boldsymbol{\mu})\mathbf{x} = (\mu_1, \dots, \mu_K)^T = \boldsymbol{\mu}$$

### 2.1.1.6. Density estimation

#### 2.1.1.6.1. Frequentist

$$\mathcal{D} = \{\mathbf{x}_1, \dots, \mathbf{x}_N\}$$

$$p(\mathcal{D}|\boldsymbol{\mu}) = \prod_{n=1}^N \prod_{k=1}^K \mu_k^{x_{nk}} = \prod_{k=1}^K \mu_k^{\sum_n x_{nk}} = \prod_{k=1}^K \mu_k^{m_k}$$

Sufficient statistic for this distribution:  $m_k = \sum_n x_{nk}$

$$\mu_k^{ML} = \frac{m_k}{N}$$

#### 2.1.1.6.2. Bayesian (Dirichlet distribution)

$$Dir(\boldsymbol{\mu}|\boldsymbol{\alpha}) = \frac{\Gamma(\alpha_0)}{\Gamma(\alpha_1) \dots \Gamma(\alpha_K)} \prod_{k=1}^K \mu_k^{\alpha_k - 1} \quad (2.18)$$

$$\alpha_0 = \sum_{k=1}^K \alpha_k$$

$$p(\boldsymbol{\mu}|\mathcal{D}, \boldsymbol{\alpha}) \propto \text{likelihood} \times \text{prior} = p(\mathcal{D}|\boldsymbol{\mu})p(\boldsymbol{\mu}|\boldsymbol{\alpha}) = Dir(\boldsymbol{\mu}|\boldsymbol{\alpha} + \mathbf{m})$$

$$= \frac{\Gamma(\alpha_0 + N)}{\Gamma(\alpha_1 + m_1) \dots \Gamma(\alpha_K + m_K)} \prod_{k=1}^K \mu_k^{\alpha_k + m_k - 1}$$

\*  $\alpha_k$  is an effective number of observations of  $x_k = 1$

## 2.3. Gaussian Distribution