

# Advanced Computational Statistics (Fall 2016)

J.P.Kim

Dept. of Statistics

Finally modified at September 8, 2016

# Preface & Disclaimer

This note is a summary of the lecture Advanced Computational Statistics (M1399.000200) held at Seoul National University, Fall 2016. Lecturer was Jung-Ho Won, and the note was summarized by J.P.Kim, who is a Ph.D student. There are few textbooks and references in this course, which are following.

- *Convex Optimization Theory, Dimitri P. Bertsekas, 2009.*
- *Optimization, Kenneth Lange, 2013.*
- *Convex Optimization, S.Boyd & L.Vandenberghe, 2004.*

Also I referred to following books when I write this note. The list would be updated continuously.

- 

If you want to correct typo or mistakes, please contact to: [joonpyokim@snu.ac.kr](mailto:joonpyokim@snu.ac.kr)

# Chapter 1

## Basic Concepts of Convex Analysis

### 1.1 Convex sets and functions

**Definition 1.1.1.** A set  $C \subseteq \mathbb{R}^n$  is **convex** if  $\alpha x + (1 - \alpha)y \in C$  for any  $x, y \in C$  and  $\alpha \in [0, 1]$ .

Note that  $\phi$  is convex by convention.

**Proposition 1.1.2.** Let  $C$  and  $C_i$  be convex sets for  $i \in I$ . Then,

(a)  $\bigcap_{i \in I} C_i$  is also a convex set.

(b)  $C_1 + C_2$  is a convex set.

(c) For any scalar  $\lambda$ ,  $\lambda C$  is a convex set. Also, for  $\lambda_1, \lambda_2 > 0$ ,  $(\lambda_1 + \lambda_2)C = \lambda_1 C + \lambda_2 C$  holds.

(d)  $cl(C)$  and  $int(C)$  are convex.

(e) For an affine function  $f$ ,  $f(C)$  or  $f^{-1}(C)$  is convex.

*Proof.* (c) Convexity is trivial. Let  $x \in (\lambda_1 + \lambda_2)C$ . Then for some  $y \in C$ ,  $x = (\lambda_1 + \lambda_2)y$  holds. Since  $\lambda_1 y \in \lambda_1 C$  and  $\lambda_2 y \in \lambda_2 C$ , we get  $x \in \lambda_1 C + \lambda_2 C$ . Thus we showed  $(\lambda_1 + \lambda_2)C \subseteq \lambda_1 C + \lambda_2 C$ .  $\supseteq$  part is similar.

(d) Let  $x, y \in cl(C)$ . Then  $\{x_k\}, \{y_k\} \subseteq C$  exist such that  $x_k \rightarrow x$  and  $y_k \rightarrow y$ . Note that for any  $\alpha \in [0, 1]$  we get  $\{\alpha x_k + (1 - \alpha)y_k\} \subseteq C$ , and so  $\alpha x + (1 - \alpha)y \in cl(C)$  from  $\alpha x_k + (1 - \alpha)y_k \rightarrow \alpha x + (1 - \alpha)y$ . Next, let  $x, y \in int(C)$ . Then there exists  $r > 0$  such that  $B(x, r) \subseteq C$  and  $B(y, r) \subseteq C$ . Note that  $B(x, r) = \{x + z : \|z\| < r\}$ . It's enough to show that  $B(\alpha x + (1 - \alpha)y, r) \subseteq C$ . Now  $B(\alpha x + (1 - \alpha)y, r) = \{\alpha x + (1 - \alpha)y + z : \|z\| < r\}$  and hence  $\alpha x + (1 - \alpha)y + z = \alpha \underbrace{(x + z)}_{\in B(x, r) \subseteq C} + (1 - \alpha) \underbrace{(y + z)}_{\in B(y, r) \subseteq C} \in C$  for any  $z$  such that  $\|z\| < r$ .

(e) If  $x, y \in f(C)$ ,  $\exists x', y' \in C$  such that  $x = f(x')$  and  $y = f(y')$ . Since  $f$  was affine, we get

$$\alpha x + (1 - \alpha)y = \alpha f(x') + (1 - \alpha)f(y') = f(\alpha x' + (1 - \alpha)y') \in f(C)$$

from  $\alpha x' + (1 - \alpha)y' \in C$ . Rest part is similar.  $\square$

**Example 1.1.3** (Special convex sets). In this example we see some examples of convex set.

(a) *Hyperplane*  $\{x : a^T x = b\}$  is convex, for given  $a$  and  $b$ .

(b) *Half-space*  $\{x : a^T x \leq b\}$  is also convex.

(c) *Polyhedra*,  $\{x : a_j^T x \leq b_j, a_j \neq 0, b_j \in \mathbb{R}, j = 1, 2, \dots, r\}$  is intersection of half-spaces, and hence convex.

(d)  $C$  is *cone* if  $\forall x \in C \lambda x \in C$  for any  $\lambda > 0$ . Note that, cone need not be convex, nor contain the origin. (See figure 1.1.) Rather, we consider *polyhedral cone*  $\{x : a_j^T x \leq 0, j = 1, 2, \dots, r\}$ , which contains the origin at the boundary. Polyhedral cone is convex.

(e)  $S = \{x : a^T x = 0\}$  is a convex set, subspace of  $\mathbb{R}^n$ , a hyperplane, and a polyhedral cone.




Figure 1.1: (a) Convex cone. (b) Convex cone which does not contain the origin. (c) Nonconvex cone, which consists of 2 lines.

**Definition 1.1.4.** Let  $C \subseteq \mathbb{R}^n$  be a convex set, and  $f : C \rightarrow \mathbb{R}$  be a function.  $f$  is **convex** if

$$f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y) \quad \forall x, y \in C, \alpha \in [0, 1].$$

*Remark:* Domain is a convex set! Also,  $f$  is **strictly convex** if

$$f(\alpha x + (1 - \alpha)y) < \alpha f(x) + (1 - \alpha)f(y) \quad \forall x, y \in C, x \neq y, \alpha \in (0, 1).$$

Finally,  $f$  is **concave** if  $-f$  is convex.

From now on, without mention,  $C$  always denote a convex set in  $\mathbb{R}^n$ .

**Example 1.1.5.** (a) Affine function  $f(x) = a^T x + b$  is both convex and concave.

(b) Any norm  $f(x) = \|x\|$  is convex from *triangle inequality*.

**Definition 1.1.6** (level set). Let  $f : C \rightarrow \mathbb{R}$  be a convex function. Then for any given  $\gamma \in \mathbb{R}$ ,

(a)  $\{x \in C : f(x) \leq \gamma\}$  is called **sublevel set** of  $f$ .

(b)  $\{x \in C : f(x) \geq \gamma\}$  is called **superlevel set** of  $f$ .

From now on, we will call a sublevel set as a **level set** in short.

**Remark 1.1.7.** It is known that if  $f$  is a convex function, then all of its level sets are convex.

Note that converse does not hold: See figure 1.2.




Figure 1.2: (a) Level set of convex function. (b) Even if all of level sets are convex, function need not be a convex one.

In many cases, it is convenient to allow function value be  $\pm\infty$ , or domain is  $\mathbb{R}^n$ . For this, we may consider a *extended real valued functions*. Then how to define convexity of such function  $f : C \rightarrow [-\infty, \infty]$ ? The rest part of this section handles this issue.

**Example 1.1.8.** (Motivation for extension to  $\bar{\mathbb{R}}$ )

(a) We may deal with the function  $f(x) = \sup_{i \in I} f_i(x)$ . Its value may be  $\infty$ .

(b) “Conjugate function” will be handled in section 1.6. To define this notion, extension should be required. For example, conjugate function  $f^*(y)$  of  $f(x) = |x|$  is

$$f^*(y) = \begin{cases} 0 & |y| \leq 1 \\ +\infty & o.w. \end{cases}.$$

(c) Consider  $f(x) = 1/x$  on  $(0, \infty)$ . For optimization, closed domain is useful and convenient, so we may extend the domain to  $[0, \infty]$ . In here,  $f(0) = \infty$  is reasonable extension.

**Remark 1.1.9.** Note that we can extend the domain of function  $f : C \rightarrow \mathbb{R}$  to  $\mathbb{R}^n$  as letting  $f(x) = \infty$  if  $x \notin C$ . Thus allowing function to be extended real-valued, we can extend the domain of function. Then how to restrict the origin domain again? *Effective domain*, which is following, can be one answer.

**Definition 1.1.10** (epigraph). **Epigraph** of function  $f : X \rightarrow \bar{\mathbb{R}}$  is defined as

$$\text{epi}(f) = \{(x, w) : x \in X, w \in \mathbb{R}, f(x) \leq w\}.$$

Note that  $w$  is not allowed to be  $\pm\infty$ .




Figure 1.3: Epigraph of a function

**Proposition 1.1.11.** *Epigraph of convex function is a convex set.*

*Proof.* Easy. □

**Definition 1.1.12** (effective domain). Let  $f : X \rightarrow \bar{\mathbb{R}}$  be a function. **Effective domain** of  $f$  is defined as

$$\text{dom}(f) = \{x \in X : f(x) < \infty\}.$$

There are some remarks.

**Remark 1.1.13.**

- (a) Since we usually deal with a convex function  $f$ , the point whose functional value is  $-\infty$  is out of interest.
- (b) Note that

$$\text{dom}(f) = \{x \in \mathbb{R}^n : \exists w \in \mathbb{R} \text{ s.t. } (x, w) \in \text{epi}(f)\},$$

so it is “projection of  $\text{epi}(f)$  onto  $\mathbb{R}^n$ . If we want to handle real valued function, we can think restriction on  $\text{dom}(f)$ . Or, as mentioned above, we can enlarge domain from  $X$  to  $\mathbb{R}^n$ . Extended or restricted functions have the same epigraph.

**Example 1.1.14.**

(a) Consider a function  $f : [0, \infty) \rightarrow [-\infty, \infty]$  defined as

$$f(x) = \begin{cases} \frac{1}{x} & x \neq 0 \\ +\infty & x = 0 \end{cases}.$$

Then  $\text{dom}(f) = (0, \infty)$ , and

$$\text{epi}(f) = \{(x, y) : 0 < x < \infty, y \geq 1/x.\}$$

(b) Suppose that  $f(x) = -\infty$  for some  $x \in X$ . Then its epigraph  $\text{epi}(f)$  may contain a vertical line.




Figure 1.4: Epigraph of nonproper function

**Definition 1.1.15** (proper function). Let  $f : X \rightarrow \bar{\mathbb{R}}$ .  $f$  is a **proper function** if

- (1)  $f(x) < \infty$  for at least one  $x \in X$ , and
- (2)  $f(x) > -\infty$  for all  $x \in X$ .

**Remark 1.1.16.** Note that,  $f$  is proper function  $\Leftrightarrow \text{epi}(f)$  is nonempty and does not contain vertical line.

Now we can extend the definition of convex function to *extended real valued function*.

**Definition 1.1.17.**  $f : C \rightarrow \bar{\mathbb{R}}$  is a convex function if  $\text{epi}(f)$  is a convex subset of  $\mathbb{R}^{n+1}$ .

**Remark 1.1.18.** Note that this definition satisfies followings.

- (1)  $\text{dom}(f)$  is convex.
- (2) All of level sets are convex.
- (3) If  $f(x) < \infty \forall x$  or  $f(x) > -\infty \forall x$ , it satisfies Jensen's inequality.

**Definition 1.1.19** (Indicator function). Let  $X \subseteq \mathbb{R}^n$  be a set. An **indicator function**  $\delta_X$  of  $X$  is defined as

$$\delta_X(x) = \begin{cases} 0 & x \in X \\ +\infty & \text{o.w.} \end{cases}.$$

Note that effective domain of  $\delta_X$  is  $X$ . Also, note that

$$X \text{ is (strictly) convex set} \Leftrightarrow \delta_X \text{ is (strictly) convex function.}$$

Also, if  $X \neq \emptyset$ ,  $\delta_X$  is proper.

**Remark 1.1.20.** Now we can give a correspondence between convex sets and convex functions. Epigraph of convex function is convex set, and indicator of convex set is convex function.




Figure 1.5: Correspondence between convex sets and functions

Now, we are ready for extension of convexity to nonconvex domain.

**Definition 1.1.21.** Let  $C$  be a convex set, and  $C \subseteq X \subseteq \mathbb{R}^n$ . Then  $f : X \rightarrow \bar{\mathbb{R}}$  is **convex over**  $C$  if  $f|_C : C \rightarrow \bar{\mathbb{R}}$  (restriction on  $C$  of  $f$ ) is convex function.

### 1.1.1 Closedness and Semicontinuity

**Definition 1.1.22.** A function  $f : X \rightarrow \bar{\mathbb{R}}$  is a closed function if its epigraph  $\text{epi}(f)$  is closed set.



It is reasonable definition because of correspondence between sets and functions. In Appendix A, we defined lower and upper semicontinuity of function. There is an important relationship between two notions of function. In fact, *they are equivalent*.

**Theorem 1.1.23.** *Let  $f : \mathbb{R}^n \rightarrow \bar{\mathbb{R}}$  be a function. Then TFAE<sup>1</sup>*

(i) *For any  $\gamma \in \mathbb{R}$ ,  $V_\gamma = \{x : f(x) \leq \gamma\}$  is closed.*

(ii)  *$f$  is lower semicontinuous function.*

(iii)  *$f$  is closed function. ( $\text{epi}(f)$  is closed)*

*Proof.* (i) $\Rightarrow$ (ii): Suppose that  $\bar{x}$  and a sequence  $\{x_k\}$  exist such that  $x_k \xrightarrow[k \rightarrow \infty]{} \bar{x}$  and  $f(\bar{x}) > \liminf_{k \rightarrow \infty} f(x_k)$ . Then  $\exists \gamma$  such that  $f(\bar{x}) > \gamma > \liminf_{k \rightarrow \infty} f(x_k)$ . Then there is a subsequence  $\{x_{k_j}\}$  which satisfies

$$f(x_{k_j}) \leq \gamma \quad \forall j,$$

by definition of  $\liminf$ . Hence  $V_\gamma := \{x : f(x) \leq \gamma\} \supseteq \{x_{k_j}\}$ , and from closedness of  $V_\gamma$ ,  $\bar{x} \in V_\gamma$  should be held, which yields contradiction.

(ii)  $\Rightarrow$  (iii): Choose a sequence  $\{(x_k, w_k)\} \subseteq \text{epi}(f)$  such that  $(x_k, w_k) \rightarrow (\bar{x}, \bar{w})$ . Then since  $f$  is l.s.c.,

$$f(\bar{x}) \leq \liminf_{k \rightarrow \infty} f(x_k) \leq w_k$$

holds by definition of epigraph. Thus we get

$$f(\bar{x}) \leq \bar{w}$$

by letting  $k \rightarrow \infty$ . Therefore  $(\bar{x}, \bar{w}) \in \text{epi}(f)$  holds.

(iii)  $\Rightarrow$  (i): Note that  $(x, \gamma) \in \text{epi}(f) \Leftrightarrow x \in V_\gamma$ . Let  $\gamma \in \mathbb{R}$ , and  $\{x_k\} \subseteq V_\gamma$  be a sequence converging to  $\bar{x}$ . Then  $(x_k, \gamma) \in \text{epi}(f)$  and  $(x_k, \gamma) \xrightarrow[k \rightarrow \infty]{} (\bar{x}, \gamma)$  hold, which imply  $(\bar{x}, \gamma) \in \text{epi}(f)$  since  $\text{epi}(f)$  is closed. Therefore  $\bar{x} \in V_\gamma$  is obtained.  $\square$

---

<sup>1</sup>The followings are equivalent.

# Appendix

## A Mathematical Background

In this section, we introduce some basic background used oftenly.

### A.1 Basic notions

- We often consider *extended real numbers*  $\bar{\mathbb{R}} := \mathbb{R} \cup \{-\infty, \infty\}$ . Also we define
  - $x \cdot 0 = 0 \ \forall x \in \bar{\mathbb{R}}$
  - $x \cdot \infty = \infty$  if  $x > 0$
  - $x \cdot \infty = -\infty$  if  $x < 0$
  - $x + \infty = \infty, \ x - \infty = -\infty \ \forall x \in \mathbb{R},$
  - and we do not allow  $\infty - \infty$ .
- For nonempty subset  $X$  of  $\mathbb{R}$ , we define  $\sup X$  as the smallest  $y \in \mathbb{R}$  such that  $y \geq x$  for any  $x \in X$ , and if such  $y$  does not exist, we define  $\sup X = \infty$ . Also, we define  $\sup \emptyset = -\infty$ . We can define  $\inf X$  similarly.
- If  $\sup X := \bar{x}$  is contained in  $X$ , we say that  $\bar{x} = \max X$ . (“maximum is attained”)  
If  $\inf X := \bar{x}$  is contained in  $X$ , we say that  $\bar{x} = \min X$ . (“minimum is attained”)
- Vector space. In this course, we only consider  $\mathbb{R}^n$ . In here, inner product  $\langle x, y \rangle = x^T y$  is defined.
- Also, for  $x \in \mathbb{R}^n$ , define the notation  $x > 0$  or  $x \geq 0$  componentwisely. Also define  $x > y \Leftrightarrow x - y > 0$ .
- Let  $f : X \rightarrow Y$  be a function. For  $U \subseteq X$  and  $V \subseteq Y$ , we define
  - $f(U) := \{f(x) : x \in U\}$  (“image of  $U$ ”)

$$- f^{-1}(V) := \{x \in X : f(x) \in V\} \quad (\text{"inverse image of } V\text{"})$$

## A.2 Linear Algebra

- Let  $X, X_1, X_2 \subseteq \mathbb{R}^n$  and  $\lambda$  be a scalar. we define

$$- \lambda X := \{\lambda x : x \in X\}$$

$$- X_1 + X_2 := \{x_1 + x_2 : x_1 \in X_1, x_2 \in X_2\}$$

$$- \bar{x} + X := \{\bar{x}\} + X \text{ for } \bar{x} \in \mathbb{R}$$

$$- \text{and } X_1 - X_2 = \{x_1 - x_2 : x_1 \in X_1, x_2 \in X_2\}.$$

$$- \text{To prevent abuse of notation, we will use } X_1 \setminus X_2 \text{ for "set difference."}$$

- If  $X_i \subseteq \mathbb{R}^{n_i}$ ,  $i = 1, 2, \dots, m$ , we define "Cartesian Product" as

$$X_1 \times \dots \times X_m := \{(x_1, \dots, x_m) : x_i \in X_i, i = 1, 2, \dots, m\} \subseteq \mathbb{R}^{n_1 + \dots + n_m}.$$

- $S \subseteq \mathbb{R}^n$  is called subspace if  $ax + by \in S$  for any  $x, y \in S$  and  $a, b \in \mathbb{R}$ .

- Also, for  $\bar{x} \in \mathbb{R}$ ,  $X := \bar{x} + S$  is called an affine set, if  $S$  is a subspace.




Figure 6: Affine set and subspace

- Facts:

1.  $\exists$  unique subspace associate with an affine set.
2.  $X (\neq \phi)$  is a subspace *if and only if*  $0 \in X$  and  $\alpha x + (1 - \alpha)y \in X$  for any  $\alpha \in \mathbb{R}$  and  $x, y \in S$ .
3.  $X (\neq \phi)$  is an affine set *if and only if*  $\alpha x + (1 - \alpha)y \in X$  for any  $\alpha \in \mathbb{R}$  and  $x, y \in S$ .
4. Note that intersection of subspaces is also a subspace.

- $\text{span}(x_1, \dots, x_m)$  is a subspace generated by  $x_1, \dots, x_m$ , and it is a set of linear combinations.
- We say that  $x_1, \dots, x_m$  are linearly independent if  $\nexists(\alpha_1, \dots, \alpha_m) \neq 0$  such that  $\sum_{k=1}^m \alpha_k x_k = 0$ .
- Let  $S$  be a nontrivial subspace. Then  $\{x_1, \dots, x_m\}$  is a basis for  $S$  if  $x_1, \dots, x_m \in S$ ,  $\text{span}(x_1, \dots, x_m) = S$  and they are linearly independent. In this case, we say  $\dim S = m$ . Also we define  $\dim(\{0\}) = 0$ .
- Dimension of the affine set is defined as that of associated subspace. In other words,  $\dim(\bar{x} + S) = \dim S$ .
- For given  $a$  and  $b$ , we define  $\{x \in \mathbb{R}^n : a^T x = b\}$  as a hyperplane.
- Let  $X \subseteq \mathbb{R}^n$ . Then  $X^\perp := \{y : \langle y, x \rangle = 0 \ \forall x \in X\}$  is a subspace of  $\mathbb{R}^n$ . In particular, if  $S$  is a subspace, then  $S^\perp$  is an orthogonal complement of  $S$ . We can say that  $\mathbb{R}^n = S \oplus S^\perp$ , and  $(S^\perp)^\perp = S$ .
- Matrices. Let  $A \in \mathbb{R}^{m \times n}$ . Then we define  $AX := \{Ax : x \in X\}$  and  $A^{-1}Y := \{x : Ax \in Y\}$ .
- Let  $\mathbb{S}^n = \{A \in \mathbb{R}^{n \times n} : A^T = A\}$ . Then positive definite matrices are elements of the set

$$\mathbb{S}_{++}^n := \{A \in \mathbb{S}^n : x^T A x > 0 \ \forall x \in \mathbb{R}^n \setminus \{0\}\},$$

and denote as  $A \succ 0$  if  $A$  is s.p.d.. Also, we define a set of nonnegative definite matrices

$$\mathbb{S}_+^n := \{A \in \mathbb{S}^n : x^T A x \geq 0 \ \forall x \in \mathbb{R}^n \setminus \{0\}\},$$

and denote as  $A \succeq 0$  if  $A \in \mathbb{S}_+^n$ .

- If  $A \succeq 0$ , then there exists  $M$  such that  $A = M^T M$ .
- For a matrix  $A \in \mathbb{R}^{m \times n}$ , we define range and null space of  $A$  as

$$\mathcal{R}(A) = \{Ax : x \in \mathbb{R}^n\}$$

$$\mathcal{N}(A) = \{x : Ax = 0\}$$

respectively.

- Rank of matrix  $A$  is defined as  $\text{rank}(A) = \dim(\mathcal{R}(A))$ . Note that,  $\text{rank}(A) = \text{rank}(A^T)$ , and  $\mathcal{R}(A) = (\mathcal{N}(A^T))^\perp$ .
- If  $\text{rank}(A) = m \wedge n$  we say that  $A$  is of full rank.

### A.3 Basic Topology

- In here we often use the Euclidean norm  $\|x\| = \sqrt{x^T x}$ . Then, Cauchy Schwarz inequality  $|x^T y| \leq \|x\| \cdot \|y\|$  and Pythagoras theorem  $\langle x, y \rangle = 0 \Rightarrow \|x + y\|^2 = \|x\|^2 + \|y\|^2$  are known.
- Let  $\{x_k\}$  be a sequence in  $\mathbb{R}$ . We say  $\{x_k\}$  converges if  $\exists x \in \mathbb{R}$  such that  $\forall \epsilon > 0 \exists K$  s.t.  $\forall k \geq K \Rightarrow |x_k - x| < \epsilon$ .
- Also we say that  $x_k$  diverges to  $\infty$  if  $\forall b \in \mathbb{R} \exists K$  s.t.  $\forall k \geq K \ x_k \geq b$ .
- $\{x_k\}$  is bounded above if  $\exists b$  such that  $x_k \leq b$  for any  $k$ .
- We can define

$$\limsup_{k \rightarrow \infty} x_k := \inf_{m \geq 1} \sup_{k \geq m} x_k = \lim_{m \rightarrow \infty} \sup_{k \geq m} x_k$$

$$\liminf_{k \rightarrow \infty} x_k := \sup_{m \geq 1} \inf_{k \geq m} x_k = \lim_{m \rightarrow \infty} \inf_{k \geq m} x_k.$$

- Note that

$$\inf_{k \geq 1} x_k \leq \liminf_{k \rightarrow \infty} x_k \leq \limsup_{k \rightarrow \infty} x_k \leq \sup_{k \geq 1} x_k$$

holds.

- Also, if for any  $k \ x_k \leq y_k$  holds, then  $\liminf x_k \leq \liminf y_k$  and  $\limsup x_k \leq \limsup y_k$ .
- Moreover,

$$\liminf_{k \rightarrow \infty} x_k + \liminf_{k \rightarrow \infty} y_k \leq \liminf_{k \rightarrow \infty} (x_k + y_k)$$

$$\limsup_{k \rightarrow \infty} x_k + \limsup_{k \rightarrow \infty} y_k \geq \limsup_{k \rightarrow \infty} (x_k + y_k)$$

hold.

- In general, for  $\{x_k\} \subseteq \mathbb{R}^n$ , we define  $x_k \rightarrow x$  as  $k \rightarrow \infty$  if  $x_{ki} \rightarrow x_i$  as  $k \rightarrow \infty$ . (componentwisely)
- Now we consider a subsequence  $\{x_k : k \in \mathcal{K}\}$ .  $x$  is called limit point if there exists a subsequence such that converges to  $x$ .

- Then we get following **Bolzano-Weierstrass Theorem**, *every bounded sequence has at least one limit point.*
- We can define closure  $cl(X)$  and interior  $int(X)$  of  $X$ . Also we can define boundary  $bd(X) := cl(X) \setminus int(X)$  of  $X$ .
- Facts:
  - The union of a finite collection of closed sets is closed.
  - The intersection of any collection of closed sets is closed.
  - The union of any collection of open sets is open.
  - The intersection of a finite collection of open sets is open.
  - A set is open if and only if all of its elements are interior points.
  - Every subspace of  $\mathbb{R}^n$  is closed.
  - A set  $X$  is compact if and only if every sequence of elements of  $X$  has a subsequence that converges to an element of  $X$ .
  - (“Cantor’s intersection theorem”, or if underlying space is  $\mathbb{R}$ , “Nested interval theorem”) *If  $\{X_k\}$  is a sequence of nonempty and compact sets such that  $X_{k+1} \subset X_k$  for all  $k$ , then the intersection  $\bigcap_{k=0}^{\infty} X_k$  is nonempty and compact.*
- Continuity. A function  $f : X \rightarrow \mathbb{R}^n$  is continuous at  $x$  if for any sequence  $\{x_k\}$  converges to  $x$ ,  $\lim_k f(x_k) = f(x)$  holds.
- A function  $f : X \rightarrow \mathbb{R}^n$  is right-continuous (left-continuous) at  $x$  if for any sequence  $\{x_k\}$  converges to  $x$  satisfying  $x_k > x$  ( $x_k < x$ ),  $\lim_k f(x_k) = f(x)$  holds.
- A real-valued function  $f : X \rightarrow \mathbb{R}$  is upper semicontinuous (lower semicontinuous) at  $x \in X$  if  $f(x) \geq \limsup f(x_k)$  ( $f(x) \leq \liminf f(x_k)$ ) for any sequence  $\{x_k\}$  in  $X$  that converges to  $x$ .

- For example, function

$$f(x) = \begin{cases} \sin(1/x) & x \neq 0 \\ 1 & x = 0 \end{cases}$$

is upper semicontinuous (Figure 7). For more examples, see figure 8.

- Facts:




Figure 7: The graph of  $y = \sin(1/x)$ . Image from WolframAlpha.




Figure 8: (Left) Such function is upper semicontinuous. (Right) Such function is lower semicontinuous.

- Any vector norm on  $\mathbb{R}^n$  is a continuous function.
- Let  $f : \mathbb{R}^m \rightarrow \mathbb{R}^p$  and  $g : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be continuous functions. Then  $f \circ g$  is also continuous.
- Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be continuous, and  $Y$  be an open (closed) subset of  $\mathbb{R}^m$ . Then  $f^{-1}(Y)$  is open (closed).
- Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be continuous, and  $X$  be a compact subset of  $\mathbb{R}^n$ . Then  $f(X)$  is compact.
- Following is **Weierstrass' theorem**, or **max-min theorem**: A continuous function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  attains a minimum over any compact subset of  $\mathbb{R}^n$ .

*Proof.* Let  $X \subseteq \mathbb{R}^n$  be a compact set. Define a level set  $V_\gamma = \{x \in X : f(x) \leq \gamma\}$ , then it is compact since it is bounded and closed. Let  $f^* := \inf_{x \in X} f(x) < \infty$ . Then for a sequence  $\{\gamma_k\}$  such that  $\gamma_k \searrow f^*$  and  $\gamma_k > f^*$ ,  $V_{\gamma_k}$  is nonempty, so by Cantor's intersection theorem,  $\cap_k V_{\gamma_k}$  is nonempty compact set. Thus,  $X^* := \{x \in X : f(x) = f^*\} = \cap_k V_{\gamma_k}$  is nonempty.  $\square$