CPSC-354 Report

Junho Yi Chapman University

September 2, 2025

Abstract

Contents

L	Introduction	1
2	Week by Week 2.1 Week 1 2.1.1 Notes and Exploration 2.1.2 Homework 1	1
3	Essay	2
1	Evidence of Participation	2
5	Conclusion	2

1 Introduction

2 Week by Week

2.1 Week 1

2.1.1 Notes and Exploration

Place holder for notes

In abstract rewriting, an object is in normal form if it cannot be rewritten any further, i.e. it is irreducible Confluence system: in a system the result eventually converges into the same answer.

Termination: Means the system stops at some point.

Decidability:

church turing thesis:

Abstract rewriting system(ARS): mathematically the same as a directed graph A is a set of "strings" (can be anything) R is the relation

so in the MIU puzzle, A is M I U, the strings we use then R is the rules we are given. ie: (Mx,Mxx)|x e A U...

2.1.2 Homework 1

This week's HW is regarding the MU puzzle, and its relevance and application to formal systems. Here we use the MU puzzle to practice and familiarize ourselves with staying within the confines of a formal system. We are given 4 rules/restrictions, which is referred to as the "Requirement of Formality". Our formal system consists of these 4 rules:

- 1. RULE I: If you possess a string whose last letter is I, you can add on a U at the end.
- 2. RULE II: Suppose you have Mx. Then you may add Mxx to your collection.
- 3. RULE III: If III occurs in one of the strings in your collection, you may make a new string with U in place of III.
- 4. RULE IV: If UU occurs inside one of your strings, you can drop it.

With these four rules in mind, we have one objective: stay within the rules and produce "MU" from "MI". As I worked through the rules, I logically deduced these points in this order:

- 1. When applying RULE II, if I exists somewhere on the string, the parity of I becomes even until RULE III is applied again.
- 2. When applying RULE III, the I's (which are even, if RULE III applies), swaps parity, i.e. it goes from even to odd.
- 3. The lowest continuous string of I's where RULE III can be applied is four I's.
- 4. Because RULE III is the only way to reduce the number of I's, and its only possible to apply RULE III if there is a minimum of four continuous I's (due to RULE II) and an even parity of I's, using RULE III to reduce the amount of I's will always result in a remainder (leftover I).
- 5. Therefore, you can never get rid of I's fully with RULE III, or any other RULE usable by us without modifications of the rules.

From the above observations, we can see that there is no way to completely reduce the number of I's into zero with the given rules. This is my personal analysis of the MU puzzle, below is the "correct" analysis of the MU puzzle

Proof (invariant mod 3). Let n be the number of I's in the current string. Then:

```
Rule I: n \mapsto n, Rule II: n \mapsto 2n, Rule III: n \mapsto n-3, Rule IV: n \mapsto n.
```

Hence $n \mod 3$ is preserved by Rules I, III, IV, and toggles between 1 and 2 under Rule II. Initially, MI has $n=1\equiv 1\pmod 3$. No sequence of the above operations can yield $n\equiv 0\pmod 3$. But MU has 0 I's, i.e. $n=0\equiv 0\pmod 3$. Therefore MU is not derivable from MI.

- 3 Essay
- 4 Evidence of Participation
- 5 Conclusion

References

[BLA] Author, Title, Publisher, Year.