

Structural and functional MRI data differentially predict chronological age and behavioral memory performance

Joram Soch^{*,1,2,•}, Anni Richter^{*,3}, Hartmut Schütze^{4,5}, Jasmin Kizilirmak¹, Björn Schott^{1,3,6,7,•}

¹ German Center for Neurodegenerative Diseases (DZNE), Göttingen, Germany

² Bernstein Center for Computational Neuroscience (BCCN), Berlin, Germany

³ Leibniz Institute for Neurobiology (LIN), Magdeburg, Germany

⁴ German Center for Neurodegenerative Diseases (DZNE), Magdeburg, Germany


⁵ Medical Faculty, Otto von Guericke University, Magdeburg, Germany

⁶ Center for Behavioral Brain Sciences (CBBS), Magdeburg, Germany


⁷ Department of Psychiatry and Psychotherapy, University Medical Center Göttingen



We predicted chronological age and memory performance from a number of source variables / feature sets.





Age group can be classified based on all these variables.



Chronological age is best predicted from structural MRI, but memory performance is best predicted from functional MRI.



B

chronological age




C


memory performance



In order to follow up, we partitioned older subjects based on chronological age and memory performance.



There is a double dissociation between memory vs. age and functional MRI vs. structural MRI (& rs-fMRI)




Summary

- Chronological age is best predicted from structural MRI, but memory performance is best predicted from functional MRI.
- Single-value fMRI scores outperform whole-brain fMRI contrasts in predicting (independent) memory performance.


THANK YOU!
QUESTIONS?

Joram.Soch@DZNE.de



Appendix

Single-value fMRI scores outperform whole-brain fMRI contrasts in predicting independent memory performance.



The predictive utility of fMRI scores for memory performance is still moderate.

