

Review: Sporadic Parkinson's disease: development and distribution of α -synuclein pathology

K. Del Tredici and H. Braak

Clinical Neuroanatomy Section, Department of Neurology, Center for Biomedical Research, University of Ulm, Ulm, Germany

K. Del Tredici and H. Braak (2016) *Neuropathology and Applied Neurobiology* 42, 33–50

Sporadic Parkinson's disease: development and distribution of α -synuclein pathology

The development of α -synuclein immunoreactive aggregates in selectively vulnerable neuronal types of the human central, peripheral, and enteric nervous systems is crucial for the pathogenesis of sporadic Parkinson's disease. The presence of these lesions persists into the end phase of the disease, a process that is not subject to remission. The initial induction of α -synuclein misfolding and subsequent aggregation probably occurs in the olfactory bulb and/or the enteric nervous system. Each of these sites is exposed to potentially hostile environmental factors. **Once formed, the aggregates appear to be capable of propagating trans-synaptically from nerve cell to nerve cell in a virtually self-promoting pathological process.** A regional distribution pattern of aggregated α -synuclein emerges that entails the involvement of only a few types of susceptible and axonally interconnected projection neurons within the human nervous system. One major route of disease progression may originate in the

enteric nervous system and retrogradely reach the dorsal motor nucleus of the vagal nerve in the lower brainstem. From there, the disease process proceeds chiefly in a caudo-rostral direction through visceromotor and somatomotor brainstem centres to the midbrain, forebrain, and cerebral cortex. Spinal cord centres may become involved by means of descending projections from involved lower brainstem nuclei as well as by sympathetic projections connecting the enteric nervous system with postganglionic peripheral ganglia and preganglionic nuclei of the spinal cord. The development of experimental cellular and animal models is helping to explain the mechanisms of how abnormal α -synuclein can undergo aggregation and how transmission along axonal connectivities can occur, thereby encouraging the initiation of potential disease-modifying therapeutic strategies for sporadic Parkinson's disease.

Keywords: α -synuclein, autophagy, central nervous system, chaperones, dorsal motor nucleus of the vagal nerve, enteric nervous system, Lewy pathology, Parkinson's disease, peripheral nervous system, protein aggregation/misfolding

Introduction

Sporadic Parkinson's disease (PD) is a widespread human disease [1–4] that has not been reported in

nonhuman vertebrates. It involves circumscribed portions of the central, peripheral, and enteric nervous systems (CNS, PNS and ENS) (Table 1; Figure 1) [5–13]. A protracted but progressive and irreversible pathological process that usually manifests itself clinically in late midlife and lasts for decades accompanies the disease (Table 2).

Despite the more obvious motor symptoms (initially asymmetric hypokinesia, cogwheel rigidity, resting

Correspondence: Kelly Del Tredici, Clinical Neuroanatomy (Department of Neurology), Center for Biomedical Research, University of Ulm, Helmholtzstrasse 8/1, 89081 Ulm, Germany. Tel: +49 731 500 63114; Fax: +49 731 500 63133; E-mail: kelly.del-tredici@uni-ulm.de

Table 1. Distribution of Lewy pathology in the enteric, peripheral, and central nervous systems together with clinical symptoms

Stage 1	Initial development of LNs somewhere in ENS, PNS or CNS
	LNs and LBs in olfactory bulb and anterior olfactory nucleus
	LNs and LBs in dorsal motor nucleus of the vagal nerve, intermediate reticular zone
	Absence of symptoms; then a phase of prodromal symptoms, for example hyposmia, autonomic dysfunction
Stage 2	LNs in ENS, in peripheral parasympathetic and sympathetic nerves, and in peripheral autonomic ganglia
	LNs and LBs in medullary nuclei of the level setting system, for example lower raphe nuclei, locus coeruleus
	Hyposmia, autonomic dysfunction, for example gastrointestinal, urinary symptoms
	Disturbed sleep, parasomnias, mood changes
Stage 3	LNs and LBs in tegmental pedunculopontine nucleus and substantia nigra, pars compacta
	LNs in spinal cord centres reached by descending projections of level setting nuclei
	LNs and LBs in upper raphe nuclei, magnocellular nuclei of the basal forebrain, hypothalamic tuberomammillary nucleus
	LNs and LBs in central subnucleus of the amygdala
Stage 4	Disturbed sleep and possible early phase motor dysfunction: asymmetric tremor, rigidity, hypokinesia
	LNs and LBs in midline and intralaminar nuclei of the thalamus
	LNs and LBs in anteromedial temporal cortex (transentorhinal and entorhinal regions, hippocampal formation, plexus of LNs in the second sector of the Ammon's horn)
	Early phase motor dysfunction: tremor, rigidity, hypokinesia
Stage 5	LNs and LBs in superordinate cortical areas for regulation of autonomic functions
	LNs and LBs in high-order sensory association areas and prefrontal fields
	Late phase motor disability: fluctuation, falls, wheelchair bound or bedridden
	Cognitive impairment
Stage 6	LNs and LBs in first-order sensory association areas and premotor fields
	LNs and LBs in primary sensory and primary motor areas
	Late phase motor disability: fluctuation, wheelchair bound or bedridden
	Cognitive impairment, dementia

tremor, postural instability) that are associated with clinical phases of the disease, an accurate interpretation of the early prodromal dysfunctions [14, 15] and the differential diagnostics of PD as opposed to other forms of parkinsonism are complicated [16]. For this reason, autopsy-based assessment is required to confirm

the clinical diagnosis and to identify individuals with early PD-related neuropathology in the broader population [17–20].

α -Synuclein and Lewy pathology

Sporadic PD still is defined neuropathologically by the continuous formation of α -synuclein immunoreactive inclusion bodies [21–28], which develop only in a few especially susceptible neuronal types within circumscribed portions of the CNS, PNS, and ENS [29–32]. The inclusions occur in the form of spherical Lewy bodies (LBs) in cell somata [33–36], as elongated spindle-shaped or thread-like Lewy neurites (LNs) in axons and dendrites [37–40], as pale bodies [25], and sometimes as particulate (granular, dot-like, punctate) aggregates [41, 42]. These lesions, subsumed under the rubric 'Lewy pathology' (LP), very likely occupy a central role in the pathogenesis of the disease [34, 35, 43–49]. Individuals who lack LP at autopsy, but who displayed symptoms resembling those seen in PD, have atypical parkinsonian disorders other than sporadic PD [50].

The natively unfolded protein α -synuclein is present in erythrocytes and platelets [51, 52] but is a predominantly neuronal protein that occurs in most, but not all, types of mature human nerve cells [53]. It is produced in the nerve cell soma and, provided it is fully functional, is redistributed from there into the axonal compartment and ultimately resides in presynaptic terminals, where it may contribute to neurotransmission and synaptic homeostasis [54–60]. The presence of a hydrophobic NAC region within the α -synuclein molecule distinguishes it from related β - and γ -synucleins, confers it with a β -sheet potential, and is required for abnormal conformational changes that can induce the process of aggregation [61–63]. It has been proposed that, in normal nerve cells, natively monomeric α -synuclein exists in dynamic equilibrium with helical tetramers that appear to be more resistant to aggregation than natively unfolded monomers [64–67], but the evidence for a predominantly tetrameric state has been challenged [68–70]. Specific types of chaperones are present in the preterminal axoplasm and appear to regulate the functional relationships between tetrameric α -synuclein and synaptic vesicle membranes [65].

Thread-like LNs generally are the first lesions that are detectable in involved regions [31, 71–74]. In many instances, it is unclear whether the filamentous




Figure 1. Diagram showing possible routes for the transmission of α -synuclein pathology within the human nervous system between stages 1 and 6 of the PD-associated pathological process. Four degrees of blue background shading indicate the enteric and peripheral nervous systems (dark blue), portions of the lower brainstem and spinal cord (medium blue), structures in the upper brainstem, thalamus and amygdala (light blue) and the cerebral cortex (sky blue). Initial predilection sites during stage 1 are the olfactory bulb (black), anterior olfactory nucleus and dorsal motor nucleus of the vagal nerve (black). In the brain, the disease process progresses in a chiefly caudo-rostral trajectory. At stage 2, lesions are seen for the first time in the level setting nuclei: the locus coeruleus, magnocellular nuclei of the reticular formation, and lower raphe nuclei (violet). A bifurcation of the pathological process may occur during stage 3 (brown). The pathological process reaches centres of the central autonomic network (central subnucleus of the amygdala, components of the limbic circuit) and progresses into centres of the somatomotor system (pedunculopontine tegmental nucleus and pars compacta of the substantia nigra). The cerebral cortex first becomes involved at stage 4, beginning with the temporal transentorhinal region (dark red) and reaches, thereafter, the entorhinal region and the Ammon's horn (dark red). The higher order association areas of the neocortex become involved in stage 5 (red), followed by the first-order association areas and primary fields in stage 6 (light red). The spinal cord becomes involved during stage 2 possibly via descending projections from involved lower brainstem nuclei (violet) as well as by sympathetic projections connecting the enteric nervous system (black) with postganglionic peripheral ganglia (violet) and spinal cord preganglionic nuclei (violet). Prim. sensory fields, primary sensory fields; first-order assoc. fields, first-order association fields; olf. bulb, olfactory bulb; v. striatum, ventral striatum; v. pallidum, ventral pallidum; MD, mediodorsal nuclei of the thalamus; cereb. circuit, cerebellar circuit; ped. pont. nucl., pedunculopontine tegmental nucleus; parasymp. pregangl. neurons, parasympathetic preganglionic neurons of the dorsal motor nucleus of the vagal nerve; IML, intermediolateral nucleus; SPS, sacral parasympathetic nucleus; symp. postgangl. neurons, sympathetic postganglionic neurons; ENS, enteric nervous system.

Table 2. Nonselected autopsy cases ($n = 2366$) showing sporadic PD stages in different age groups by decade

	<i>LP</i>	<i>Stage</i>						
<i>Age</i>	<i>O</i>	<i>1</i>	<i>2</i>	<i>3</i>	<i>4</i>	<i>5</i>	<i>6</i>	<i>Totals</i>
0–9	7	0	0	0	0	0	0	0
10–19	22	0	0	0	0	0	0	0
20–29	66	0	0	0	0	0	0	0
30–39	95	0	0	0	0	0	0	0
40–49	170	0	0	0	0	0	0	0
50–59	316	3	4	2	1	0	0	10
60–69	465	5	2	8	2	5	0	22
70–79	498	11	10	15	17	12	1	66
80–89	450	10	13	14	14	16	8	75
90–100	90	4	0	4	3	3	0	14
Totals	2179	33	29	43	37	36	9	187

LP, Lewy pathology.

lesions are dendritic or axonal inclusions; however, inasmuch as in mature neurons α -synuclein is localized chiefly in the axon, it is possible that the initial thread-like lesions preferentially develop in preterminal portions of the axonal compartment [75], thereby leaving uninvolved only the presynaptic terminals and the initial axon segment [31]. This means that the protein would have to undergo a slight relocation prior to its aggregation within the axon. The very first small aggregates may possibly be transported in an anterograde as well as retrograde direction but rapidly consolidate to form larger LNs that remain stationary in the axon.

Shortly after the appearance of axonal LNs, additional aggregates develop and rapidly accumulate in the *somatodendritic compartments* of involved neurons. The sources of this abnormal material are unclear. The initial aggregation process most likely results in the reduction of monomeric α -synuclein in the axon, a state that could rapidly induce renewed production of monomers in the soma, which then – at the wrong location – form oligomers, insoluble protofibrils and, subsequently, insoluble fibrillar somatic LBs and dendritic LNs [58, 76–78]. In addition to α -synuclein, the aggregates can contain a variety of other components, including ubiquitin, tyrosine hydroxylase, synaptophysin, heat shock and autophagosomal proteins [79–86].

The insoluble intra-axonal and cytoplasmic aggregates possess amyloid characteristics [87]. More importantly, they are resistant to proteasomal recycling,

autophagy or other endogenous cellular removal mechanisms [86, 88–94]. The larger spindle-shaped axonal aggregates, which can also be connected to each other, are distributed across extensive portions of the axon, and do not shift in an anterograde or retrograde direction [40, 73, 74, 95–97]. These spindles likely reduce normal axonal transport functions [73, 96, 98, 99].

Postmitotic neurons suspend their cell cycle for purposes of survival [100] and possess numerous surveillance mechanisms for repairing transient damage and eliminating defective proteins [101, 102]. For these reasons, it is unlikely that such cells would ‘permit’ spontaneous aggregation of α -synuclein molecules that subsequently cannot be degraded or exocytosed. Indeed, spontaneous aggregation of α -synuclein barely occurs under optimal experimental conditions, where amyloid-like fibrils develop only after prolonged incubation [56, 69, 103]. Moreover, were the natively unfolded protein to be present in tetrameric form, its tendency to resist aggregation would be, presumably, even greater. Because there are major differences between the axonal and somatodendritic compartments, the mechanisms by which axonal compartments deal with misfolded and aggregated proteins merit closer study.

Just as there are mechanisms to repair damage in postmitotic nerve cells or to stabilize partially folded protein intermediates [104, 105], mechanisms might also exist that support, and possibly accelerate, the processes leading to the production of abnormal α -synuclein aggregates. Instead of guardian chaperones, for instance, endogenous or exogenous proteins could ‘coax’ or ‘mislead’ the protein α -synuclein into undergoing conformational changes and then building pathological aggregates [106]. Such putative molecules possessing what we would call ‘Siren-like’ functions may waylay the host nerve cells during an assault on axons and thereby annul the otherwise unlikely scenario of spontaneous aggregation [107].

Specific types of nerve cells develop PD-associated Lewy pathology during the prodromal and clinical disease phases

As pointed out previously, very few types of nerve cells within the entire human nervous system are capable of developing LP, whereas other neuronal types, including directly adjacent nerve cells, remain intact for the

duration of the illness. Only projection cells with a disproportionately long, thin, and weakly myelinated or nonmyelinated axon become involved; by contrast, interneurons and projection cells with short axons or those with a heavily myelinated axon are not susceptible [30, 31, 108]. The specific regional distribution of the lesions within portions of the nervous system is a mirror of this selective vulnerability.

Because LP formation is not limited to dopaminergic projection neurons of the substantia nigra but also occurs in glutamatergic, noradrenergic, serotonergic, histaminergic and cholinergic projection cells, neurotransmitters *per se* are not adequate predictors of which neurons are predisposed or resistant to the pathological process. During the past decade, PD has come to be recognized as a multisystem disorder rather than a chiefly monosystemic disorder with selective dysfunction and loss of nigral neurons [11, 12, 20, 109–113].

The combination of especially susceptible neuronal types and the peculiarities of their distribution pattern make it possible not only to differentiate sporadic PD from other synucleinopathies but also to recognize mild forms of the pathological process in nonsymptomatic individuals. Those lacking clinically manifest motor symptoms but displaying LP in the same neuronal types and at the same sites as PD patients most likely represent individuals in a prodromal phase of PD [19, 112, 114–130].

Cases in an early phase are exceptionally valuable because they are temporally close to the prevailing circumstances at the outset of the disease process. Thus, evaluation of this phase is crucial to all other efforts directed at replacing currently symptomatic treatments with potentially causally effective therapies [131–135].

The pathological process underlying PD consists of a very early nonsymptomatic period, followed by a prodromal phase, often characterized by olfactory dysfunction [118, 136–144], autonomic dysregulation [145–156], as well as sleep, and/or mood disturbances [134, 157–159]. This phase transitions into a phase that is accompanied by the classical somatomotor symptoms and impaired cognitive functioning (Table 1) [2, 139, 159–166]. The duration of each phase varies considerably from one individual to another [134, 139, 167–169]; nevertheless, by the time of the initial PD diagnosis, when patients have crossed the threshold to clinically manifest somatomotor disease, the pathological

process within the nervous system is already remarkably advanced (Table 1).

Interindividual differences with respect to disposition, pre-existing comorbidities, and environmental influences most probably have an impact on the threshold at which the conversion to the final phase occurs as well as on the reported phenotypical heterogeneity of initially presenting symptoms, disease course, and rates of disease progression [169–172]. Grading according to six neuropathological stages can incorporate all three phases, provided the assumption is correct that non-symptomatic, prodromal, and symptomatic motor phases combined represent a biological continuum [19, 20, 31, 71, 112, 134, 171, 173–177].

Regional distribution pattern of the α -synuclein lesions within the nervous system in PD

Within the brain, the nonrandom development of the pathological process generally progresses caudo-rostrally through susceptible regions of the lower brain-stem (the dorsal motor nucleus of the vagal nerve, lower raphe nuclei, magnocellular nuclei of the reticular formation, locus coeruleus) into midbrain tegmental nuclei (pedunculopontine nucleus, dopaminergic projection neurons of the substantia nigra, nuclei of the upper raphe system) and noncortical centres of the forebrain (amygdala, hypothalamic tuberomammillary nucleus, magnocellular nuclei of the basal forebrain, midline and intralaminar nuclei of the thalamus). Then, it reaches the cerebral cortex (transentorhinal and entorhinal regions, Ammon's horn, cortical visceromotor areas, and finally the entire neocortex) (Figure 1) [47, 71, 134, 173, 176–178].

Cases with the earliest signs of α -synuclein immunoreactivity frequently display LP both in anterior olfactory structures (that is, in mitral cells and tufted cells of the olfactory bulb and in projection neurons of the anterior olfactory nucleus) and in preganglionic parasympathetic neurons of the dorsal motor nucleus of the vagal nerve, including the adjoining intermediate reticular zone, during stage 1 [19, 31, 42, 71, 137, 138, 179, 180]. Thread-like LNs appear in unmyelinated long and thin axons that course in small bundles obliquely through the medulla oblongata and as spindle-shaped dendritic LNs and globular somatic LBs in visceromotor projection neurons of the vagal dorsal

motor nucleus. The motor cells in the ambiguous nucleus that send well-myelinated thick axons to striated muscles of the proximal oesophagus are spared [71, 179].

The nerve cells within the olfactory epithelium have not been reported to develop α -synuclein aggregates [181]; but even if the contrary were to prove true, the presence of subtle LP might well escape neuropathological detection because the nerve cells there are constantly being replaced and defective or damaged neurons are subject to removal from the functional neuroepithelium. This region, which is directly connected to the CNS and, at the same time, provides direct access to the upper respiratory tract, has effective mechanisms for repulsing invading pathogens [182].

It should also be noted that stage 1 involvement can be confined to the olfactory bulb alone [31, 175, 183]. For this reason, the possibility cannot be excluded that a toxin or pathogen may penetrate beyond the olfactory epithelium into the olfactory bulb, where it initiates the formation of α -synuclein aggregates or, alternatively, whence α -synuclein aggregation is induced in more distant interconnected regions [167, 184–188]. The site or region where the earliest aggregates occur is important for potential therapeutic strategies. In this context, the intimate spatial relationships between the nervous system and external as well as internal surfaces of the body (CNS – olfactory epithelium, ENS – gastrointestinal lumen) as potential entry points for environmental toxins and pathogens bears repeating [106, 189]. Swallowed nasal secretions, for example, come directly into contact with the upper gastrointestinal tract [78, 139, 167, 168, 190, 191].

The pathological process in the brain progresses into the serotonergic nuclei of the lower raphe system, the magnocellular portions of the reticular formation, and the noradrenergic locus coeruleus in stage 2 [31, 71, 192]. Descending projections from all of these ‘level setting’ nuclei regulate medullary and spinal cord centres for somato- and viscerosensory input and viscerosomatomotor output (Figure 1) [193–195]. Short and especially strong projections reach the dorsal motor nucleus of the vagal nerve. Spinal cord involvement begins during stage 2 with spindle-shaped axonal LNs in long descending projections generated from the involved supraspinal level setting nuclei that terminate in layer 7 preganglionic sympathetic (IML) and parasympathetic (SPS) nuclei (Figure 1) [7, 13, 19]. This sequence of spinal cord involvement indicates that

aggregated α -synuclein could progress in an antero-grade direction to spinal cord axonal networks in lateral portions of layer 7 (visceromotor autonomic system; IML, SPS) [13, 196, 197].

During stage 3, the somatodendritic compartments of neurons in the spinal cord that receive supraspinal projections from the level setting nuclei develop LP: dorsal horn layer 1 projection neurons (nociception) [198] and layer 9 ventral horn α -motoneurons, including the motoneurons of Onuf’s nucleus (somatomotor system) (Figure 1). As in stage 2, the pathology could progress anterogradely and trans-synaptically into the somatodendritic compartments of vulnerable nerve cells [13]. A route that would involve retrograde axonal transport of the lesions to level setting nuclei and to layer 1 of the dorsal horn could originate in the plexus of the ENS. From there, aggregated α -synuclein could spread to preganglionic sympathetic neurons of the IML in layer 7 via postganglionic sympathetic projection neurons located in the coeliac ganglion (Figure 1) [7, 19, 198, 199].

In the brain, the caudo-rostral trajectory extends in stage 3 towards the mesencephalic tegmentum and basal portions of the prosencephalon – at this point, the pathological process bifurcates, entering, first, the central subnucleus and, subsequently, the basolateral subnuclei of the amygdala (Figure 1). The central subnucleus of the amygdala is reciprocally connected with the level setting nuclei and with the vagal dorsal motor nucleus (Figure 1) [37], thereby providing dense supervening limbic input to these supraspinal nuclei [200, 201]. Essential somatomotor centres also become involved at this stage, namely, the tegmental pedunclopontine nucleus [202] and the pars compacta of the substantia nigra (Figure 1). The pathological process in the substantia nigra may be induced via its interconnectivities with the central subnucleus of the amygdala or via direct coeruleo-nigral and pedunclopontine-nigral projections [203, 204]. During stage 3, LP also develops in the serotonergic nuclei of the upper raphe system, the histaminergic tuberomammillary nucleus and the cholinergic magnocellular nuclei of the basal forebrain [31, 71, 173].

The routes described up to this point are also used by neurotropic viruses (e.g., pseudorabies) which, in experimental animals, have been inoculated into the gastric epithelium and progress sequentially through the ENS, the vagal nerve (including the dorsal motor

nucleus) and the nuclei of the level setting system to arrive at the central subnucleus of the amygdala and beyond [167, 205–209]. Nonetheless, the viruses utilized reach all of these sites in a matter of days [210], whereas the PD-related pathological process takes years to do so [139, 211].

Additional regions of the forebrain and, for the first time, a specific portion of the cerebral cortex, the transentorhinal region, become involved in stage 4. This specialized region of the cortex mediates between the temporal allocortex (hippocampal formation and entorhinal region) and the high-order association areas of the temporal neocortex. The region is highly developed only in higher primates, and above all in humans [212, 213]. The fact that the region is lacking in most experimental animals is probably the reason why the routes taken by the viral agents do not progress beyond the level of the amygdala [188]. By contrast, in humans, the transentorhinal region develops the most severe LP within the entire cerebral cortex [31].

The locus coeruleus, the basolateral nuclei of the amygdala, and the magnocellular nuclei of the basal forebrain are bidirectionally connected with the transentorhinal region, and these connectivities could lead the pathological process to the cortex. As if through a bottle-neck, the data stream from neocortical high-order sensory association areas must transit through the transentorhinal/entorhinal regions and, from there, is transferred via the perforant path to reach the hippocampal formation (Figure 1) [214, 215]. Via the perforant path and, additionally, along axons generated from the tuberomammillary nucleus to the second sector of the Ammon's horn, the pathological process could enter the hippocampal formation [31, 39].

The fullest topographic extent of the PD-related process is reached during stages 5 and 6. Beginning with the transentorhinal region, LP gradually penetrates the entire neocortex [31, 116, 216]. During stage 5, the lesions develop not only in all high-order sensory association areas but also in prefrontal fields of the neocortex (at first, in superordinate fields that process interoceptive data and regulate visceromotor functions). In stage 6, the pathological process finally makes inroads into the neocortical first-order sensory association areas, premotor fields, and even the primary sensory and motor fields (Figure 1) [31, 71]. The small- to medium-sized pyramidal cells that contain α -synuclein

aggregates occur chiefly in neocortical layers V and VI [160, 217].

It still is unknown whether the involved pyramidal cells generate cortico-striatal or cortico-cortical projections, or both. The progression of the disease process throughout the neocortex, however, appears to take place via cortico-cortical projections directed from high-order association areas (that is, the prefrontal fields) to first-order association areas (that is, premotor fields) and, from there, to primary areas. These projections possess diffuse synaptic contacts in neocortical layers I–III and V–VI but very few in layer IV [218–220]. Involvement of these interconnectivities is consistent with the fact that layer IV remains virtually intact in PD, whereas LP develops mostly in layers V and VI. Thus, we have been inclined to view the disease process within the cerebral cortex as most likely progressing chiefly via these cortico-cortical pathways, supported additionally by coeruleo-cortical projections and projections from upper raphe nuclei and magnocellular nuclei of the basal forebrain (Figure 1).

Developing new models of α -synuclein spreading along axonal connectivities

We formerly speculated that an environmental or neurotropic pathogen might adhere to the mucous membrane of the upper gastrointestinal tract and/or nose and penetrate into neuronal processes of neurons there, where it might trigger a conformational change in α -synuclein. Then, via transneuronal and retrograde axonal transport along unmyelinated axons of the vagal nerve, α -synuclein aggregates from the ENS could reach the preganglionic neurons of the dorsal motor nucleus of the vagal nerve [116, 167, 168, 210, 211, 221–224]. Inasmuch as the vagus nerve exerts its influence chiefly on the distal portions of the oesophagus and following proximal portions of the stomach [225–227], these regions are among the more likely sites for a transfer and propagation of small α -synuclein aggregates between interconnected neurons. Indeed, we recently reported a case in which LP was present only in the olfactory bulb and in the submucosal plexus of the lower oesophagus [13]. Of course, it is impossible to extrapolate to a general rule from isolated occurrences [12, 228, 229]. At the same time, however, experimental models have made it possible to detect seeding mechanisms [230, 231] and spreading

over considerable distances by α -synuclein aggregates from the periphery via the vagus nerve to the CNS following intragastric, intraduodenal, and peripheral vagal nerve inoculations [199, 232–234]. There is some experimental evidence for retrograde axonal transport from the PNS to the CNS [235], but the connectivities between the ENS and the CNS via the vagus nerve probably play a pivotal role in the progression of PD [236, 237].

Experimental models involving the ENS and CNS have received additional support from a very recent large-scale epidemiologic evaluation of vagotomies that were performed in the past to treat peptic ulcers [238]. Full vagotomy is defined by resection of both vagal trunks, whereas selective vagotomy entails resection of only terminal branches supplying the fundus and corpus of the stomach [238–240]. The results of the study illustrate that the risk of having developed sporadic PD at follow-up 10 years later was significantly reduced in individuals who had undergone full truncal vagotomy but remained nearly similar to the risk of the general population in individuals having undergone selective vagotomy [238].

Taken together, the latest evidence indicates that the brain regions and the types of nerve cells that become sequentially in PD involved are anatomically interconnected, often over long distances, and that physical contacts between nerve cells as well as axonal transport are involved in PD pathogenesis [31, 221, 241]. Misfolded and slightly aggregated α -synuclein seeds can spread trans-synaptically along multisynaptic pathways and function, probably in a strain-dependent manner, as self-propagating pathogens for disease progression [70, 78, 99, 222, 223, 242–261]. Whether neuronal loss or impairment of cellular functions, the disease-related damage revolves around superordinate centres of the limbic, visceromotor, and somatomotor systems (Table 1, Figure 1). With the exception of olfaction and portions of the pain system (that is, the large neurons in layer 1 of the spinal cord), the sensorium is affected to a lesser extent [10, 262–267].

Studies of pathological mechanisms that underlie the conformational change and aggregation of α -synuclein, as well as those involving the formation of pathogens that can be transported axonally and transmitted trans-synaptically to interconnected nerve cells, where they induce α -synuclein misfolding and aggregation in a previously uninvolved cell, cannot be performed on

formalin-fixed autopsy tissue. Nonetheless, experimental cellular models can only approximate the cellular milieu found in the human nervous system, where LP (as opposed to 'Lewy-like' lesions in nonhuman models) develops only in a few susceptible types of long-lived projection neurons. Resected tissue culled from surgical interventions on the human gastrointestinal tract, for example, contains functional groups of postmitotic myenteric plexus nerve cells [268–270]. Although admittedly challenging to design, a human-based ENS model might yield even more accurate insights into the mechanisms of α -synuclein aggregation and, possibly, also transmission in PD [78, 116, 225, 226].

Acknowledgements

The authors thank Mr. David Ewert for technical assistance and support (figures), Ms. Simone Feldengut (organization of the Tables), the Michael J. Fox Foundation for Parkinson's Research, the Goethe University Frankfurt (Braak Collection), and the families and patients who, by means of brain and organ donations, make our work possible.

Author contributions

Both authors contributed to the conception of the text. The first author drafted and revised the manuscript. The senior author designed and drafted the figure.

References

- 1 de Rijk MC, Launer LJ, Berger K, Breteler MM, Dartigues JF, Baldereschi M, Fratiglioni L, Lobo A, Martinez-Lage J, Trenkwalder C, Hofman A. Prevalence of Parkinson's disease in Europe: A collaborative study of population-based cohorts. *Neurology* 2000; **54**(Suppl 5): 21–23.
- 2 Fahn S. Description of Parkinson's disease as a clinical syndrome. *Ann N Y Acad Sci* 2003; **991**: 1–14.
- 3 Dorsey ER, Constantinesco R, Thompson JP, Biglan KM, Holloway RG, Kieburtz K, Marshall FJ, Ravina BM, Schifitto G, Siderowf A, Tanner CM. Projected number of people with Parkinson's disease in the most populous nations, 2005 through 2030. *Neurology* 2005; **68**: 384–6.
- 4 de Lau L, Breteler MM. Epidemiology of Parkinson's disease. *Lancet Neurol* 2006; **5**: 525–35.
- 5 den Hartog Jager WA, Bethlem J. The distribution of Lewy bodies in the central and autonomic nervous systems in idiopathic paralysis agitans. *J Neurol Neurosurg Psychiatry* 1960; **23**: 283–90.

- 6 Wakabayashi K, Takahashi H, Takeda S, Ohama E, Ikuta F. Parkinson's disease: the presence of Lewy bodies in Auerbach's and Meissner's plexuses. *Acta Neuropathol* 1988; **76**: 217–21.
- 7 Wakabayashi K, Takahashi H. The intermediolateral nucleus and Clarke's column in Parkinson's disease. *Acta Neuropathol* 1997; **94**: 287–9.
- 8 Deguchi K, Sasaki I, Tsukaguchi M, Kamoda M, Touge T, Takeuchi H, Kuriyama S. Abnormalities of rate-corrected QT intervals in Parkinson's disease – a comparison with multiple system atrophy and progressive supranuclear palsy. *J Neurol Sci* 2002; **199**: 31–7.
- 9 Orimo S, Takahashi A, Uchihara T, Mori F, Kakita A, Wakabayashi K, Takahashi H. Degeneration of cardiac sympathetic nerve begins in the early disease process of Parkinson's disease. *Brain Pathol* 2007; **17**: 24–30.
- 10 Ikemura M, Saito Y, Sengoku Y, Sakiyama Y, Hatsuta H, Kanemaru K, Sawabe M, Arai T, Ito G, Iwatsubo T, Fukayama M, Murayama S. Lewy body pathology involves cutaneous nerves. *J Neuropathol Exp Neurol* 2008; **67**: 945–53.
- 11 Djaldetti R, Lev N, Melamed E. Lesions outside the CNS in Parkinson's disease. *Mov Disord* 2009; **24**: 793–800.
- 12 Beach TG, Adler CH, Sue LI, Vedders L, Lue L, White CL III, Akiyama H, Caviness JN, Shill HA, Sabbagh MN, Walker DG. Arizona Parkinson's Disease Consortium. Multi-organ distribution of phosphorylated α -synuclein histopathology in subjects with Lewy body disorders. *Acta Neuropathol* 2010; **119**: 689–702.
- 13 Del Tredici K, Braak H. Spinal cord lesions in sporadic Parkinson's disease. *Acta Neuropathol* 2012; **124**: 643–66.
- 14 Cersósimo MG, Tumilasci OR, Raina GB, Benarroch EE, Cardoso EM, Micheli F, Pazo JH. Hyposialorrhea as an early manifestation of Parkinson disease. *Auton Neurosci* 2009; **150**: 150–1.
- 15 Gallagher DA, Lees AJ, Schrag A. What are the most important nonmotor symptoms in patients with Parkinson's disease and are we missing them? *Mov Disord* 2010; **25**: 2493–500.
- 16 Ali K, Morris HR. Parkinson's disease: chameleons and mimics. *Pract Neurol* 2015; **15**: 14–25.
- 17 Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry* 1992; **55**: 181–4.
- 18 Koller WC. How accurately can Parkinson's disease be diagnosed? *Neurology* 1992; **42**: 6–16.
- 19 Bloch A, Probst A, Bissig H, Adams H, Tolnay M. α -Synuclein pathology of the spinal and peripheral autonomic nervous system in neurologically unimpaired elderly subjects. *Neuropathol Appl Neurobiol* 2006; **12**: 284–95.
- 20 Dickson DW, Braak H, Duda JE, Duyckaerts C, Gasser T, Halliday GM, Hardy J, Leverenz JB, Del Tredici K, Wszolek ZK, Litvan I. Diagnostic criteria for the neuropathological assessment of Parkinson disease. *Lancet Neurol* 2009; **8**: 1150–7.
- 21 Goedert M. The awakening of α -synuclein. *Nature* 1997; **388**: 232–3.
- 22 Spillantini MG, Schmidt ML, Lee VM, Trojanowski JQ, Jakes R, Goedert M. α -Synuclein in Lewy bodies. *Nature* 1997; **388**: 839–40.
- 23 Wakabayashi K, Matsumoto K, Takayama K, Yoshimoto M, Takahashi A. NACP, a presynaptic protein, immunoreactivity in Lewy bodies in Parkinson's disease. *Neurosci Lett* 1997; **239**: 45–8.
- 24 Duda JE, Lee VM, Trojanowski JQ. Neuropathology of synuclein aggregates: new insights into mechanism of neurodegenerative diseases. *J Neurosci Res* 2000; **61**: 121–7.
- 25 Takahashi H, Wakabayashi K. The cellular pathology of Parkinson's disease. *Neuropathology* 2001; **21**: 315–22.
- 26 Trojanowski JQ, Lee VM. Parkinson's disease and related synucleinopathies are a class of nervous system amyloidoses. *Neurotoxicology* 2002; **23**: 457–60.
- 27 Trojanowski JQ, Lee VM. Meeting summary – cell biology of Parkinson's disease and related neurodegenerative disorders. *Sci Aging Knowl Environ* 2003; **13**: 23.
- 28 Jellinger KA, Mizuno Y. Parkinson's disease. In *Neurodegeneration: The Molecular Pathology of Dementia and Movement Disorders* 1st edn. Ed Dickson DW. Basel: ISN Neuropathol Press, 2003; 159–87.
- 29 Ohama E, Ikuta F. Parkinson's disease: distribution of Lewy bodies and monoamine neuron system. *Acta Neuropathol* 1976; **34**: 311–9.
- 30 Braak H, Del Tredici K. Poor and protracted myelination as a contributory factor to neurodegenerative disorders. *Neurobiol Aging* 2004; **25**: 19–23.
- 31 Braak H, Del Tredici K. Neuroanatomy and pathology of sporadic Parkinson's disease. *Adv Anat Embryol Cell Biol* 2009; **201**: 1–119.
- 32 Malek N, Swallow D, Grosset KA, Anichtchik O, Spillantini M, Grosset DG. Alpha-synuclein in peripheral tissues and body fluids as a biomarker for Parkinson's disease – a systematic review. *Acta Neurol Scand* 2014; **130**: 59–72.
- 33 Lewy FH. Paralysis agitans. I. Pathologische Anatomie. In *Handbuch der Neurologie*. Ed Lewandowski M. Berlin: Springer, 1912; 920–33.
- 34 Gibb WRG, Lees AJ. The relevance of the Lewy body to the pathogenesis of idiopathic Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1988; **51**: 745–52.
- 35 Gibb WRG, Lees AJ. The significance of the Lewy body in the diagnosis of idiopathic Parkinson's disease. *Neuropathol Appl Neurobiol* 1989; **15**: 27–44.
- 36 Lowe J. Lewy bodies. In *DP Calne*. Ed N Diseases. Philadelphia: Saunders, 1994; 51–69.

- 37 Braak H, Braak E, Yilmazer D, de Vos RAl, Jansen ENH, Bohl J, Jellinger K. Amygdala pathology in Parkinson's disease. *Acta Neuropathol* 1994; **88**: 493–500.
- 38 Braak H, Sandmann-Keil D, Gai WP, Braak E. Extensive axonal Lewy neurites in Parkinson's disease: a novel pathological feature revealed by α -synuclein immunocytochemistry. *Neurosci Lett* 1999; **265**: 67–9.
- 39 Dickson DW, Schmidt ML, Lee VMY, Zhao ML, Yen SH, Trojanowski JQ. Immunoreactivity profile of hippocampal CA2/3 neurites in diffuse Lewy body disease. *Acta Neuropathol* 1994; **87**: 269–76.
- 40 Gai WP, Blessing WW, Blumbergs PC. Ubiquitin-positive degenerating neurites in the brainstem in Parkinson's disease. *Brain* 1995; **118**: 1447–59.
- 41 Kuusisto E, Parkkinen L, Alafuzoff I. Morphogenesis of Lewy bodies: dissimilar incorporation of α -synuclein, ubiquitin, and p62. *J Neuropathol Exp Neurol* 2003; **62**: 1241–53.
- 42 Saito Y, Kawashima A, Ruberu NN, Fujiwara H, Koyama S, Sawabe M, Arai T, Nagura H, Yamanouchi H, Hasegawa M, Iwatsubo T, Murayama S. Accumulation of phosphorylated α -synuclein in aging human brain. *J Neuropathol Exp Neurol* 2003; **62**: 644–54.
- 43 Fearnley JM, Lees AJ. Pathology of Parkinson's disease. In *Neurodegenerative Diseases*. Ed DB Calne. Philadelphia: Saunders, 1994; 545–54.
- 44 Forno LS. Neuropathology of Parkinson's disease. *J Neuropathol Exp Neurol* 1996; **55**: 259–72.
- 45 Dickson DW. α -Synuclein and the Lewy body disorders. *Curr Opin Neurol* 2001; **14**: 423–32.
- 46 Goedert M. The significance of tau and α -synuclein inclusions in neurodegenerative diseases. *Curr Opin Genet Dev* 2001; **11**: 343–51.
- 47 Goedert M, Spillantini MG, Del Tredici K, Braak H. 100 years of Lewy pathology. *Nat Rev Neurol* 2013; **9**: 13–24.
- 48 Norris EH, Giasson BI, Lee VM. α -Synuclein: normal function and role in neurodegenerative diseases. *Curr Top Dev Biol* 2004; **60**: 17–54.
- 49 Tofaris GK, Spillantini MG. Physiological and pathological properties of alpha-synuclein. *Cell Mol Life Sci* 2007; **64**: 2194–201.
- 50 Stamelou M, Hoeglenger GU. Atypical parkinsonism: an update. *Curr Opin Neurol* 2013; **26**: 401–5.
- 51 El-Agnaf OM, Salem SA, Paleologou KE, Cooper LJ, Fullwood NJ, Gibson MJ, Curran MD, Court JA, Mann DM, Ikeda S, Cookson MR, Hardy J, Allsop D. Alpha-synuclein implicated in Parkinson's disease is present in extracellular biological fluids, including human plasma. *FASEB J* 2003; **17**: 1945–7.
- 52 Barbour R, Kling K, Anderson JP, Banducci K, Cole T, Diep L, Fox M, Goldstein JM, Soriano F, Seubert P, Chilcote TJ. Red blood cells are the major source of alpha-synuclein in blood. *Neurodegener Dis* 2008; **5**: 55–9.
- 53 Braak H, Del Tredici K, Gai WP, Braak E. Alpha-synuclein is not a requisite component of synaptic boutons in the adult human central nervous system. *J Chem Neuroanat* 2001; **20**: 245–52.
- 54 Clayton DF, George JM. Synucleins in synaptic plasticity and neurodegenerative disorders. *J Neurosci Res* 1999; **58**: 120–9.
- 55 Fortin DL, Nemani VM, Voglmaier SM, Anthony MD, Ryan TA, Edwards RH. Neural activity controls the synaptic accumulation of α -Synuclein. *Neurobiol Dis* 2005; **25**: 10913–21.
- 56 Stefanis L. α -Synuclein in Parkinson's disease. *Cold Spring Harb Perspect Med* 2012; **4**: a009399.
- 57 Anichtchik O, Calo L, Spillantini MG. Synaptic dysfunction in synucleinopathies. *CNS Neurol Disord Drug Targets* 2013; **12**: 1094–100.
- 58 Lashuel HA, Overk CA, Queslatti A, Masliah E. The many faces of α -synuclein: from structure and toxicity to therapeutic target. *Nat Rev Neurosci* 2013; **14**: 38–48.
- 59 Silva BA, Breydo L, Uversky VN. Targeting the chameleon: a focused look at α -synuclein and its roles in neurodegeneration. *Mol Neurobiol* 2013; **47**: 446–59.
- 60 Goedert M. Alzheimer's and Parkinson's diseases: the prion concept in relation to assembled A β , tau, and α -synuclein. *Science* 2015; **349**: 125555.
- 61 Giasson BI, Murray IV, Trojanowski JQ, Lee VM. A hydrophobic stretch of 12 amino acid residues in the middle of alpha-synuclein is essential for filament assembly. *J Biol Chem* 2001; **276**: 2380–6.
- 62 Rodriguez JA, Ivanova MI, Sawaya MR, Cascio D, Reyes FE, Shi D, Sangwan S, Guenther EL, Johnson LM, Zhang M, Jiang L, Arbing MA, Nannenga BI, Hattne J, Whitelegge J, Brewster AS, Messerschmidt M, Boutet S, Sauter NK, Gonen T, Eisenberg DS. Structure of the toxic core of α -synuclein from invisible crystals. *Nature* 2015; **525**: 486–90.
- 63 Goedert M, Cheng Y. Parkinson's disease: crystals of a toxic core. *Nature* 2015; **525**: 458–9.
- 64 Bartels T, Choi JG, Selkoe DJ. α -synuclein occurs physiologically as a helically folded tetramer that resists aggregation. *Nature* 2011; **477**: 107–10.
- 65 Witt SN. Molecular chaperones, α -synuclein, and neurodegeneration. *Mol Neurobiol* 2013; **47**: 552–60.
- 66 Selkoe D, Dettmer U, Luth E, Kim N, Newman A, Bartels T. Defining the native state of α -synuclein. *Neurodegener Dis* 2014; **13**: 114–7.
- 67 Dettmer U, Newman AJ, Soldner F, Luth ES, Kim NC, vonSaucken VE, Sanderson JB, Jaenisch R, Bartels T, Selkoe D. Parkinson-causing α -synuclein missense mutations shift native tetramers to monomers as a mechanism of disease. *Nat Commun* 2015; **6**: 7314.
- 68 Fauvet B, Mbefo MK, Fares MB, Desobry C, Michael S, Ardah MT, Tsika E, Coune P, Prudent M, Lion N, Eliezer D, Moore DJ, Schneider B, Aebischer P,

- El-Agnaf OM, Masliah E, Lashuel HA. α -Synuclein in central nervous system and from erythrocytes, mammalian cells, and *Escherichia coli* exists predominantly as disordered monomer. *J Biol Chem* 2012; **287**: 15345–64.
- 69 Burré J, Vivona S, Sidao J, Sharma M, Brunger AT, Sudhof TC. Properties of native brain α -Synuclein. *Nature* 2013; **498**: E4–7.
- 70 Kim WS, Kägedal K, Halliday GM. Alpha-synuclein biology in Lewy body disease. *Alzheimers Res Ther* 2014; **6**: 73.
- 71 Braak H, Del Tredici K, Rüb U, de Vos RA, Jansen Steur EN, Braak E. Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging* 2003; **24**: 197–210.
- 72 Orimo S, Amino T, Itoh Y, Takahashi A, Kojo T, Uchihara T, Tsuchiya K, Mori F, Wakabayashi K, Takahashi H. Cardiac sympathetic denervation precedes neuronal loss in the sympathetic ganglia in Lewy body disease. *Acta Neuropathol* 2005; **109**: 583–8.
- 73 Volpicelli-Daley LA, Luk KC, Patel TP, Tanik SA, Riddle DM, Stieber A, Meaney DF, Trojanowski JQ, Lee VM, et al. Exogenous α -synuclein fibrils induce Lewy body pathology leading to synaptic dysfunction and neuron death. *Neuron* 2011; **72**: 57–71.
- 74 Recasens K, Dehay B, Bove J, Carballo-Carbaja L, Dovero S, Pérez-Villaba A, Fernagut PO, Biesa J, Parent A, Perier C, Fariñas I, Obeso JA, Bezard E, Vila M. Lewy body extracts from Parkinson disease brains trigger alpha-synuclein pathology and neurodegeneration in mice and monkeys. *Ann Neurol* 2014; **75**: 351–62.
- 75 Tanji K, Mori F, Mimura J, Itho K, Kakita A, Takahashi H, Wakabayashi K. Proteinase K-resistant α -synuclein is deposited in presynapses in human Lewy body disease and A53T α -synuclein transgenic mice. *Acta Neuropathol* 2010; **120**: 145–54.
- 76 Pieri L, Madiona K, Bousset L, Melki R. Fibrillar α -synuclein and huntingtin exon 1 assemblies are toxic to cells. *Biophys J* 2012; **102**: 2894–905.
- 77 Prasad K, Beach TG, Hedreen J, Richfield EK. Critical role for truncated α -synuclein and aggregates in Parkinson's disease and incidental Lewy body disease. *Brain Pathol* 2012; **22**: 811–25.
- 78 Dehay B, Bourdenx M, Gorry P, Przedborski S, Vila M, Hunot S, Singleton A, Olanow CW, Merchant KM, Bezard E, Petsko GA, Meissner WG. Targeting α -synuclein for treatment of Parkinson's disease: mechanistic and therapeutic considerations. *Lancet Neurol* 2015; **14**: 855–66.
- 79 Leigh PN, Probst A, Dale GE, Power DP, Brion JP, Dodson A, Anderton BH. New aspects of the pathology of neurodegenerative disorders as revealed by ubiquitin antibodies. *Acta Neuropathol* 1989; **79**: 61–72.
- 80 Lennox G, Lowe J, Landon M, Byrne EJ, Mayer RJ, Godwin-Austen RB. Diffuse Lewy body disease: correlative neuropathology using anti-ubiquitin immunocytochemistry. *J Neurol Neurosurg Psychiatry* 1989; **52**: 1236–47.
- 81 Wakabayashi K, Takahashi H, Ohama E, Ikuta F. Parkinson's disease: an immunohistochemical study of Lewy body-containing neurons in the enteric nervous system. *Acta Neuropathol* 1990; **79**: 581–3.
- 82 Wakabayashi K, Tanji K, Mori F, Takahashi H. The Lewy body in Parkinson's disease: molecules implicated in the formation and degradation of alpha-synuclein aggregates. *Neuropathology* 2007; **27**: 494–506.
- 83 Wakabayashi K, Tanji K, Odagiri S, Odagiri S, Miki Y, Mori F, Takahashi H. The Lewy body in Parkinson's disease and related neurodegenerative disorders. *Mol Neurobiol* 2013; **47**: 495–508.
- 84 Katsuse O, Iseki E, Marui W, Kosaka K. Developmental stages of cortical Lewy bodies and their relation to axonal transport blockage in brains of patients with dementia with Lewy bodies. *J Neurol Sci* 2003; **211**: 29–35.
- 85 Dugger BN, Dickson DW. Cell type specific sequestration of choline acetyltransferase and tyrosine hydroxylase within Lewy bodies. *Acta Neuropathol* 2010; **120**: 633–9.
- 86 Miki Y, Tanji K, Mori F, Utsumi J, Sasaki H, Kakita A, Takahashi H, Wakabayashi K. Alteration of upstream autophagy-related proteins (ULK1, ULK2, Beclin1, VPS34, and AMBRA1) in Lewy body disease. *Brain Pathol* 2015; doi: 10.1111/bpa.12297.
- 87 Volles M, Lansbury PJ. Zeroing in on the pathogenic form of alpha-synuclein and its mechanism of neurotoxicity in Parkinson's disease. *Biochemistry* 2003; **42**: 7871–8.
- 88 Chung KK, Dawson VL, Dawson TM. The role of the ubiquitin-proteasomal pathway in Parkinson's disease and other neurodegenerative disorders. *Trends Neurosci* 2001; **24**: 7–14.
- 89 McNaught KS, Jenner P. Proteasomal function is impaired in substantia nigra in Parkinson's disease. *Neurosci Lett* 2001; **297**: 191–4.
- 90 Cuervo AM, Stefanis L, Fredenburg R, Lansbury PT, Sulzer D. Impaired degradation of mutant α -synuclein by chaperone-mediated autophagy. *Science* 2004; **305**: 1292–5.
- 91 Martinez-Vincente M, Cuervo AM. Autophagy and neurodegeneration: when the cleaning crew goes on strike. *Lancet Neurol* 2007; **6**: 352–61.
- 92 Chu Y, Dodiya H, Aebischer P, Olanow CW, Kordower JH. Alterations in lysosomal and proteasomal markers in Parkinson's disease: relationship to alpha-synuclein inclusions. *Neurobiol Dis* 2009; **35**: 385–98.

- 93 Dehay B, Bové J, Rodriguez-Muela N, Recasens A, Boya P, Vila M. Pathogenic lysosomal depletion in Parkinson's disease. *J Neurosci* 2010; **30**: 12535–44.
- 94 Lynch-Day M, Mao K, Wang K, Zhao M, Klionsky DJ. The role of autophagy in Parkinson's disease. *Cold Spring Harb Perspect Med* 2012; **2**: a009357.
- 95 Duda JE. Pathology and neurotransmitter abnormalities of dementia with Lewy bodies. *Dement Geriatr Cogn Disord* 2004; **17**: 3–14.
- 96 Saha AR, Hill J, Utton MA, Assuni AA, Ackerley S, Grierson AJ, Miller CC, Davies AM, Buchman VL, Anderton BH, Hanger DP. Parkinson's disease alpha-synuclein mutations exhibit defective axonal transport in cultured neurons. *J Cell Sci* 2004; **117**: 1017–24.
- 97 Kanazawa T, Adachi E, Orimo S, Nakamura A, Mizusawa H, Uchichara T. Pale neurites, premature α -synuclein aggregates with centripetal extension from axon collaterals. *Brain Pathol* 2012; **22**: 67–78.
- 98 Roy S, Zhang B, Lee VM, Trojanowski JQ. Axonal transport defects: a common theme in neurodegenerative diseases. *Acta Neuropathol* 2005; **109**: 5–13.
- 99 Lamberts JT, Hildebrandt EN, Brundin P. Spreading of α -synuclein in the face of axonal transport deficits in Parkinson's disease: a speculative synthesis. *Neurobiol Dis* 2015; **77**: 276–83.
- 100 Herrup K, Yang Y. Cell cycle regulation in the post-mitotic neuron: oxymoron or new biology? *Nat Rev Neurosci* 2007; **8**: 368–78.
- 101 Webb J, Ravikumar B, Athkins J, Skepper J, Rubinsztajn D. Alpha-Synuclein is degraded by both autophagy and the proteasome. *J Biol Chem* 2003; **278**: 25009–13.
- 102 Levine B, Kroemer G. Autophagy in the pathogenesis of disease. *Cell* 2008; **132**: 27–42.
- 103 Breydo L, Wu JW, Uversky VN. A-synuclein misfolding and Parkinson's disease. *Biochim Biophys Acta* 2012; **1822**: 261–85.
- 104 Bandopadhyay R, de Belleruche J. Pathogenesis of Parkinson's disease: emerging role of molecular chaperones. *Trends Mol Med* 2010; **16**: 27–36.
- 105 Cox D, Carber JA, Ecroyd H. Preventing α -synuclein aggregation: the role of the small heat-shock molecular chaperone proteins. *Biochim Biophys Acta* 2014; **1842**: 1830–43.
- 106 Zhou L, Miranda-Saksena M, Saksena NK. Viruses and neurodegeneration. *J Virol* 2013; **10**: 172–86.
- 107 Marreiros R, Müller-Schiffmann A, Bader V, Selvarajah S, Dey D, Lingappa VR, Korth C. Viral capsid assembly as a model for protein aggregation diseases: active processes catalyzed by cellular assembly machines composing novel drug target. *Virus Res* 2015; **207**: 155–64.
- 108 Orimo S, Kanazawa T, Itoh Y, Wakabayashi K, Kakita A. Unmyelinated axons are more vulnerable to degeneration than myelinated axons of the cardiac nerve in Parkinson's disease. *Neuropathol Appl Neurobiol* 2011; **37**: 791–802.
- 109 Lang AE, Obeso JA. Challenges in Parkinson's disease: restoration of the nigrostriatal dopamine system is not enough. *Lancet Neurol* 2004; **3**: 309–16.
- 110 Langston JW. The Parkinson's complex: parkinsonism is just the tip of the iceberg. *Ann Neurol* 2006; **59**: 591–6.
- 111 Ahlskog JE. Beating a dead horse: Dopamine and Parkinson's disease. *Neurology* 2007; **69**: 1701–11.
- 112 Dickson DW, Fujishiro H, Orr C, DelleDonne A, Josephs KA, Frigerio R, Burnett M, Parisi JE, Klos KJ, Ahlskog JE. Neuropathology of non-motor features of Parkinson's disease. *Parkinsonism Rel Disord* 2009; **15**(Suppl 3): S1–S5.
- 113 Halliday GM, Lees AJ, Stern M. Milestones in Parkinson's disease – clinical and pathologic features. *Mov Disord* 2011; **26**: 1015–21.
- 114 Wakabayashi K, Takahashi H, Ohama E, Takeda S, Ikuta F. Lewy bodies in the visceral autonomic nervous system in Parkinson's disease. *Adv Neurol* 1993; **60**: 609–12.
- 115 Iwanaga K, Wakabayashi K, Yoshimoto M, Tomita I, Satoh H, Takashima H, Satoh A, Seto M, Tsujihata M, Takahashi H. Lewy body-type degeneration in cardiac plexus in Parkinson's and incidental Lewy body diseases. *Neurology* 1999; **52**: 1269–71.
- 116 Braak H, de Vos RAI, Bohl J, Del Tredici K. Gastric α -synuclein immunoreactive inclusions in Meissner's and Auerbach's plexuses in cases staged for Parkinson's disease-related brain pathology. *Neurosci Lett* 2006; **396**: 67–72.
- 117 Klos KJ, Ahlskog JE, Josephs KA, Apaydin H, Parisi JE, Boeve BF, DeLucia MW, Dickson DW. α -Synuclein pathology in the spinal cord of neurologically asymptomatic aged individuals. *Neurology* 2006; **66**: 1100–2.
- 118 Ross GW, Abbott RD, Petrovitch H, Tanner CM, Davis DG, Nelson J, Markesbery WR, Hardman J, Masaki K, Launer L, White LR. Association of olfactory dysfunction with incidental Lewy bodies. *Mov Disord* 2006; **21**: 2062–7.
- 119 Abbott RD, Ross GW, Petrovitch H, Tanner CM, Davis DG, Masaki KH, Launer LJ, Curb JD, White LR. Bowel movement frequency in late-life and incidental Lewy bodies. *Mov Disord* 2007; **22**: 1581–6.
- 120 Minguez-Castellanos A, Chamorro CE, Escamilla-Sevilla F, Ortega-Moreno A, Rebollo AC, Gomez-Rio M, Concha A, Munoz DG. Do α -synuclein aggregates in autonomic plexuses predate Lewy body disorders? A cohort study *Neurology* 2007; **68**: 1012–8.
- 121 DelleDonne A, Klos KJ, Fujishiro H, Ahmed Z, Parisi JR, Josephs KA, Frigerio R, Burnett M, Wszolek ZK, Uitti RJ, Ahlskog JE, Dickson DW. Incidental Lewy body disease and preclinical Parkinson disease. *Arch Neurol* 2008; **65**: 1074–80.

- 122 Dickson DW, Fujishiro H, DelleDonne A, Menke J, Zeshan A, Klos K, Josephs KA, Frigerio R, Burnett M, Parisi JE, Ahlskog JE. Evidence that incidental Lewy body disease is presymptomatic Parkinson's disease. *Acta Neuropathol* 2008; **115**: 437–44.
- 123 Markesbery WR, Jicha GA, Liu H, Schmitt FA. Lewy body pathology in normal elderly subjects. *J Neuropathol Exp Neurol* 2009; **68**: 816–22.
- 124 Savica R, Rocca WA, Ahlskog R. When does Parkinson disease start? *Arch Neurol* 2010; **67**: 798–801.
- 125 Del Tredici K, Hawkes CH, Ghebremedhin E, Braak H. Lewy pathology in the submandibular gland of individuals with incidental Lewy body disease and sporadic Parkinson's disease. *Acta Neuropathol* 2010; **119**: 703–13.
- 126 Del Tredici K, Braak H. Lewy pathology and neurodegeneration in premotor Parkinson's disease. *Mov Disord* 2012; **27**: 597–607.
- 127 Shannon KM, Deshavarzian A, Mutlu E, Dodiya D, Jaglin JA, Kordower JH. Alpha-synuclein in colonic submucosa in early untreated Parkinson's disease. *Mov Disord* 2012; **27**: 709–15.
- 128 Tamura T, Yoshida M, Hashizume Y, Sobue G. Lewy body-related α -synucleinopathy in the spinal cord of cases with incidental Lewy body disease. *Neuropathology* 2012; **32**: 13–22.
- 129 Navarro-Otano J, Gelpi E, Mestres CA, Quintana E, Rauek S, Ribalta T, Santiago V, Tolosa E. Alpha-synuclein aggregates in epicardial fat tissue in living subjects without parkinsonism. *Parkinsonism Rel Disord* 2013; **19**: 27–31.
- 130 Milber JM, Noorigian JV, Morley JF, Petrovitch H, White L, Ross GW, Duda JE. Lewy pathology is not the first sign of degeneration in vulnerable neurons in Parkinson's disease. *Neurology* 2012; **79**: 2307–14.
- 131 Tolosa E, Gaig C, Santamaria J, Compta Y. Diagnosis and the premotor phase of Parkinson disease. *Neurology* 2009; **72**(7 Suppl): S12–S20.
- 132 Schapira AH, Tolosa E. Molecular and clinical prodrome of Parkinson disease: implications for treatment. *Nat Rev Neurol* 2010; **6**: 309–17.
- 133 Olanow CW, Obeso JA. The significance of defining preclinical or prodromal Parkinson's disease. *Mov Disord* 2012; **27**: 666–9.
- 134 Boeve BF. Idiopathic REM sleep behaviour disorder in the development of Parkinson's disease. *Lancet Neurol* 2013; **12**: 469–82.
- 135 Chen H, Burton EA, Ross GW, Huang X, Savica R, Abbott RD, Ascherio A, Caviness JN, Gao X, Gray KA, Hong JS, Kamel F, Jennings D, Kirshner A, Lawler C, Liu R, Miller GW, Nussbaum R, Peddada SD, Rick AC, Ritz B, Siderowf AD, Tanner CM, Tröster AI, Zhang J. Research on the premotor symptoms of Parkinson's disease: clinical and etiological implications. *Environ Health Perspect* 2013; **121**: 1245–52.
- 136 Doty RL, Stern MB, Pfeiffer C, Gollomp SM, Hurtig HI. Bilateral olfactory dysfunction in early stage treated and untreated idiopathic Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1992; **55**: 138–42.
- 137 Hawkes CH, Shephard BC, Daniel SE. Olfactory dysfunction in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1997; **62**: 436–46.
- 138 Hawkes CH, Shephard BC, Daniel SE. Is Parkinson's disease a primary olfactory disorder? *Q J Med* 1999; **92**: 473–80.
- 139 Hawkes CH, Del Tredici K, Braak H. A timeline for Parkinson's disease. *Parkinsonism Rel Disord* 2010; **16**: 79–84.
- 140 Doty RL. Olfaction. *Ann Rev Psychol* 2001; **52**: 423–52.
- 141 Doty RL. Olfactory dysfunction in Parkinson disease. *Nat Rev Neurol* 2012; **8**: 329–39.
- 142 Hawkes CH. Olfaction in neurodegenerative disorder. *Mov Disord* 2003; **18**: 364–72.
- 143 Ponsen MM, Stoffers D, Boij J, van Eck-Smit BL, Wolters EC, Berendse HW. Idiopathic hyposmia as a preclinical sign of Parkinson's disease. *Ann Neurol* 2004; **56**: 173–81.
- 144 Haehner A, Hummel T, Hummel C, Sommer U, Jughannhans S, Reichmann H. Olfactory loss may be a first sign of idiopathic Parkinson's disease. *Mov Disord* 2007; **30**: 839–42.
- 145 Korczyn AD. Autonomic nervous system disturbances in Parkinson's disease. *Adv Neurol* 1990; **53**: 463–8.
- 146 Edwards LI, Pfeiffer RF, Quigley EM, Hofman R, Balluff M. Gastrointestinal symptoms in Parkinson's disease. *Mov Disord* 1991; **6**: 151–6.
- 147 Abbot RD, Petrovitch H, White LR, Masaki KH, Tanner CM, Curb JD, Grandinetti A, Blanchette PL, Popper JS, Ross GW. Frequency of bowel movements and the future risk of Parkinson's disease. *Neurology* 2001; **57**: 456–62.
- 148 Chaudhuri KR, Healy DG, Schapira AH; National Institute for Clinical Excellence. Non-motor symptoms of Parkinson's disease: diagnosis and management. *Lancet Neurol* 2006; **5**: 235–45.
- 149 Poewe W. Non-motor symptoms in Parkinson's disease. *Eur J Neurol* 2008; **15**(Suppl 1): 14–20.
- 150 Lang AE. A critical appraisal of the premotor symptoms of Parkinson's disease: potential usefulness in early diagnosis and design of neuroprotective trials. *Mov Disord* 2011; **26**: 775–83.
- 151 Pfeiffer RF. Gastrointestinal dysfunction in Parkinson's disease. *Parkinsonism Rel Disord* 2011; **17**: 10–5.
- 152 Cersósimo MG, Benarroch EE. Autonomic involvement in Parkinson's disease: pathology pathophysiology, clinical features and possible peripheral biomarkers. *J Neurol Sci* 2012; **313**: 57–63.
- 153 Cersósimo MG, Perandonas C, Micheli FE, Raina GB, Beron AM, Nassetter G, Radrizanni M, Benarroch EE. Alpha-synuclein immunoreactivity in minor salivary

- gland biopsies of Parkinson's disease patients. *Mov Disord* 2011; **26**: 188–90.
- 154 Postuma RB, Aarsland D, Barone P, Burn DJ, Hawkes CH, Oertel W, Ziemssen T. Identifying prodromal Parkinson's disease: pre-motor disorders in Parkinson's disease. *Mov Disord* 2012; **27**: 617–26.
 - 155 Khoo TK, Yarnall AJ, Duncan GW, Coleman S, O'Brien JT, Brooks RA, Burn DJ. The spectrum of nonmotor symptoms in early Parkinson disease. *Neurology* 2013; **80**: 276–81.
 - 156 Hilton D, Stephens M, Kirk L, Edwards P, Potter R, Zajicek J, Broughton E, Hagan H, Carroll C. Accumulation of α -synuclein in the bowel of patients in the pre-clinical phase of Parkinson's disease. *Acta Neuropathol* 2014; **127**: 235–41.
 - 157 Stiasny-Kolster K, Doerr Y, Möller JC, Höffken H, Behr TM, Oertel WH, Mayer G. Combination of 'idiopathic' REM sleep behaviour disorder and olfactory dysfunction as possible indicator for alpha-synucleinopathy demonstrated by dopamine transporter FP-CIT-SPECT. *Brain* 2005; **128**: 126–37.
 - 158 Boeve BF, Saper CB. REM sleep behavior disorder: a possible early marker for synucleinopathies. *Neurology* 2006; **28**: 796–7.
 - 159 Korczyn A, Gurevich T. Parkinson's disease: before the motor symptoms and beyond. *J Neurol Sci* 2010; **289**: 2–6.
 - 160 Hurtig HI, Trojanowski JQ, Galvin J, Ewbank D, Schmidt ML, Lee VM, Clark CM, Glosser G, Stern MB, Gollomp SM, Arnold SE. Alpha-synuclein cortical Lewy bodies correlate with dementia in Parkinson's disease. *Neurology* 2000; **54**: 1916–21.
 - 161 Apaydin H, Ahlskog J, Parisi J, Boeve B, Dickson D. Parkinson's disease neuropathology: later-developing dementia and loss of the levodopa response. *Arch Neurol* 2002; **59**: 102–12.
 - 162 Hely MA, Reid WG, Adena MA, Halliday GM, Morris JG. The Sydney multicenter study of Parkinson's disease: the inevitability of dementia at 20 years. *Mov Disord* 2008; **23**: 837–44.
 - 163 Sabbagh MN, Adler CH, Lahti TJ, Connor DJ, Vedders L, Peterson LK, Caviness JN, Shill HA, Sue LI, Ziabreva I, Perry E, Ballard CG, Aarsland D, Walker DG, Beach TG. Parkinson's disease with dementia: comparing patients with and without Alzheimer pathology. *Alzheimer Dis Assoc Disord* 2009; **23**: 295–7.
 - 164 Reid WG, Hely MA, Morris JG, Loy C, Halliday GM. Dementia in Parkinson's disease: a 20-year neuropsychological study (Sydney Multicentre Study). *J Neurol Neurosurg Psychiatry* 2011; **82**: 1033–7.
 - 165 Irwin DJ, White MT, Toledo JB, Xie SX, Robinson JL, Van Deerlin V, Lee VM, Leverenz JB, Montine TJ, Duda JE, Hurtig HI, Trojanowski JQ. Neuropathologic substrates of Parkinson disease dementia. *Ann Neurol* 2012; **72**: 587–98.
 - 166 Pinter B, Diem-Zangert A, Wenning GK, Scherfler C, Oberger W, Seppi K, Poewe W. Mortality in Parkinson's disease: a 38-year follow-up study. *Mov Disord* 2015; **30**: 266–9.
 - 167 Hawkes CH, Del Tredici K, Braak H. Parkinson's disease: a dual-hit hypothesis. *Neuropathol Appl Neurobiol* 2007; **33**: 599–614.
 - 168 Hawkes CH, Del Tredici K, Braak H. Parkinson's disease: the dual hit theory revisited. *Ann N Y Acad Sci* 2009; **1170**: 615–22.
 - 169 Gaig C, Tolosa E. When does Parkinson's disease begin? *Mov Disord* 2009; **24**(Suppl 2): S656–S664.
 - 170 Halliday GM, Holton JL, Revesz R, Dickson DW. Neuropathology underlying clinical variability in patients with synucleinopathies. *Acta Neuropathol* 2011; **122**: 187–204.
 - 171 van de Berg WD, Hepp DH, Dijkstra AA, Rozemuller JA, Berendse HW, Foncke E. Patterns of alpha-synuclein pathology in incidental cases and clinical subtypes of Parkinson's disease. *Parkinsonism Relat Disord* 2012; **18**(Suppl 1): S28–S30.
 - 172 Jenner P, Morris HR, Robbins TW, Goedert M, Hardy H, Ben-Schlomo Y, Bolam P, Burn D. Parkinson's disease: the debate on the clinical phenomenology, aetiology, pathology and pathogenesis. *J Parkinsons Dis* 2013; **3**: 1–11.
 - 173 Braak H, Ghebremedhin E, Rüb U, Bratzke H, Del Tredici K. Stages in the development of Parkinson's disease-related pathology. *Cell Tiss Res* 2004; **318**: 121–34.
 - 174 Braak H, Müller CM, Bohl JR, Rüb U, de Vos RAI, Del Tredici K. Stanley Fahn Lecture 2005: the staging procedure for the inclusion body pathology associated with sporadic Parkinson disease reconsidered. *Mov Disord* 2006; **21**: 2042–51.
 - 175 Sengoku R, Saito Y, Ikemura M, Hatsuta H, Sakiyama Y, Kanemaru K, Arai T, Sawabe M, Tanaka N, Mochizuki H, Inoue K, Shigeo S. Incidence and extent of Lewy body-related alpha-synucleinopathy in aging human olfactory bulb. *J Neuropathol Exp Neurol* 2008; **67**: 1072–83.
 - 176 Dickson DW, Uchikado H, Fujishiro H, Tsuboi Y. Evidence in favor of Braak staging of Parkinson's disease. *Mov Disord* 2010; **25**(Suppl 1): S78–S82.
 - 177 Halliday GM, McCann H, Shepherd C. Evaluation of the Braak hypothesis: how far can it explain the pathogenesis of Parkinson's disease? *Exp Rev Neurother* 2012; **12**: 673–86.
 - 178 Uchihara T, Giasson BI. Propagation of alpha-synuclein pathology: hypotheses, discoveries, and yet unresolved questions from experimental and human studies. *Acta Neuropathol* 2016; **131**: 49–73.
 - 179 Del Tredici K, Rüb U, de Vos RA, Bohl JR, Braak H. Where does Parkinson disease pathology begin in the brain? *J Neuropathol Exp Neurol* 2002; **61**: 413–26.

- 180 Pearce RK, Hawkes CH, Daniel SE. The anterior olfactory nucleus in Parkinson's disease. *Mov Disord* 1995; **10**: 283–7.
- 181 Duda JE, Shah U, Arnold SE, Lee VM, Trojanowski JQ. The expression of alpha-, beta-, and gamma-synucleins in olfactory mucosa. *Exp Neurol* 1999; **60**: 515–22.
- 182 Watelet JR, Strolin-Benedetti M, Whomsley R. Defense mechanisms of olfactory neuro-epithelium: mucosa regeneration, metabolising enzymes and transporters. *B-ENT* 2009; **5**(Suppl 13): 21–37.
- 183 Beach TG, Adler CH, Lue LI, Bachalakuri J, Henry-Watson J, Sasse J, Boyer S, Shirohi S, Brooks R, Eschbacher J, White CL III, Akiyama H, Caviness J, Shill HA, Connor DJ, Sabbagh MN, Walker DG. Unified staging system for Lewy body disorders: correlation with nigrostriatal degeneration, cognitive impairment and motor dysfunction. *Acta Neuropathol* 2009; **117**: 613–34.
- 184 Baker H, Genter MB. The olfactory system and the nasal mucosa as portals of entry of viruses, drugs, and other exogenous agents into the brain. In *Handbook of Olfaction and Gustation*. Ed Doty RL. New York: Marcel Dekker, 2003; 549–73.
- 185 Hubbard PS, Esiri MM, Reading M, McShane R, Nagy Z. Alpha-synuclein pathology in the olfactory pathways of dementia patients. *J Anat* 2007; **211**: 117–24.
- 186 Doty RL. The olfactory vector hypothesis of neurodegenerative disease: is it viable? *Ann Neurol* 2008; **63**: 7–15.
- 187 Lerner A, Bagic A. Olfactory pathogenesis of idiopathic Parkinson's disease revisited. *Mov Disord* 2008; **23**: 1076–84.
- 188 Rey NL, Petit GH, Bousset L, Melki R, Brundin P. Transfer of human α -synuclein from the olfactory bulb to interconnected brain regions in mice. *Acta Neuropathol* 2013; **126**: 555–73.
- 189 Forsyth CB, Shannon KM, Kordower JH, Voigt RM, Shaikh M, Jaglin JA, Estes JD, Dodiya HB, Keshavarzian A. Increased intestinal permeability correlates with sigmoid mucosa alpha-synuclein staining and endotoxin exposure markers in early Parkinson's disease. *PLoS ONE* 2011; **6**: e28032.
- 190 Kuo YM, Li Z, Jiao Y, Gaborit N, Pani AK, Orison BM, Bruneau BG, Giasson BI, Smeyne RJ, Gershon MD, Nussbaum RL. Extensive enteric nervous system abnormalities in mice transgenic for artificial chromosomes containing Parkinson disease-associated alpha-synuclein gene mutations precede central nervous system changes. *Hum Mol Genet* 2010; **19**: 1633–50.
- 191 De Chiara G, Marcocci ME, Sgarbanti R, Civitelli L, Ripoli C, Piacentini R, Garaci E, Grassi C, Palamara AT. Infectious agents in neurodegeneration. *Mol Neurobiol* 2012; **46**: 614–38.
- 192 Benarroch EE. The locus ceruleus norepinephrine system. Functional organization and potential clinical significance. *Neurology* 2009; **17**: 1699–704.
- 193 Nieuwenhuys R. The greater limbic system, the emotional motor system and the brain. *Prog Brain Res* 1996; **107**: 551–80.
- 194 Nieuwenhuys R, Voogd J, van Huijzen C. *The Human Central Nervous System*, 4th edn. Berlin, Heidelberg: Springer, 2008.
- 195 Braak H, Rüb U, Sandmann-Keil D, Gai WP, de Vos RA, Jansen Steur EN, Arai K, Braak E. Parkinson's disease: affection of brain stem nuclei controlling premotor and motor neurons of the somatomotor system. *Acta Neuropathol* 2000; **99**: 489–95.
- 196 Sumikura H, Takao M, Hatsuta H, Ito S, Uchino A, Nogami A, Saito Y, Mochizuki H, Murayama S. Distribution of α -synuclein in the spinal cord and dorsal root ganglia in an autopsy cohort of elderly persons. *Acta Neuropathol Comm* 2015; **3**: 57.
- 197 VanderHorst V, Samardzic T, Saper CB, Anderson MP, Nag S, Schneider JA, Bennett DA, Buchman AS. α -Synuclein pathology accumulates in sacral spinal visceral sensory pathways. *Ann Neurol* 2015; **78**: 142–9.
- 198 Braak H, Sastre M, Bohl JRE, de Vos RA, Del Tredici K. Parkinson's disease: lesions in dorsal horn layer I, involvement of parasympathetic and sympathetic pre- and postganglionic neurons. *Acta Neuropathol* 2007; **113**: 421–9.
- 199 Pan-Montojo F, Anichtchik O, Dening Y, Knels L, Pursche S, Jung R, Jackson S, Gille G, Spillantini MG, Reichmann H, Funk RHW. Progression of Parkinson's disease pathology is reproduced by intragastric administration of rotenone in mice. *PLoS ONE* 2010; **5**: e8762.
- 200 Sims KS, Williams RS. The human amygdaloid complex: a cytologic and histochemical atlas using Nissl, myelin, acetylcholinesterase and nicotinamide adenine dinucleotide phosphate diaphorase staining. *Neuroscience* 1990; **36**: 449–72.
- 201 Liubashina O, Jolkonen E, Pitkänen A. Projections from the central nucleus of the amygdala to the gastric related area of the dorsal vagal complex: a Phaseolus vulgaris leucoagglutinin study in rat. *Neurosci Lett* 2000; **291**: 85–8.
- 202 Pahapill PA, Lozano AM. The pedunculopontine nucleus and Parkinson's disease. *Brain* 2000; **123**: 1767–83.
- 203 Collingridge GL, James TA, MacLeod NK. Neurochemical and electrophysiological evidence for a projection from the locus coeruleus to the substantia nigra. *J Physiol* 1979; **290**: 44.
- 204 Fudge JL, Haber SN. The central nucleus of the amygdala projection to dopamine subpopulations in primates. *Neuroscience* 2000; **97**: 479–94.
- 205 Strack AM, Sawyer WB, Hughes JH, Platt KB, Loewy AD. A general pattern of CNS innervation of the sym-

- thetic outflow demonstrated by transneuronal pseudorabies viral infections. *Brain Res* 1989; **491**: 156–62.
- 206 Morrison LA, Sidman RL, Fields BN. Direct spread of reovirus from the intestinal lumen to the central nervous system through vagal autonomic nerve fibers. *Proc Natl Acad Sci USA* 1991; **88**: 3852–6.
 - 207 Kristensson K. Potential role of viruses in neurodegeneration. *Mol Chem Neuropathol* 1992; **16**: 45–58.
 - 208 Card JP. Exploring brain circuitry with neurotropic viruses: new horizons in neuroanatomy. *Anat Rec* 1998; **253**: 176–85.
 - 209 Rinaman L, Levitt P, Card JP. Progressive postnatal assembly of limbic-autonomic circuits revealed by central transneuronal transport of pseudorabies virus. *J Neurosci* 2000; **20**: 2731–41.
 - 210 Jang H, Boltz D, Sturm-Ramirez K, Shepherd KR, Jiao Y, Webster R, Smeyne RJ. Highly pathogenic H5N1 influenza virus can enter the central nervous system and induce neuroinflammation and neurodegeneration. *Proc Natl Acad Sci USA* 2009; **106**: 14063–8.
 - 211 McCann H, Cartwright H, Halliday GM. Neuropathology of α -synuclein propagation and Braak hypothesis. *Mov Disord* 2015; doi: 10.1002/mds.26421.
 - 212 Braak H, Braak E. The human entorhinal cortex: normal morphology and lamina-specific pathology in various diseases. *Neurosci Res* 1992; **15**: 6–31.
 - 213 Braak H, Del Tredici K. The preclinical phase of the pathological process underlying sporadic Alzheimer's disease. *Brain* 2015; **138**: 2814–33.
 - 214 Gloor P. *The Temporal Lobe and Limbic System*. New York: Oxford University Press, 1997.
 - 215 Amunts K, Kedo O, Kindler M, Pieperhoff P, Mohlbach H, Shah NJ, Habel U, Schneider F, Zilles K. Cytoarchitectonic mapping of the human amygdala, hippocampal region and entorhinal cortex: intersubject variability and probability maps. *Anat Embryol* 2005; **210**: 343–52.
 - 216 Amunts K, Zilles K. Advances in cytoarchitectonic mapping of the human cerebral cortex. *Neuroimaging Clin N Am* 2001; **11**: 151–69.
 - 217 Wakabayashi K, Hansen LA, Masliah E. Cortical Lewy body-containing neurons are pyramidal cells: laser confocal imaging of double-immunolabeled sections with anti-ubiquitin and SMI32. *Acta Neuropathol* 1995; **89**: 404–8.
 - 218 Barbas H, Rempel-Clower N. Cortical structure predicts the pattern of corticocortical connections. *Cereb Cortex* 1997; **7**: 635–46.
 - 219 Bannister AP. Inter- and intra-laminar connections of pyramidal cells in the neocortex. *Neurosci Res* 2005; **53**: 95–103.
 - 220 Barbas H. Specialized elements of orbitofrontal cortex in primates. *Ann N Y Acad Sci* 2007; **1121**: 10–32.
 - 221 Braak H, Rüb U, Gai WP, Del Tredici K. Idiopathic Parkinson's disease: possible routes by which vulnerable neuronal types may be subject to neuroinvasion by an unknown pathogen. *J Neural Transm* 2003; **110**: 517–36.
 - 222 Reichmann H. View point: etiology in Parkinson's disease. Dual hit or spreading intoxication. *J Neurol Sci* 2011; **310**: 9–11.
 - 223 Pan-Montojo F, Reichmann H. Considerations on the role of environmental toxins in idiopathic Parkinson's disease pathophysiology. *Transl Neurodegener* 2014; **3**: 10.
 - 224 Nakamura K, Mori F, Tanji K, Miki Y, Toyoshima Y, Kakita A, Takahashi H, Yamada M, Wakabayashi K. α -Synuclein pathology in cranial and spinal nerves in Lewy body disease. *Neuropathology* 2015; doi: 10.1111/neup.12269.
 - 225 Richards WG, Sugarbaker DJ. Neuronal control of esophageal function. *Chest Surg Clin N Am* 1995; **5**: 157–71.
 - 226 Benarroch EE. Enteric nervous system: functional organization and neurologic implications. *Neurology* 2007; **69**: 1953–7.
 - 227 Cersósimo MG, Benarroch EE. Neural control of the gastrointestinal tract: implications for Parkinson disease. *Mov Disord* 2008; **23**: 1065–75.
 - 228 Fuminura Y, Ikemura M, Saito Y, Sengoku R, Kanemaru K, Sawabe M, Arai T, Ito G, Iwatsubo T, Fukayama M, Mizusawa H, Murayama S. Analysis of the adrenal gland is useful for evaluating pathology of the peripheral autonomic nervous system in Lewy body disease. *J Neuropathol Exp Neurol* 2007; **66**: 354–62.
 - 229 Miki Y, Mori F, Wakabayashi K, Kuroda N, Orimo S. Incidental Lewy body disease restricted to the heart and stellate ganglia. *Mov Disord* 2009; **24**: 2299–301.
 - 230 Desplats P, Lee H-J, Bae E-J, Patrick C, Rockenstein E, Crews L, Spencer B, Masliah E, Lee S-J. Inclusion formation and neuronal cell death through neuron-to-neuron transmission of α -synuclein. *Proc Natl Acad Sci USA* 2009; **106**: 13010–5.
 - 231 Hansen C, Angot E, Bergström AL, Steiner JA, Pieri L, Paul G, Outeiro TF, Melki R, Kallunki P, Fog K, Li JY, Brundin P. α -Synuclein propagates from mouse brain to grafted dopaminergic neurons and seeds aggregation in cultured human cells. *J Clin Invest* 2011; **121**: 715–25.
 - 232 Miwa H, Kubo T, Suzuki A, Kondo T. Intragastric proteasome inhibition induces alpha-synuclein-immunopositive aggregations in neurons in the dorsal motor nucleus of the vagus in rats. *Neurosci Lett* 2006; **401**: 146–9.
 - 233 Ulusoy A, Rusconi R, Perez-Revuelta BI, Musgrove RE, Helwig M, Winzen-Reichert B, Di Monte DA. Caudorostral brain spreading of alpha synuclein through vagal connections. *EMBO Mol Med* 2013; **5**: 1051–9.
 - 234 Holmqvist S, Chutna O, Bousset L, Aldrin-Kirk P, Li W, Björklund T, Wang ZY, Roybon L, Melki R, Li J-Y. Direct evidence of Parkinson pathology spread from

- the gastrointestinal tract to the brain in rats. *Acta Neuropathol* 2014; **128**: 805–20.
- 235 Sacino AN, Brooks J, Thomas MA, McKinney AB, Lee S, Regenhardt RW, McGarvey NH, Ayers JL, Notterpek L, Borchelt DR, Golde TE, Giasson BI. Intramuscular injection of α -synuclein induces CNS α -synuclein pathology and a rapid-onset motor phenotype in transgenic mice. *Proc Natl Acad Sci USA* 2014; **111**: 10732–7.
 - 236 Phillips RJ, Walter GC, Wilder SL, Baronowski EA, Powley TL. Alpha-synuclein immunopositive myenteric neurons and vagal preganglionic terminals: autonomic pathway implicated in Parkinson's disease? *Neuroscience* 2008; **153**: 733–50.
 - 237 Norrian AR, Rha J, Annerino DM, Bernhard D, Taylor GM, Greene JG. Alpha-synuclein transgenic mice display age-related slowing of gastrointestinal motility associated with transgene expression in the vagal system. *Neurobiol Dis* 2012; **48**: 9–19.
 - 238 Svensson E, Horváth-Puhó E, Thomsen RW, Djurhuus JC, Pedersen L, Borghammer P, Sørensen HT. Vagotomy and subsequent risk of Parkinson's disease. *Ann Neurol* 2015; **78**: 522–9.
 - 239 Hopkins DA, Bieger D, de Vente J, Steinbusch HWM. Vagal efferent projections: viscerotopy, neurochemistry and effects of vagotomy. *Progr Brain Res* 1996; **107**: 79–96.
 - 240 Pan-Montojo F, Schwarz M, Winkler C, Arnhold M, O'Sullivan GA, Pal A, Said J, Marsico G, Verbavatz JM, Rodrigo-Angulo M, Gille G, Funk RH, Reichmann H. Environmental toxins trigger PD-like progression via increased alpha-synuclein release from enteric neurons in mice. *Sci Rep* 2012; **2**: 898.
 - 241 Saper CB, Wainer BH, German DC. Axonal and transneuronal transport in the transmission of neurological disease: potential role in system degenerations, including Alzheimer's disease. *Neuroscience* 1987; **23**: 389–98.
 - 242 Brundin P, Li JY, Holton JL, Lindvall O, Revesz T. Research in motion: the enigma of Parkinson's disease pathology spread. *Nat Rev Neurosci* 2008; **9**: 741–5.
 - 243 Angot E, Steiner JA, Hansen C, Li JY, Brundin P. Are synucleinopathies prion-like disorders? *Lancet Neurol* 2010; **9**: 1128–38.
 - 244 Goedert M, Clavaguera F, Tolnay M. The propagation of prion-like protein inclusions in neurodegenerative diseases. *Trends Neurosci* 2010; **33**: 317–25.
 - 245 Goedert M, Falcon B, Clavaguera F, Tolnay M. Prion-like mechanisms in the pathogenesis of tauopathies and synucleinopathies. *Curr Neurol Neurosci Rep* 2014; **14**: 495.
 - 246 Steiner JA, Angot E, Brundin P. A deadly spread: cellular mechanisms of α -synuclein transfer. *Cell Death Differ* 2011; **18**: 1425–33.
 - 247 Danzer KM, Kranich LR, Ruf WP, Cagsal-Getkin O, Winslow AR, Zhu L, Vanderburg CR, McLean PJ. Exosomal cell-to-cell transmission of alpha synuclein oligomers. *Mol Neurodegener* 2012; **7**: 42.
 - 248 Dunning CJ, Reyes JF, Steiner JA, Brundin P. Can Parkinson's disease pathology be propagated from neuron to another? *Prog Neurobiol* 2012; **97**: 205–19.
 - 249 Hansen C, Li JY. Beyond alpha-synuclein transfer: pathology propagation in Parkinson's disease. *Trends Mol Med* 2012; **18**: 248–55.
 - 250 Bousset L, Pieri L, Ruiz-Arlandis G, Gath J, Jensen PH, Habenstein B, Madiona K, Olieric V, Böckmann A, Meier BH, Melki R. Structural and functional characterization of two alpha-synuclein strains. *Nat Commun* 2013; **4**: 2575.
 - 251 George S, Rey NL, Reichenbach N, Steiner JA, Brundin P. α -Synuclein; the long distance runner. *Brain Pathol* 2013; **23**: 350–7.
 - 252 Jucker M, Walker LC. Self-propagation of pathogenic protein aggregates in neurodegenerative diseases. *Nature* 2013; **501**: 45–51.
 - 253 Olanow CW, Brundin P. Parkinson's disease and alpha synuclein: is Parkinson's disease a prion-like disorder? *Mov Disord* 2013; **28**: 31–40.
 - 254 Sacino AN, Giasson BI. Does a prion-like mechanism play a major role in the apparent spread of α -synuclein pathology? *Alzheimers Res Ther* 2013; **4**: 48.
 - 255 Luk KC, Lee VM. Modeling Lewy body propagation in Parkinson's disease. *Parkinsonism Related Disord* 2014; **20 Suppl 1**: S85–S87.
 - 256 Recasens A, Dehay B. Alpha-synuclein spreading in Parkinson's disease. *Front Neuroanat* 2014; **8**: 1–9.
 - 257 Tran HT, Chung CH, Iba M, Zhang B, Trojanowski JQ, Luk KC, Lee VM. A-synuclein immunotherapy blocks uptake and templated propagation of misfolded α -synuclein and neurodegeneration. *Cell Rep* 2014; **7**: 2054–65.
 - 258 Herva ME, Spillantini MG. Parkinson's disease as a member of prion-like disorders. *Virus Res* 2015; **207**: 38–46.
 - 259 Melki R. Role of different alpha-synuclein strains in synucleinopathies, similarities with other neurodegenerative diseases. *J Parkinson's Dis* 2015; **5**: 217–27.
 - 260 Peelaerts W, Bousset L, Van der Peren A, Moskalyuk A, Pulizzi R, Giugliano M, Van den Haute C, Melki R, Baekelandt V. α -Synuclein strains cause distinct synucleinopathies after local and systemic administration. *Nature* 2015; **522**: 340–4.
 - 261 Kamel F, Goldman SM, Umbach DM, Chen H, Richardson G, Barber RM, Meng C, Marras C, Korell M, Kasten M, Hoppin JA, Comyns K, Chade A, Blair A, Bhudhikanok GS, Webster Ross G, Langston JW, Sandler DP, Tanner CM. Dietary fat intake, pesticide use, and Parkinson's disease. *Parkinsonism Relat Disord* 2015; **20**: 82–7.

- 262 Büttner T, Patzold T, Kuhn W, Müller T, Przuntek H. Impaired color discrimination in Parkinson's disease. *Neuroophthalmol* 1994; **14**: 91–5.
- 263 Haug BA, Kolle RU, Trenkwalder C, Oertel WH, Paulus W. Predominant affection of the blue cone pathway in Parkinson's disease. *Brain* 1995; **118**: 771–8.
- 264 Nolano M, Provitera V, Estraneo A, Selim MM, Caa-poraso G, Stancanelli A, Saltalamacchia AM, Lanzillo B, Santoro L. Sensory deficit in Parkinson's disease: evidence of a cutaneous denervation. *Brain* 2008; **131**: 1903–11.
- 265 Doppler K, Ebert S, Uçeyler N, Trenkwalder C, Ebentheuer J, Volkmann J, Sommer C. Cutaneous neuropathy in Parkinson's disease: a window into brain pathology. *Acta Neuropathol* 2014; **128**: 99–109.
- 266 Kaur M, Saxena R, Singh D, Behari M, Sharma P, Menon V. Correlation between structural and functional retinal changes in Parkinson disease. *J Neuroophthalmol* 2015; **35**: 254–8.
- 267 Mu L, Chen J, Sobotka S, Nyirenda T, Benson B, Gupta F, Sanders I, Adler CH, Caviness JN, Shill HA, Sabbagh M, Samanta JE, Sue LI, Beach TG; Arizona Parkinson's Disease Consortium. Alpha-synuclein pathology in sensory nerve terminals of the upper aerodigestive tract of Parkinson's disease patients. *Dysphagia* 2015; **30**: 404–17.
- 268 Costa M, Brookes SJ, Hennig GW. Anatomy and physiology of the enteric nervous system. *Gut* 2000; **47**(Suppl 4): 15–19.
- 269 Furness JB. Types of neurons in the enteric nervous system. *J Auton Nerv Syst* 2000; **81**: 87–96.
- 270 Aldecoa I, Navarro-Otano J, Stefanova N, Sprenger FS, Seppi K, Poewe W, Cuatrecasas M, Valdeoriola F, Gelpi E, Tolosa E. Alpha-synuclein immunoreactivity patterns in the enteric nervous system. *Neurosci Lett* 2015; **602**: 145–9.

Received 18 October 2015

Accepted after revision 4 December 2015

Published online Article Accepted on 13 December 2015