


*Use of **G**eographic **I**nformation **S**ystems in Crop Protection Warning Service in Germany*

3rd Conference on PRECISION CROP PROTECTION
Bonn, September 19 - 21 2010

Zeuner & Kleinhenz

- Introduction
- Spatial input data
 - Temperature and relative humidity
 - Precipitation
- Risk maps
- Summary & Conclusion

Structure of ZEPP



Aims in Crop Protection Warning Service by ZEPP

Improvement of Decision Support Systems (DSS)

- ➔ Forecasting the first appearance of plant diseases
- ➔ Identifying areas of maximum and minimum infection risk
- ➔ Giving best control for pests and diseases
- ➔ Optimising spraying strategy
- ➔ Reducing the fungicide intensity

Models

in practical use	22
<u>under development</u>	<u>18</u>
total models	40

Use of internet warning service

met. data



- temperature
- relative humidity
- precipitation

field data

- cultivar
- plant emergence
- crop rotation
- geographic data
- etc.



Forecasting models




(www.isip.de)

Technical advance

GIS

ZEPP




- Introduction
- **Spatial input data**
 - Temperature and relative humidity
 - Precipitation
- Risk maps
- Summary & Conclusion

Storage of spatial met. data



- virtual weather database
- grid of 1 km²
- virtual met. stations
- Germany: 357 050 km²
- 360 000 virtual stations




3 met. stations against 1367 virtual stations on this map

- Introduction
- **Spatial input data**
 - **Temperature and relative humidity**
 - Precipitation
- Risk maps
- Summary & Conclusion


Interpolation of temperature and relative humidity

Deterministic interpolation methods

IDW next neighbor method




spline




Geostatistic interpolations methods


kriging



multiple regression



Interpolation with multiple regression (MR)



best correlations with altitude, longitude and elevation

Validation of model results



Validation of the method


year	days/ Year	hours/ Day	stations	temperature	humidity	n
2004	366	24	676	x	x	11.875.968
2005	365	24	676	x	x	11.843.520
2006	365	24	676	x	x	11.843.520
2007	365	24	676	x	x	11.843.520
2008	366	24	676	x	x	11.875.968
alle						59.282.496

Number of Boxplots


year	stations	parameter	whole year	spring/ summer	April	May	Juni	Juli	August	frequence	Boxplots
2004	676	2	x	x	x	x	x	x	x	3	28.392
2005	676	2	x	x	x	x	x	x	x	3	28.392
2006	676	2	x	x	x	x	x	x	x	3	28.392
2007	676	2	x	x	x	x	x	x	x	3	28.392
2008	676	2	x	x	x	x	x	x	x	3	28.392
alle											141.960

Validation of model results

Temperatur




relative Luftfeuchte



- similar results for other Boxplots
- Statistic check (T-Test): not significant
- useful as input for forecasting models

- Introduction
- **Spatial input data**
 - Temperature and relative humidity
 - **Precipitation**
- Risk maps
- Summary & Conclusion

One hour with precipitation



Spatial precipitation with radar data

- Spatial measured precipitation data is collected by the German Met. Service
- 16 radar stations measure the data nationwide over Germany
- The radar data is calibrated by 1600 classic automatic precipitation stations (Ombrometer) which are located over the whole area of Germany
- The radar data is collected in a frequent of one hour and in a spatial resolution of 1km²

Ombrometer




+

radar station





=

precipitation map



Comparison between radar data and independently classically collected precipitation data




- Introduction
- Spatial input data
 - Temperature and relative humidity
 - Precipitation
- **Risk maps**
- Summary & Conclusion

Risk maps




spatial input data

temperature



relative humidity



precipitation



warning service


field specific input data
cultivar, plant emergence,
crop rotation, etc.

forecasting models



spatial result presented as
risk map

risk map presentation



Forecasting model runs with spatial input data




Two type of models used by ZEPP

First Type

plant disease outbreak


- date of first monitoring
- date of first treatment



Second Type


plant disease infection risk

- high/low risk situations
- spraying strategy




First type: Risk maps of late blight with SIMBLIGHT1

Forecast of the outbreak of potato late blight in Germany



Second type: Risk maps of infection pressure with SIMPHYT3

- The output of SIMPHYt3 is a daily met. data based infection pressure for late blight by checking at met. data of the last 14 days



- Introduction
- Spatial input data
 - Temperature and relative humidity
 - Precipitation
- Risk maps
- **Summary & Conclusion**

- Virtual met. data were produced and tested on validity
 - interpolation method for temperature and relative humidity
 - radar data for precipitation
- Virtual met. data were used as input for forecasting models to produce risk maps
 - risk maps show hot spots of disease outbreak
 - risk maps show daily infection pressure for plant diseases
- Forecasting results as risk maps are easier to understand and to interpret
- GIS helps to obtain more detailed calculations and results with higher accuracy than before
- Risk maps are available under www.isip.de since April 2010

**This leads to a reduced pesticide use and
an economical and environmental friendly crop protection strategy**



Thanks for your attention

SIMPHYT3

2007