

Coalgebraic Representation of Büchi Automata

Research Internship Presentation

Jorrit de Boer


17 January 2025

Outline:


1. Büchi Automata
2. Coalgebraic Representation Deterministic Finite Automata
3. Coalgebraic Representation Nondeterministic Finite Automata
4. Coalgebraic Representation Possibly Infinite Behavior Nondeterministic Finite Automata
5. Coalgebraic Representation Büchi Automata
6. Outline Derivation using Game Semantics

Büchi Automata

Büchi Automata




Büchi Automata



Definition: A (nondeterministic) *Büchi Automaton*

$A = \langle S, \Sigma, \delta, s_0, F \rangle$, where:

- S : finite set of states
- Σ : alphabet
- $s_0 \in S$: initial state
- $\delta : S \times \Sigma \rightarrow \mathcal{P}(S)$: transition function
- $F \subseteq S$: set of *final* (or *accepting*) states.




Definition: A (nondeterministic) *Büchi Automaton*

$A = \langle S, \Sigma, \delta, s_0, F \rangle$, where:

- S : finite set of states
- Σ : alphabet
- $s_0 \in S$: initial state
- $\delta : S \times \Sigma \rightarrow \mathcal{P}(S)$: transition function
- $F \subseteq S$: set of *final* (or *accepting*) states.

A *run* of A on an ω -word $w = \sigma_0\sigma_1\ldots \in \Sigma^\omega$ is an infinite sequence of states $s_0, s_1, \ldots \in S^\omega$ such that for all n , $s_{n+1} \in \delta(s_n, \sigma_n)$

A run is *accepted* if it visits F infinitely often.



Definition: A (nondeterministic) *Büchi Automaton*

$A = \langle S, \Sigma, \delta, s_0, F \rangle$, where:

- S : finite set of states
- Σ : alphabet
- $s_0 \in S$: initial state
- $\delta : S \times \Sigma \rightarrow \mathcal{P}(S)$: transition function
- $F \subseteq S$: set of *final* (or *accepting*) states.

A *run* of A on an ω -word $w = \sigma_0\sigma_1\ldots \in \Sigma^\omega$ is an infinite sequence of states $s_0, s_1, \ldots \in S^\omega$ such that for all n , $s_{n+1} \in \delta(s_n, \sigma_n)$

A run is *accepted* if it visits F infinitely often.

Accepted language:

$(\text{request} \cdot \text{process}^* \cdot \text{return})^\omega$

Coalgebra

Coalgebraic Representation Deterministic Finite Automata

$\langle S, \Sigma, \delta, o \rangle$ with states S , alphabet Σ , transition function $\delta : S \times \Sigma \rightarrow S$, $o : S \rightarrow 2$ ($2 = \{0, 1\}$). Can be represented by a coalgebra $\langle o, \delta \rangle : S \rightarrow 2 \times S^\Sigma$ for functor $FS = 2 \times S^\Sigma$

Coalgebraic Representation Deterministic Finite Automata

$\langle S, \Sigma, \delta, o \rangle$ with states S , alphabet Σ , transition function $\delta : S \times \Sigma \rightarrow S$, $o : S \rightarrow 2$ ($2 = \{0, 1\}$). Can be represented by a coalgebra $\langle o, \delta \rangle : S \rightarrow 2 \times S^\Sigma$ for functor $FS = 2 \times S^\Sigma$

The final coalgebra for F is $\langle e, d \rangle : 2^{\Sigma^*} \rightarrow 2 \times (2^{\Sigma^*})^\Sigma$. Where

- $e(L) = L(\varepsilon)$, i.e. $e(L) = 1$ iff $\varepsilon \in L$
- $d(L)(\sigma) = L_\sigma$ where $L_\sigma(w) = L(\sigma w)$ so $w \in d(L)(\sigma)$ iff $\sigma w \in L$

Coalgebraic Representation Deterministic Finite Automata

$\langle S, \Sigma, \delta, o \rangle$ with states S , alphabet Σ , transition function $\delta : S \times \Sigma \rightarrow S$, $o : S \rightarrow 2$ ($2 = \{0, 1\}$). Can be represented by a coalgebra $\langle o, \delta \rangle : S \rightarrow 2 \times S^\Sigma$ for functor $FS = 2 \times S^\Sigma$

The final coalgebra for F is $\langle e, d \rangle : 2^{\Sigma^*} \rightarrow 2 \times (2^{\Sigma^*})^\Sigma$. Where

- $e(L) = L(\varepsilon)$, i.e. $e(L) = 1$ iff $\varepsilon \in L$
- $d(L)(\sigma) = L_\sigma$ where $L_\sigma(w) = L(\sigma w)$ so $w \in d(L)(\sigma)$ iff $\sigma w \in L$


$$\begin{array}{ccc}
 2 \times S^\Sigma & \xrightarrow{\text{id} \times \text{beh}^\Sigma} & 2 \times (2^{\Sigma^*})^\Sigma \\
 \uparrow \langle o, \delta \rangle & & \uparrow \langle e, d \rangle \\
 S & \xrightarrow{\text{beh}} & 2^{\Sigma^*}
 \end{array}$$

Coalgebraic Representation Deterministic Finite Automata

$\langle S, \Sigma, \delta, o \rangle$ with states S , alphabet Σ , transition function $\delta : S \times \Sigma \rightarrow S$, $o : S \rightarrow 2$ ($2 = \{0, 1\}$). Can be represented by a coalgebra $\langle o, \delta \rangle : S \rightarrow 2 \times S^\Sigma$ for functor $FS = 2 \times S^\Sigma$

The final coalgebra for F is $\langle e, d \rangle : 2^{\Sigma^*} \rightarrow 2 \times (2^{\Sigma^*})^\Sigma$. Where

- $e(L) = L(\varepsilon)$, i.e. $e(L) = 1$ iff $\varepsilon \in L$
- $d(L)(\sigma) = L_\sigma$ where $L_\sigma(w) = L(\sigma w)$ so $w \in d(L)(\sigma)$ iff $\sigma w \in L$



Following the paths through the diagram we obtain:


- $\text{beh}(s)(\varepsilon) = e(\text{beh}(s)) = o(s)$, and
- $\text{beh}(s)(\sigma w) = \text{beh}(s)_a(w) = d(\text{beh}(s))(\sigma) = \text{beh}(\delta(s)(\sigma))(w)$,

Coalgebraic Representation Deterministic Finite Automata

$\langle S, \Sigma, \delta, o \rangle$ with states S , alphabet Σ , transition function $\delta : S \times \Sigma \rightarrow S$, $o : S \rightarrow 2$ ($2 = \{0, 1\}$). Can be represented by a coalgebra $\langle o, \delta \rangle : S \rightarrow 2 \times S^\Sigma$ for functor $FS = 2 \times S^\Sigma$

The final coalgebra for F is $\langle e, d \rangle : 2^{\Sigma^*} \rightarrow 2 \times (2^{\Sigma^*})^\Sigma$. Where

- $e(L) = L(\varepsilon)$, i.e. $e(L) = 1$ iff $\varepsilon \in L$
- $d(L)(\sigma) = L_\sigma$ where $L_\sigma(w) = L(\sigma w)$ so $w \in d(L)(\sigma)$ iff $\sigma w \in L$




Following the paths through the diagram we obtain:

- $\text{beh}(s)(\varepsilon) = e(\text{beh}(s)) = o(s)$, and
- $\text{beh}(s)(\sigma w) = \text{beh}(s)_a(w) = d(\text{beh}(s))(\sigma) = \text{beh}(\delta(s)(\sigma))(w)$,


So beh captures exactly the accepted language of the automaton!

Nondeterministic Finite Automata



Might be modeled by coalgebra $c : S \rightarrow 2 \times \mathcal{P}(\Sigma \times S)$.

Nondeterministic Finite Automata



Might be modeled by coalgebra $c : S \rightarrow 2 \times \mathcal{P}(\Sigma \times S)$.

A final coalgebra $z : Z \rightarrow 2 \times \mathcal{P}(\Sigma \times Z)$ cannot exist. Lambek's lemma says z would have to be an isomorphism, which would imply $Z \cong \mathcal{P}(Z)$

Kleisli Category of the monad \mathcal{P} :

A coalgebra $c : S \rightarrow \Sigma \times S$ in $\mathcal{Kl}(\mathcal{P})$ is $c : S \rightarrow \mathcal{P}(\Sigma \times S)$ in **Sets**.

Kleisli Category of the monad \mathcal{P} :

A coalgebra $c : S \rightarrow \Sigma \times S$ in $\mathcal{K}\ell(\mathcal{P})$ is $c : S \rightarrow \mathcal{P}(\Sigma \times S)$ in **Sets**. Concretely:

- $\eta_X : X \rightarrow \mathcal{P}(X)$: $\eta_X(x) = \{x\}$
- $\mu_X : \mathcal{P}(\mathcal{P}(X)) \rightarrow \mathcal{P}(X)$: $\mu_X(A) = \bigcup_{a \in A} a$.

For $f : X \rightarrow Y$, $\mathcal{P}(f) : \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$ by $\mathcal{P}(f)(A) = \{f(a) \mid a \in A\}$

- **objects**: the same as for **Sets**, sets
- **morphisms**: $f : X \rightarrow Y$ in $\mathcal{K}\ell(\mathcal{P})$ is $f : X \rightarrow \mathcal{P}(Y)$ in **Sets**.

For morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ in $\mathcal{K}\ell(\mathcal{P})$ we define

$$g \odot f := X \xrightarrow{f} \mathcal{P}(Y) \xrightarrow{\mathcal{P}(g)} \mathcal{P}(\mathcal{P}(Z)) \xrightarrow{\mu_Y} \mathcal{P}(Z)$$

Model NFA $\langle S, \Sigma, \delta, o \rangle$ by coalgebra $c : S \rightarrow 1 + \Sigma \times S$ for the functor $FS = 1 + \Sigma \times S$, which is $c : S \rightarrow \mathcal{P}(1 + \Sigma \times S)$ in **Sets**.

Lift functor F in **Sets** to \overline{F} in $\mathcal{K}\ell(\mathcal{P})$

Model NFA $\langle S, \Sigma, \delta, o \rangle$ by coalgebra $c : S \rightarrow 1 + \Sigma \times S$ for the functor $FS = 1 + \Sigma \times S$, which is $c : S \rightarrow \mathcal{P}(1 + \Sigma \times S)$ in **Sets**.

Lift functor F in **Sets** to \overline{F} in $\mathcal{K}\ell(\mathcal{P})$

Theorem [Hasuo, Jacobs, Sokolova 2007]: An initial algebra $\alpha : FA \rightarrow A$ for the functor F in **Sets** yields the final coalgebra for \overline{F} in $\mathcal{K}\ell(\mathcal{P})$:

$$(\eta_{FA} \circ \alpha^{-1}) : A \rightarrow \overline{F}A \text{ in } \mathcal{K}\ell(\mathcal{P})$$

The initial algebra for $FS = 1 + \Sigma \times S$ is $[\text{nil}, \text{cons}] : 1 + \Sigma \times \Sigma^* \rightarrow \Sigma^*$:

- $\text{nil}(\ast) = \varepsilon$ $\text{cons}(\sigma, w) = \sigma w$

Coalgebraic Representation Nondeterministic Automata

The initial algebra for $FS = 1 + \Sigma \times S$ is $[\text{nil}, \text{cons}] : 1 + \Sigma \times \Sigma^* \rightarrow \Sigma^*$:

- $\text{nil}(\ast) = \varepsilon$ $\text{cons}(\sigma, w) = \sigma w$

$$\begin{array}{ccc}
 & 1 + \Sigma \times \text{beh} & \\
 1 + \Sigma \times S & \xrightarrow{\quad\quad\quad} & 1 + \Sigma \times \Sigma^* \\
 \uparrow c & & \uparrow \eta_{1+\Sigma \times \Sigma^*} \circ [\text{nil}, \text{cons}]^{-1} \text{ in } \mathcal{K}\ell(\mathcal{P}). \\
 S & \xrightarrow{\quad\quad\quad \text{beh} \quad\quad\quad} & \Sigma^*
 \end{array}$$

Coalgebraic Representation Nondeterministic Automata

The initial algebra for $FS = 1 + \Sigma \times S$ is $[\text{nil}, \text{cons}] : 1 + \Sigma \times \Sigma^* \rightarrow \Sigma^*$:

- $\text{nil}(\ast) = \varepsilon$ $\text{cons}(\sigma, w) = \sigma w$

$$\begin{array}{ccc}
 & 1 + \Sigma \times \text{beh} & \\
 1 + \Sigma \times S & \xrightarrow{\quad\quad\quad} & 1 + \Sigma \times \Sigma^* \\
 \uparrow c & & \uparrow \eta_{1+\Sigma \times \Sigma^*} \circ [\text{nil}, \text{cons}]^{-1} \text{ in } \mathcal{K}\ell(\mathcal{P}). \\
 S & \xrightarrow{\quad\quad\quad \text{beh} \quad\quad\quad} & \Sigma^*
 \end{array}$$

$$\varepsilon \in \text{beh}(s) \iff \ast \in c(s) \iff \text{state } s \text{ is accepting}$$

$$\sigma w \in \text{beh}(s) \iff (\sigma, w) \in ((\Sigma \times \text{beh}) \circ c)(s) = \{(\sigma, \text{beh}(t)) \mid (\sigma, t) \in c(s)\} \iff \exists t. (t \in \delta(s)(\sigma) \wedge w \in \text{beh}(t)).$$

Theorem [Jacobs 2004]: A final coalgebra $\xi : Z \rightarrow FZ$ yields a *weakly final* coalgebra

$$(\eta_{FZ} \circ \xi) : Z \rightarrow \overline{F}(Z) \text{ in } \mathcal{K}\ell(\mathcal{P})$$

Theorem [Jacobs 2004]: A final coalgebra $\xi : Z \rightarrow FZ$ yields a *weakly final* coalgebra

$$(\eta_{FZ} \circ \xi) : Z \rightarrow \overline{F}(Z) \text{ in } \mathcal{K}\ell(\mathcal{P})$$

$$\begin{array}{ccc} \overline{F}S & \overset{\overline{F}(\text{beh})}{\rightsquigarrow} & \overline{F}Z \\ \uparrow c & & \uparrow \eta_{FZ} \circ \xi \\ S & \overset{\text{beh}}{\rightsquigarrow} & Z \end{array} \text{ in } \mathcal{K}\ell(\mathcal{P}),$$

beh is not unique. However, we can take beh^∞ , the maximal mapping with respect to inclusion.

$\xi : \Sigma^\infty \rightarrow 1 + \Sigma \times \Sigma^\infty$ is the final F -coalgebra, defined by $\xi(\varepsilon) = * \in 1$ and $\xi(\sigma w) = (\sigma, w)$
($\Sigma^\infty = \Sigma^* \cup \Sigma^\omega$).

Possibly Infinite Behavior

$\xi : \Sigma^\infty \rightarrow 1 + \Sigma \times \Sigma^\infty$ is the final F -coalgebra, defined by $\xi(\varepsilon) = * \in 1$ and $\xi(\sigma w) = (\sigma, w)$ ($\Sigma^\infty = \Sigma^* \cup \Sigma^\omega$).

$$\begin{array}{ccc}
 & 1 + \Sigma \times \text{beh}_c^\infty & \\
 1 + \Sigma \times S & \rightsquigarrow & 1 + \Sigma \times \Sigma^\infty \\
 \uparrow c & & \uparrow J\xi \\
 S & \rightsquigarrow_{\text{beh}_c^\infty} & \Sigma^\infty
 \end{array}
 \quad \text{in } \mathcal{K}\ell(\mathcal{P}).$$

$\varepsilon \in \text{beh}^\infty(s) \iff * \in c(s) \iff \text{state } s \text{ is accepting}$

$\sigma w \in \text{beh}^\infty(s) \iff \exists t. (s \xrightarrow{\sigma} t \wedge w \in \text{beh}^\infty(t)).$

Idea: split $S = S_1 \cup S_2$ for S_1 non-accepting and S_2 accepting

Idea: split $S = S_1 \cup S_2$ for S_1 non-accepting and S_2 accepting

$$\begin{array}{ccc}
 \Sigma \times [\text{beh}_1, \text{beh}_2] & & \Sigma \times [\text{beh}_1, \text{beh}_2] \\
 \Sigma \times S \rightsquigarrow \Sigma \times \Sigma^\omega & & \Sigma \times S \rightsquigarrow \Sigma \times \Sigma^\omega \\
 \uparrow c_1 & \overset{=}{\mu} & \uparrow c_2 \\
 S_1 \rightsquigarrow \Sigma^\omega & & S_2 \rightsquigarrow \Sigma^\omega \\
 \text{beh}_1 & & \text{beh}_2
 \end{array}
 \quad
 \begin{array}{ccc}
 \Sigma \times [\text{beh}_1, \text{beh}_2] & & \Sigma \times [\text{beh}_1, \text{beh}_2] \\
 \Sigma \times S \rightsquigarrow \Sigma \times \Sigma^\omega & & \Sigma \times S \rightsquigarrow \Sigma \times \Sigma^\omega \\
 \uparrow c_2 & \overset{=}{\nu} & \uparrow \eta_{\Sigma^\omega} \circ d \\
 S_2 \rightsquigarrow \Sigma^\omega & & \Sigma^\omega
 \end{array}
 \quad
 \text{in } \mathcal{K}\ell(\mathcal{P}).$$

Idea: split $S = S_1 \cup S_2$ for S_1 non-accepting and S_2 accepting

$$\begin{array}{ccc}
\Sigma \times [\text{beh}_1, \text{beh}_2] & & \Sigma \times [\text{beh}_1, \text{beh}_2] \\
\Sigma \times S \rightsquigarrow \Sigma \times \Sigma^\omega & \xrightarrow{\quad \mu \quad} & \Sigma \times S \rightsquigarrow \Sigma \times \Sigma^\omega \\
\uparrow c_1 & & \uparrow c_2 \\
S_1 \rightsquigarrow \Sigma^\omega & \xrightarrow{\quad \text{beh}_1 \quad} & S_2 \rightsquigarrow \Sigma^\omega \\
& & \text{beh}_2
\end{array}
\quad \text{in } \mathcal{K}\ell(\mathcal{P}).$$

$$\text{beh}_1 \stackrel{\mu}{=} (\eta_{\Sigma^\omega} \circ d)^{-1} \odot \overline{F}[\text{beh}_1, \text{beh}_2] \odot c_1$$

$$\text{beh}_2 \stackrel{\nu}{=} (\eta_{\Sigma^\omega} \circ d)^{-1} \odot \overline{F}[\text{beh}_1, \text{beh}_2] \odot c_2$$

$$\text{beh}_1 \stackrel{\mu}{=} (\eta_{\Sigma^\omega} \circ d)^{-1} \odot \overline{F}[\text{beh}_1, \text{beh}_2] \odot c_1$$

$$\text{beh}_2 \stackrel{\nu}{=} (\eta_{\Sigma^\omega} \circ d)^{-1} \odot \overline{F}[\text{beh}_1, \text{beh}_2] \odot c_2$$

$$\text{beh}_1 \stackrel{\mu}{=} (\eta_{\Sigma^\omega} \circ d)^{-1} \odot \overline{F}[\text{beh}_1, \text{beh}_2] \odot c_1 \qquad \text{beh}_2 \stackrel{\nu}{=} (\eta_{\Sigma^\omega} \circ d)^{-1} \odot \overline{F}[\text{beh}_1, \text{beh}_2] \odot c_2$$

Rewrite to:

$$\text{beh}_1 \stackrel{\mu}{=} \Diamond_\delta([\text{beh}_1, \text{beh}_2]) \upharpoonright S_1 \qquad \text{beh}_2 \stackrel{\nu}{=} \Diamond_\delta([\text{beh}_1, \text{beh}_2]) \upharpoonright S_2$$

Where $\Diamond_\delta : (\mathcal{P}(\Sigma^\omega))^S \rightarrow (\mathcal{P}(\Sigma^\omega))^S$ is given by

$$\Diamond_\delta(\text{beh})(s) = \{\sigma \cdot w \mid s' \in \delta(s)(\sigma), w \in \text{beh}(s')\}.$$

$$\text{beh}_1 \stackrel{\mu}{=} (\eta_{\Sigma^\omega} \circ d)^{-1} \odot \overline{F}[\text{beh}_1, \text{beh}_2] \odot c_1 \qquad \text{beh}_2 \stackrel{\nu}{=} (\eta_{\Sigma^\omega} \circ d)^{-1} \odot \overline{F}[\text{beh}_1, \text{beh}_2] \odot c_2$$

Rewrite to:

$$\text{beh}_1 \stackrel{\mu}{=} \Diamond_\delta([\text{beh}_1, \text{beh}_2]) \upharpoonright S_1 \qquad \text{beh}_2 \stackrel{\nu}{=} \Diamond_\delta([\text{beh}_1, \text{beh}_2]) \upharpoonright S_2$$

Where $\Diamond_\delta : (\mathcal{P}(\Sigma^\omega))^S \rightarrow (\mathcal{P}(\Sigma^\omega))^S$ is given by

$$\Diamond_\delta(\text{beh})(s) = \{\sigma \cdot w \mid s' \in \delta(s)(\sigma), w \in \text{beh}(s')\}.$$

Definition: The *solution* to this *equational system* is calculated as follows:

- Intermediate solution $l_1^{(1)} := \mu u_1 \cdot f_1(u_1, u_2)$
- $l^{\text{sol}} := \nu u_2 \cdot f_2(l_1^{(1)}(u_2), u_2)$
- $l_1^{\text{sol}} = l_1^{(1)}(l_2^{\text{sol}})$

$$\text{beh}_1 \stackrel{\mu}{=} (\eta_{\Sigma^\omega} \circ d)^{-1} \odot \overline{F}[\text{beh}_1, \text{beh}_2] \odot c_1 \quad \text{beh}_2 \stackrel{\nu}{=} (\eta_{\Sigma^\omega} \circ d)^{-1} \odot \overline{F}[\text{beh}_1, \text{beh}_2] \odot c_2$$

Rewrite to:

$$\text{beh}_1 \stackrel{\mu}{=} \Diamond_\delta([\text{beh}_1, \text{beh}_2]) \upharpoonright S_1 \quad \text{beh}_2 \stackrel{\nu}{=} \Diamond_\delta([\text{beh}_1, \text{beh}_2]) \upharpoonright S_2$$

Where $\Diamond_\delta : (\mathcal{P}(\Sigma^\omega))^S \rightarrow (\mathcal{P}(\Sigma^\omega))^S$ is given by

$$\Diamond_\delta(\text{beh})(s) = \{\sigma \cdot w \mid s' \in \delta(s)(\sigma), w \in \text{beh}(s')\}.$$

Definition: The *solution* to this *equational system* is calculated as follows:

- Intermediate solution $l_1^{(1)} := \mu u_1 \cdot f_1(u_1, u_2)$
- $l_2^{\text{sol}} := \nu u_2 \cdot f_2(l_1^{(1)}(u_2), u_2)$
- $l_1^{\text{sol}} = l_1^{(1)}(l_2^{\text{sol}})$

Concretely:

- $l_1^{(1)} := \mu u_1 \cdot \Diamond_\delta([u_1, u_2]) \upharpoonright S_1$
- $l_2^{\text{sol}} := \nu u_2 \cdot u_2 \Diamond_\delta([\mu u_1 \cdot \Diamond_\delta([u_1, u_2]) \upharpoonright S_1, u_2]) \upharpoonright S_2, u_2)$
- $l_1^{\text{sol}} = \mu u_1 \cdot \Diamond_\delta([u_1, \nu u_2 \cdot u_2 \stackrel{\nu}{=} \Diamond_\delta([\mu u_1' \cdot \Diamond_\delta([u_1', u_2]) \upharpoonright S_1, u_2]) \upharpoonright S_2, u_2]) \upharpoonright S_1$

Let $A = \langle S, \Sigma, \delta, s_0, F \rangle$ be a Büchi automaton. Take $S_1 = S \setminus F$, $S_2 = F$. Model δ by coalgebras $c_1 : S_1 \rightarrow \mathcal{P}(\Sigma \times S)$, $c_2 : S_2 \rightarrow \mathcal{P}(\Sigma \times S)$. Take the initial algebra $d : \Sigma^\omega \rightarrow \Sigma \times \Sigma^\omega$ defined by $d(\sigma w) = (\sigma, w)$ in **Sets**.

$$\begin{array}{ccc}
 \begin{array}{ccc}
 \Sigma \times [\text{beh}_1, \text{beh}_2] \\
 \Sigma \times S \rightsquigarrow \Sigma \times \Sigma^\omega \\
 \uparrow c_1 \quad \quad \quad \uparrow \eta_{\Sigma^\omega} \circ d \\
 S_1 \rightsquigarrow \Sigma^\omega \\
 \text{beh}_1
 \end{array}
 & \overset{\mu}{=} &
 \begin{array}{ccc}
 \Sigma \times [\text{beh}_1, \text{beh}_2] \\
 \Sigma \times S \rightsquigarrow \Sigma \times \Sigma^\omega \\
 \uparrow c_2 \quad \quad \quad \uparrow \eta_{\Sigma^\omega} \circ d \\
 S_2 \rightsquigarrow \Sigma^\omega \\
 \text{beh}_2
 \end{array}
 \end{array}
 \quad \text{in } \mathcal{K}\ell(\mathcal{P}).$$

$$\text{beh}_1 \overset{\mu}{=} \Diamond_\delta([\text{beh}_1, \text{beh}_2]) \upharpoonright S_1 \quad \quad \text{beh}_2 \overset{\nu}{=} \Diamond_\delta([\text{beh}_1, \text{beh}_2]) \upharpoonright S_2$$

Where $\Diamond_\delta : (\mathcal{P}(\Sigma^\omega))^S \rightarrow (\mathcal{P}(\Sigma^\omega))^S$ is given by

$$\Diamond_\delta(\text{beh})(s) = \{\sigma \cdot w \mid s' \in \delta(s)(\sigma), w \in \text{beh}(s')\}.$$

Let $A = \langle S, \Sigma, \delta, s_0, F \rangle$ be a Büchi automaton. Take $S_1 = S \setminus F$, $S_2 = F$. Model δ by coalgebras $c_1 : S_1 \rightarrow \mathcal{P}(\Sigma \times S)$, $c_2 : S_2 \rightarrow \mathcal{P}(\Sigma \times S)$. Take the initial algebra $d : \Sigma^\omega \rightarrow \Sigma \times \Sigma^\omega$ defined by $d(\sigma w) = (\sigma, w)$ in **Sets**.

$$\begin{array}{ccc}
 \Sigma \times [\text{beh}_1, \text{beh}_2] & & \Sigma \times [\text{beh}_1, \text{beh}_2] \\
 \Sigma \times S \rightsquigarrow \Sigma \times \Sigma^\omega & & \Sigma \times S \rightsquigarrow \Sigma \times \Sigma^\omega \\
 \uparrow c_1 & \overset{=}{\mu} & \uparrow c_2 \\
 S_1 \rightsquigarrow \Sigma^\omega & & S_2 \rightsquigarrow \Sigma^\omega \\
 \text{beh}_1 & & \text{beh}_2
 \end{array}
 \quad \eta_{\Sigma^\omega} \circ d \quad \text{in } \mathcal{Kl}(\mathcal{P}).$$

$$\text{beh}_1 \overset{\mu}{=} \Diamond_\delta([\text{beh}_1, \text{beh}_2]) \upharpoonright S_1 \quad \text{beh}_2 \overset{\nu}{=} \Diamond_\delta([\text{beh}_1, \text{beh}_2]) \upharpoonright S_2$$

Where $\Diamond_\delta : (\mathcal{P}(\Sigma^\omega))^S \rightarrow (\mathcal{P}(\Sigma^\omega))^S$ is given by

$$\Diamond_\delta(\text{beh})(s) = \{\sigma \cdot w \mid s' \in \delta(s)(\sigma), w \in \text{beh}(s')\}.$$

Theorem [Urabe, Shimizu, Hasuo 2016]: The solutions $\text{beh}_1, \text{beh}_2$ to the system of equations coincide with the accepted language of the Büchi Automaton A .

Alternate Proof of Coincidence Result

Problem: system of fixed point equations is convoluted.

Alternate Proof of Coincidence Result

Problem: system of fixed point equations is convoluted.

Alternate derivation using game semantics:

Game Semantics For Modal Mu-Calculus: $s \models^T \varphi \iff$ verifier has a winning strategy in $\mathcal{G}(\varphi, T)$

Outline:

- Convert system of equations to modal mu-calculus formula
- Apply game semantics of modal mu-calculus
- Prove: V has a winning strategy in $\mathcal{G}(\varphi, T)$ from state $(s_i, w) \iff w \in \text{beh}(s_i)$

Converting formula:

$$l_{\text{sol}}^2 = \nu u_2. \Diamond_{\delta}^2 [(\mu u'_1. \Diamond_{\delta}^1 [u'_1, u_2]), u_2]$$

$$\overline{\varphi_2} = \nu u_2. (p_2 \wedge \Diamond((\mu u'_1. p_1 \wedge \Diamond((p_1 \wedge u'_1) \vee (p_2 \wedge u_2))) \vee (p_2 \wedge u_2))))$$

Converting formula:

$$l_{\text{sol}}^2 = \nu u_2. \Diamond_{\delta}^2 [(\mu u'_1. \Diamond_{\delta}^1 [u'_1, u_2]), u_2]$$

$$\overline{\varphi_2} = \nu u_2. (p_2 \wedge \Diamond((\mu u'_1. p_1 \wedge \Diamond((p_1 \wedge u'_1) \vee (p_2 \wedge u_2))) \vee (p_2 \wedge u_2))))$$

Defining Transition System for Büchi Automaton A :

Let $A = (S_1 \cup S_2, \Sigma, \delta)$ be a Büchi automaton. Let Transition System (TS) over the set of propositional variables $\{p_1, p_2\}$, denoted as T_A , as follows:

- States: (s, w) for $s \in S$ and $w \in \Sigma^\omega$
- Transition $(s, \sigma w) \rightarrow (s', w)$ for $s, s' \in S, \sigma \in \Sigma, w \in \Sigma^\omega$, iff $s' \in \delta(s)(\sigma)$
- Labeling function: $\lambda((s, w)) = \{p_i\}$ iff $s \in S_i$

Converting formula:

$$l_{\text{sol}}^2 = \nu u_2. \Diamond_{\delta}^2 [(\mu u_1'. \Diamond_{\delta}^1 [u_1', u_2]), u_2]$$

$$\overline{\varphi_2} = \nu u_2. (p_2 \wedge \Diamond((\mu u_1'. p_1 \wedge \Diamond((p_1 \wedge u_1') \vee (p_2 \wedge u_2))) \vee (p_2 \wedge u_2))))$$

Defining Transition System for Büchi Automaton A :

Let $A = (S_1 \cup S_2, \Sigma, \delta)$ be a Büchi automaton. Let Transition System (TS) over the set of propositional variables $\{p_1, p_2\}$, denoted as T_A , as follows:

- States: (s, w) for $s \in S$ and $w \in \Sigma^\omega$
- Transition $(s, \sigma w) \rightarrow (s', w)$ for $s, s' \in S, \sigma \in \Sigma, w \in \Sigma^\omega$, iff $s' \in \delta(s)(\sigma)$
- Labeling function: $\lambda((s, w)) = \{p_i\}$ iff $s \in S_i$

Game Semantics For Modal Mu-Calculus: $(s, w) \stackrel{T}{\models} \varphi$ iff Verifier has a winning strategy in $\mathcal{G}(\varphi, T_A)$ from state $(\varphi, (s, w))$

Converting formula:

$$l_{\text{sol}}^2 = \nu u_2. \Diamond_{\delta}^2 [(\mu u'_1. \Diamond_{\delta}^1 [u'_1, u_2]), u_2]$$

$$\overline{\varphi_2} = \nu u_2. (p_2 \wedge \Diamond((\mu u'_1. p_1 \wedge \Diamond((p_1 \wedge u'_1) \vee (p_2 \wedge u_2))) \vee (p_2 \wedge u_2))))$$

Defining Transition System for Büchi Automaton A :

Let $A = (S_1 \cup S_2, \Sigma, \delta)$ be a Büchi automaton. Let Transition System (TS) over the set of propositional variables $\{p_1, p_2\}$, denoted as T_A , as follows:

- States: (s, w) for $s \in S$ and $w \in \Sigma^\omega$
- Transition $(s, \sigma w) \rightarrow (s', w)$ for $s, s' \in S, \sigma \in \Sigma, w \in \Sigma^\omega$, iff $s' \in \delta(s)(\sigma)$
- Labeling function: $\lambda((s, w)) = \{p_i\}$ iff $s \in S_i$

Game Semantics For Modal Mu-Calculus: $(s, w) \models^T \varphi$ iff Verifier has a winning strategy in $\mathcal{G}(\varphi, T_A)$ from state $(\varphi, (s, w))$

Lemma: Verifier has a winning strategy in $\mathcal{G}(\varphi, T_A)$ from state iff the Büchi automaton accepts w from s .

1. Büchi Automata
 - Modeling infinite behavior
2. Coalgebraic Representation Nondeterministic Finite Automata
 - Work in $\mathcal{KL}(\mathcal{P})$
 - Initial coalgebra in **Sets** yields final coalgebra in $\mathcal{KL}(\mathcal{P})$
3. Coalgebraic Representation Possibly Infinite Behavior Nondeterministic Finite Automata
 - Final coalgebra in **Sets** yields weakly final coalgebra in $\mathcal{KL}(\mathcal{P})$
 - This adds infinite behavior
4. Coalgebraic Representation Büchi Automata
 - Split $S = S_1 \cup S_2$
 - Take those traces which are solution to system of fixed point equations
5. Outline Derivation using Game Semantics
 - Use game semantics for modal mu-calculus to obtain more comprehensive proof of coincidence