

PUBLISHED

Question	Answer	Marks																				
9(b)(i)	any four from : 1 increase then decrease ; 2 optimum temperature at 29 °C ; 3 data quote – two temperatures and two rates with units ; <table><tr><th>temperature / °C ±0.5</th><th>rate of photosynthesis / mg g⁻¹ h⁻¹ ±0.05</th></tr><tr><td>2.5</td><td>0.75</td></tr><tr><td>5</td><td>1.1</td></tr><tr><td>10</td><td>1.8</td></tr><tr><td>20</td><td>2.95</td></tr><tr><td>29</td><td>3.5</td></tr><tr><td>30</td><td>3.45</td></tr><tr><td>40</td><td>2.65</td></tr><tr><td>45</td><td>1.5</td></tr><tr><td>48</td><td>0.5</td></tr></table> increase / before 29 °C 4 more kinetic energy for, enzyme (activity) / AW ; 5 temperature is the limiting factor ; decrease / after 29 °C 6 denaturation of, rubisco / enzymes / oxygen-evolving complex ; 7 AVP ; e.g. photorespiration limiting the rate at above 25 °C	temperature / °C ±0.5	rate of photosynthesis / mg g ⁻¹ h ⁻¹ ±0.05	2.5	0.75	5	1.1	10	1.8	20	2.95	29	3.5	30	3.45	40	2.65	45	1.5	48	0.5	4
temperature / °C ±0.5	rate of photosynthesis / mg g ⁻¹ h ⁻¹ ±0.05																					
2.5	0.75																					
5	1.1																					
10	1.8																					
20	2.95																					
29	3.5																					
30	3.45																					
40	2.65																					
45	1.5																					
48	0.5																					