

Question	Answer	Marks
1(a)	I All the following data recorded. - two burette readings AND titre for the rough titration - initial and final readings for two (or more) accurate titrations II Titre values recorded for accurate titrations, AND correct headings and units in the accurate titration table - initial / start AND (burette) reading / volume - final / end AND (burette) reading / volume - titre OR volume used / added OR FB 2 used / added - unit: / cm³ OR (cm³) OR in cm³ (for each heading) OR cm³ unit given for each volume recorded. III All accurate burette readings recorded to 0.05 cm³. IV The final accurate titre recorded must be within 0.10 cm³ of any other accurate titre.	7
	Accuracy (Q) marks Round burette readings to the nearest 0.05 cm³. Check and correct titre subtractions where necessary. Select the best mean titre, using the following hierarchy: - two (or more) accurate identical titres (ignoring any that are labelled “rough”), <i>then</i> - two (or more) accurate titres within 0.05 cm³, <i>then</i> - two(or more) accurate titres within 0.10 cm³, <i>etc.</i> Calculate the Supervisor’s mean titre to two decimal places. Calculate the Candidate’s mean titre to two decimal places. Calculate the difference (δ) between the Supervisor’s and Candidate’s mean titre.	
	Award accuracy Q marks as follows: V award if $\delta \leq 0.50 \text{ cm}^3$ VI award if $\delta \leq 0.30 \text{ cm}^3$ VII award if $\delta \leq 0.20 \text{ cm}^3$	