

Linear Algebra: Applications of Matrices

Glenn Bruns
CSUMB

Some of the material in these slides comes from Géron's notes:
https://github.com/ageron/handson-ml/blob/master/math_linear_algebra.ipynb

Learning outcomes

After this lecture you should be able to:

1. Use matrix multiplication to:
 - create derive new features from existing features
 - make predictions from a linear model
 - perform projections

Review: Matrix multiplication

$$\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} =$$

$$\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} =$$

Application: creating new features

Suppose we want to create a new feature that is derived from existing features.

feature data $\rightarrow \begin{pmatrix} 1 & -5 & 3 \\ 2 & -6 & 4 \end{pmatrix} \begin{pmatrix} 1 \\ .5 \\ .2 \end{pmatrix}$ \leftarrow take this linear combination of existing features

$$= 1 \begin{pmatrix} 1 \\ 2 \end{pmatrix} + .5 \begin{pmatrix} -5 \\ -6 \end{pmatrix} + .2 \begin{pmatrix} 3 \\ 4 \end{pmatrix}$$
$$= \begin{pmatrix} 1 \\ 2 \end{pmatrix} + \begin{pmatrix} -2.5 \\ -3 \end{pmatrix} + \begin{pmatrix} 0.6 \\ 0.8 \end{pmatrix}$$
$$= \begin{pmatrix} -0.9 \\ -0.2 \end{pmatrix}$$

Another view of matrix multiplication

Each column of AB is a linear combination of the columns in A .

$$\begin{pmatrix} 1 & 2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 2 & 3 \end{pmatrix} = \begin{pmatrix} 5 & 7 \\ 8 & 11 \end{pmatrix}$$

$$1 \begin{pmatrix} 1 \\ 2 \end{pmatrix} + 2 \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 5 \\ 8 \end{pmatrix}$$

First column of B defines a linear combination of the columns of A

$$1 \begin{pmatrix} 1 \\ 2 \end{pmatrix} + 3 \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 7 \\ 11 \end{pmatrix}$$

Second column of B defines a linear combination of the columns of A

Create multiple new features

Suppose we want to create **3** new features derived from existing features.

$$\begin{pmatrix} 1 & -5 & 3 \\ 2 & -6 & 4 \end{pmatrix} \begin{pmatrix} 1 & 1/3 & 1/2 \\ .5 & 1/3 & 1/2 \\ .2 & 1/3 & 0 \end{pmatrix}$$

this describes
how to get third
new feature


$$= \begin{pmatrix} -0.9 & -\frac{1}{3} & -2 \\ -0.2 & 0 & -2 \end{pmatrix}$$

$$1 \begin{pmatrix} 1 \\ 2 \end{pmatrix} + 0.5 \begin{pmatrix} -5 \\ -6 \end{pmatrix} + 0.2 \begin{pmatrix} 3 \\ 4 \end{pmatrix} = ?$$

Yet another view

You can also think of matrix multiplication AB as:
matrix A times every column of B

$$\begin{pmatrix} 1 & -5 & 3 \\ 2 & -6 & 4 \end{pmatrix} \begin{pmatrix} 1 & 1/3 & 1/2 \\ .5 & 1/3 & 1/2 \\ .2 & 1/3 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & -5 & 3 \\ 2 & -6 & 4 \end{pmatrix} \begin{pmatrix} 1 \\ .5 \\ .2 \end{pmatrix} \quad \begin{pmatrix} 1 & -5 & 3 \\ 2 & -6 & 4 \end{pmatrix} \begin{pmatrix} 1/3 \\ 1/3 \\ 1/3 \end{pmatrix} \quad \begin{pmatrix} 1 & -5 & 3 \\ 2 & -6 & 4 \end{pmatrix} \begin{pmatrix} 1/2 \\ 1/2 \\ 0 \end{pmatrix}$$

$$= \begin{pmatrix} -0.9 & -\frac{1}{3} & -2 \\ -0.2 & 0 & -2 \end{pmatrix}$$

Application: linear regression

We make a prediction using a linear model:

$$\hat{y} = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \cdots + \theta_n x_n$$

where x_1, x_2, \dots, x_p are the predictors and $\theta_0, \theta_1, \dots, \theta_n$ are the model coefficients.

Example: predict car mileage from engine size and care weight. After training our model we have:

$$\hat{y} = 50 - 1.5 \text{ weight} - 5.2 \text{ size}$$

If weight (in tons) is 2.5, and engine size (in 100's of cubic inches) is 3, what is predicted mileage?

Application: linear regression

Making a prediction:

$$\hat{y} = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \cdots + \theta_n x_n$$

vectorized form:

$$\hat{y} = \theta^T \cdot \mathbf{x}$$

θ, \mathbf{x} are
column
vectors

$$(\theta_0, \theta_1, \theta_2) \begin{pmatrix} 1 \\ x_1 \\ x_2 \end{pmatrix} = \theta_0 + \theta_1 x_1 + \theta_2 x_2$$

Mileage model example:

$$\hat{y} = 50 - 1.5 \text{ weight} - 5.2 \text{ size}$$

$$\text{If } \mathbf{x} = \begin{pmatrix} 1 \\ 2.5 \\ 3 \end{pmatrix}, \text{ then } \hat{y} = (50, -1.5, -5.2) \begin{pmatrix} 1 \\ 2.5 \\ 3 \end{pmatrix} = 30.65$$

Matrices as vector transformers

Here's a function that maps a vector to another vector:

$$f\left(\begin{pmatrix} x \\ y \\ z \end{pmatrix}\right) = \begin{pmatrix} 2x \\ x + y \\ 3z \end{pmatrix}$$

Q: what is $f\left(\begin{pmatrix} -1 \\ 2 \\ 3 \end{pmatrix}\right)$?

Another way to write it:

$$f(u) = \begin{pmatrix} 2 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 3 \end{pmatrix} u$$

$$3 \times 3 \quad 3 \times 1 \rightarrow 3 \times 1$$


Lots of functions on vectors could be written like this:

$$f(u) = Au \quad (A \text{ is } n \times n, \text{ where } u \text{ has length } n)$$

You can think of u being "transformed" by A .

Projection matrices

What is the projection of the vector onto the x axis?




$$\begin{pmatrix} & \end{pmatrix} \begin{pmatrix} 3.5 \\ 3 \end{pmatrix} = \begin{pmatrix} 3.5 \\ 0 \end{pmatrix}$$

What values should be in the 2 by 2 projection matrix to give the desired projection?

Projection matrices

What is the projection of the vector onto the x,y plane?



$$\begin{pmatrix} & & \end{pmatrix} \begin{pmatrix} 4 \\ 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 4 \\ 2 \\ 0 \end{pmatrix}$$

What values should be in the 3 by 3 projection matrix to give the desired projection?

And to project the same vector onto the z axis?

Summary

Some applications of matrix multiplication to machine learning

- ❑ derive new features as the linear combination of existing features
- ❑ compute predictions from a linear model
- ❑ project vectors