

1 Gauge Theory

Gauge Transformation

Local transformation that leaves the Lagrangian/Hamiltonian unchanged.

Schrödinger equation is invariant under global transformations of the wave function

$$\psi \rightarrow \psi' = e^{i\theta} \psi \quad (1)$$

Electrodynamics can be derived from a gauge transformation of the wave function

$$\psi \rightarrow \psi' = e^{i\theta(t,x)} \psi \quad (2)$$

we write

$$\theta(t, x) \equiv \frac{q}{\hbar c} \Lambda(t, x) \quad (3)$$

Schrödinger eq. is invariant under a gauge transformation of the wave function by introducing the gauge fields \vec{A} and φ satisfying the transformations

$$\vec{A}' = \vec{A} + \nabla \Lambda \quad (4)$$

$$\varphi' = \varphi - \frac{1}{c} \frac{\partial \Lambda}{\partial t} \quad (5)$$

We define the gauge invariant fields

$$\vec{B} = \nabla \times \vec{A} \quad (6)$$

$$\vec{E} = -\nabla \varphi - \frac{1}{c} \frac{\partial \vec{A}}{\partial t} \quad (7)$$

The *magnetic field* \vec{B} does no work. The power we need to move a particle subject to a Lorentz force, i.e. against the fields is

$$P = -q \vec{E} \cdot \vec{v} \quad (8)$$

We define the current of charge particles as

$$I = \vec{j} \cdot \Delta \vec{S} \quad (9)$$

$$nq = \rho \quad \text{charge density} \quad (10)$$

$$\vec{j} = \rho \vec{v} \quad \text{current density} \quad (11)$$

Lorentz Force

In the presence of electric and magnetic fields, a particle with mass m and charge q experience a force given by

$$\vec{F} = q(\vec{E} + \frac{\vec{v}}{c} \times \vec{B}) \quad (12)$$

ALL CONSERVATION LAWS HAVE TO BE *LOCAL* in order to respect the relativity of simultaneity.

Conservation of charge

$$\frac{dQ_V}{dt} + \int_{\partial V} \vec{j} \cdot d\vec{a} = 0 \quad (13)$$

Generic conservation law

$$\frac{dC_V}{dt} + \int_{\partial V} \underbrace{\vec{j}}_{\text{density flux}} \cdot d\vec{a} = \underbrace{F_V}_{\text{sources within } V} \quad (14)$$

Conservation of energy

$$\frac{d}{dt} \int_V d^3\vec{r} u + \int_{\partial V} d\vec{a} \cdot \vec{S} = - \int_V d^3\vec{r} \vec{j} \cdot \vec{E} \quad (15)$$

where $u = \frac{E^2+B^2}{8\pi}$ is the e-m energy density and $\vec{S} = \frac{c}{4\pi} \vec{E} \times \vec{B}$ is the Poynting vector.

Maxwell equations

$$\nabla \cdot \vec{B} = 0 \quad (16)$$

$$\nabla \times \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t} \quad (17)$$

$$\nabla \cdot \vec{E} = 4\pi\rho \quad (18)$$

$$\nabla \times \vec{B} = \frac{4\pi}{c} \vec{j} + \frac{1}{c} \frac{\partial \vec{E}}{\partial t} \quad (19)$$

Gauss's law	$\int_{\partial V} d\vec{a} \cdot \vec{E} = 4\pi \int_V d^3r \rho \quad (20)$	The flux of \vec{E} through a closed surface equals the enclosed charge.
—	$\int_{\partial V} d\vec{a} \cdot \vec{B} = 0 \quad (21)$	Magnetic charges are absent.
Faraday's law	$\oint_C d\vec{r} \cdot \vec{E} = - \int_S d\vec{a} \cdot \frac{1}{c} \frac{\partial \vec{B}}{\partial t} \quad (22)$	A varying magnetic flux generates an electromotive force.
Ampere's law	$\oint_C d\vec{r} \cdot \vec{B} = \int_S d\vec{a} \cdot \left(\frac{4\pi}{c} \vec{j} + \frac{1}{c} \frac{\partial \vec{E}}{\partial t} \right) \quad (23)$	The circulation of \vec{B} along a closed curve C bordering a surface S equals the flux thorough S the sum of hte usual current plus Maxwell's displacement current $\frac{1}{c} \frac{\partial \vec{E}}{\partial t}$.

Helmholtz Theorem

Equations for the potentials

$$\nabla^2 \varphi + \frac{1}{c} \frac{\partial}{\partial t} \nabla \cdot \vec{A} = -4\pi\rho \quad (24)$$

$$\nabla^2 \vec{A} - \nabla \left(\nabla \cdot \vec{A} + \frac{1}{c} \frac{\partial \varphi}{\partial t} \right) - \frac{1}{c^2} \frac{\partial^2 \vec{A}}{\partial t^2} = -\frac{4\pi}{c} \vec{j} \quad (25)$$

Gauge Fixing

$$\textbf{Lorenz} \quad \nabla \cdot \vec{A} \equiv -\frac{1}{c} \frac{\partial \varphi}{\partial t} \quad (26)$$

$$\textbf{Coulomb} \quad \nabla \cdot \vec{A} \equiv 0 \quad (27)$$

$$\textbf{Hamilton} \quad \varphi \equiv 0 \quad (28)$$

$$\textbf{Landau} \quad A_x \equiv 0 \text{ (Quantum Hall Effect)} \quad (29)$$

Using the Lorenz gauge

$$\nabla^2 \varphi - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \varphi = -4\pi \rho \quad (30)$$

$$\nabla^2 \vec{A} - \frac{1}{c^2} \frac{\partial^2 \vec{A}}{\partial t^2} = -\frac{4\pi}{c} \vec{j} \quad (31)$$

Using the Coulomb gauge

$$\nabla^2 \varphi = -4\pi \rho \quad (32)$$

$$\nabla^2 \vec{A} - \frac{1}{c^2} \frac{\partial^2 \vec{A}}{\partial t^2} = -\frac{4\pi}{c} \vec{j}_T \quad (33)$$

In Coulomb's gauge and applying the fourier transform in the spatial derivatives

$$\underbrace{\frac{d^2 A_k^\alpha}{dt^2}}_{\text{Harm. osc. } \omega_k=kc} = -k^2 c^2 A_k^\alpha + \underbrace{4\pi c j_k^\alpha}_{\text{Ext. rest. force}} \quad (34)$$

Casimir Effect

Consider in the vacuum two perfect mirrors in front of each other at a distance L . Inside the mirrors suppose there is a potential \vec{A} . Each mode have a wave length given by the boundary conditions and can only take discrete values

$$\lambda = \frac{2L}{m}, \Rightarrow k = \frac{m\pi}{L}, m = 1, 2, \dots \quad (35)$$

Each of these modes have an energy

$$\epsilon_m = \hbar \omega_m (n_{m,\alpha} + 1/2) = \frac{\hbar m \pi c}{L} (n_{m,\alpha} + 1/2), \quad (36)$$

n counts photons —not a well defined quantity—, m enumerates modes and $\alpha = 1, 2$ enumerates the degrees of freedom. The total energy is then

$$U = \sum_{m\alpha} \frac{\hbar m \pi c}{L} (\langle n_{m,\alpha} \rangle + 1/2). \quad (37)$$

In the lowest energy state with temperature $T = 0$ we have no photons $n = 0$. Thus, the energy is

$$U = 2 \sum_m \frac{\hbar m \pi c}{2L} = -\frac{\hbar \pi c}{12L}. \quad (38)$$

And the force

$$F = -\frac{d}{dL} U(L) = -\frac{\hbar \pi c}{12L^2} \quad (39)$$

The apparence of an attractive force between the mirrors is called the *Casimir effect*.
Pahse differences

Aharonov-Bohm Effect

Suppose a double slit experiment with electrons, between the double slti and the screen we put a solenoid with a current circulating in the azimuthal direction. Inside the solenoid there is a magnetic field $\vec{B} \neq 0$, but outside $\vec{B} = 0$.

Over trajewctory $\mathcal{L}_{1,2}$ we have $\vec{B} = \nabla \times \vec{A} = 0 \Rightarrow \vec{A} = \nabla \Lambda_{1,2} \Rightarrow \Lambda_{1,2} = \int_{I \rightarrow F \text{ over } \mathcal{L}_{1,2}} d\vec{l} \cdot \vec{A}$.

Let $\Psi_{01,02}$ be the wave function of an electron when \vec{B} is off over trajectory $\mathcal{L}_{1,2}$. When \vec{B} is on the wave function $\Psi_{01,02}$ become $\Psi_{1,2}$

$$\Psi_{01,02} \rightarrow \Psi_{1,2} \propto e^{ikL_{1,2} + \frac{q}{\hbar c} \Lambda_{1,2}} \quad (40)$$

a phase difference in the fringe pattern appears

$$\Delta\varphi = k\Delta L + \frac{q}{\hbar c}(\Lambda_1 - \Lambda_2) \quad (41)$$

where

$$\Lambda_1 - \Lambda_2 = \oint d\vec{l} \cdot \vec{A} = \int d\vec{a} \cdot \nabla \times \vec{A} = \Phi_B \quad (42)$$

The shift in the fringe pattern i a double slit experiment in a region where $\vec{B} = 0$ is called the *Aharonov-Bohm effect*.




Figure 1: Thick, \vec{B} off. Dashed, \vec{B} on.

2 Electrostatics

Electric Field of a point charge

$$\vec{E}(\vec{r}) = \frac{q}{r^3} \vec{r} \quad (43)$$

Dirac Delta function

$$\int_{-\infty}^{\infty} \frac{dk}{2\pi} e^{ik(x-x')} = \delta(x' - x) \quad (44)$$

Work to move a charge against \vec{E}

$$W = -q \int_{C'} d\vec{l} \cdot \vec{E}(\vec{r}) \quad (45)$$

Potential of a point charge due to \vec{E}

$$\varphi(\vec{r}) = - \int_{\vec{O}}^{\vec{r}} d\vec{l} \cdot \vec{E}(\vec{r}) \quad (46)$$

Charged cylinder

Charged plane

Consider a square plane of length a with a charge q , then its superficial charge density is $\sigma = \frac{q}{a^2}$

Charged plaque

Capacitance and capacitors

The capacitance C is the ability of a body to hold charge. And it is defined through

$$Q = CV \quad (47)$$

A capacitor is a device that stores energy in a electrical field.

Poisson's Equation

Theorem Let $\rho(\vec{r}) \in V$ and the potential in the boundary is given $F(\vec{r}) = \varphi(\vec{r}) \in \partial V$, then the potential in the volume $\varphi(\vec{r})$, $\vec{r} \in V$ is unique.

Proof:

Suppose φ_1 and φ_2 are two different solutions of Laplace equation and $\varphi_1 = \varphi_2 = F$ in the boundary. Let $\Psi = \varphi_2 - \varphi_1$, then $\nabla^2 \Psi = 0$ and $\Psi = 0$ in the boundary. Then

$$\int_V d^3r |\nabla \Psi|^2 = \int_V d^3r \nabla \cdot (\Psi \nabla \Psi) - \int_V d^3r \nabla^2 \Psi = \int_{\partial V} d\vec{a} \cdot \Psi \nabla \Psi = 0 \quad (48)$$

therefore $|\nabla \Psi|^2 = 0$, $\forall \vec{r} \in V \Rightarrow \nabla \Psi = 0 \Rightarrow \Psi = \text{cte}$. Thus $\Psi = 0$. QED.

Theorem Let $\rho(\vec{r}) \in V$ and the directional derivative —along the perpendicular of the volume— of the potential in the boundary is given $D(\vec{r}) = \partial_n \varphi(\vec{r}) \in \partial V$, then the potential in the volume $\varphi(\vec{r})$, $\vec{r} \in V$ is unique up to a constant term.

Proof:

Suppose φ_1 and φ_2 are two different solutions of Laplace equation and $\partial_n \varphi_1 = \partial_n \varphi_2 = D$ in the boundary. Let $\Psi = \varphi_2 - \varphi_1$, then $\nabla^2 \Psi = 0$ and $\partial_n \Psi = 0$ in the boundary. Then

$$\int_V d^3r |\nabla \Psi|^2 = \int_{\partial V} d\vec{a} \cdot \Psi \nabla \Psi = \int_{\partial V} da \cdot (\hat{n} \nabla \Psi) \Psi = \int_{\partial V} da (\partial_n \Psi) \Psi = 0 \quad (49)$$

therefore $\nabla \Psi = 0 \Rightarrow \Psi = \text{cte}$. Thus $\varphi_2 = \varphi_1 + \text{cte}$. QED.

Neumann and Dirichlet Boundary Conditions

Spontaneous Symmetry Breaking

Green's function

Let $G(\vec{r}, \vec{r}')$ be such that

$$(\nabla')^2 G(\vec{r}, \vec{r}') = -4\pi \delta(\vec{r} - \vec{r}') \quad (50)$$

$$\begin{aligned}
\varphi(\vec{r}) &= \int_V d^3r' \varphi(\vec{r}') \delta(\vec{r} - \vec{r}') = -\frac{1}{4\pi} \int_V d^3r' \varphi(\vec{r}') (\nabla')^2 G(\vec{r}, \vec{r}') \\
&= -\frac{1}{4\pi} \int_V d^3r' \nabla' \cdot (\varphi(\vec{r}') \nabla' G(\vec{r}, \vec{r}')) + \frac{1}{4\pi} \int_V d^3r' (\nabla' \varphi(\vec{r}')) \cdot \nabla' G(\vec{r}, \vec{r}') \\
&= -\frac{1}{4\pi} \int_{\partial V} d\vec{a} \cdot \varphi(\vec{r}') \nabla' G(\vec{r}, \vec{r}') + \frac{1}{4\pi} \int_V d^3r' \nabla' \cdot (\nabla' \varphi(\vec{r}')) G(\vec{r}, \vec{r}') \\
&\quad - \frac{1}{4\pi} \int_V d^3r' ((\nabla')^2 \varphi(\vec{r}')) G(\vec{r}, \vec{r}') \tag{51}
\end{aligned}$$

$$= -\frac{1}{4\pi} \int_{\partial V} da \varphi(\vec{r}') \partial_{n'} G(\vec{r}, \vec{r}') + \frac{1}{4\pi} \int_{\partial V} da G(\vec{r}, \vec{r}') \partial_{n'} \varphi(\vec{r}') \tag{52}$$

$$+ \int_V d^3r' G(\vec{r}, \vec{r}') \rho(\vec{r}') \tag{53}$$

Dirichlet Green's Function

If we have Dirichlet's boundary conditions then

$$\varphi(\vec{r}) = \int_V d^3r' G_D(\vec{r}, \vec{r}') \rho(\vec{r}') - \frac{1}{4\pi} \int_{\partial V} da \varphi(\vec{r}') \partial_{n'} G_D(\vec{r}, \vec{r}') \tag{54}$$

Neumann Green's Function

With Neumann boundary conditions, having $\partial_{n'} G_N(\vec{r}, \vec{r}') = 0$, $\vec{r}' \in \partial V$ doesn't work since

$$\int_{\partial V} da' \partial_{n'} G(\vec{r}, \vec{r}') = -4\pi \tag{55}$$

The next easiest ansatz for G_N is a constant. Let $\partial_{n'} G_N = K = \text{cte}$. Then

$$\int_{\partial V} da' \partial_{n'} G(\vec{r}, \vec{r}') = K A = -4\pi \Rightarrow K = -\frac{4\pi}{A} \tag{56}$$

where A is the area of the region. And finally

$$\varphi(\vec{r}') = \int_V d^3r' G_N(\vec{r}, \vec{r}') + \langle \varphi \rangle_A + \frac{1}{4\pi} \int_{\partial V} da' G_N(\vec{r}, \vec{r}') \partial_{n'} \varphi(\vec{r}') \tag{57}$$

where $\langle \varphi \rangle_A = \frac{1}{A} \int_{\partial V} da' \varphi(\vec{r}')$ is the average of φ over the surface.

Green's function in Quantum Mechanics

Method of images

Consider a system composed by an infinite plane whose electric potential is known and given by a function $f(\vec{r})$ for \vec{r} in the plane, but is unknown everywhere else.

If the plane is on $z = 0$ as shown in image 2, the question is, what is the electric potential for $z > 0$? guessing there are some distributed charges upon it.

This problem is solved finding the Dirichlet function and using it in (54), but how to compute Dirichlet Green's function?, finding a potential produced by a charge placed upon the plane, with Dirichlet boundary conditions in the plane, this is $G_D(x, y, z = 0) = 0$.

This problem is exactly the same one that the system shown in 3, which is solved with images method.

The initial situation is a grounded metallic surface whit an electric charge above it, then by the images method we have to put an image charge q_i under the plane at the same




Figure 2: Plane at $\psi(\vec{r})$




Figure 3: An electrical charge q upon a metallic plane at $\psi(x, y, z = 0) = 0$ electric potential

distance from it than the real one, and with the same charge but with opposed sign, then the potential for $z > 0$ is,

$$\psi(\vec{r}) = \frac{q}{||\vec{r} - \vec{r}_0||} - \frac{q}{||\vec{r} - \vec{r}_i||}, \quad (58)$$

donde $\vec{r}_i = (x_0, y_0, -z_0)$,

$$\psi(x, y, z > 0) = \frac{q}{((x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2)^{\frac{1}{2}}} - \frac{q}{((x - x_0)^2 + (y - y_0)^2 + (z + z_0)^2)^{\frac{1}{2}}},$$

then it's Laplacian is

$$\nabla^2 \psi(\vec{r}) = -4\pi q \delta(\vec{r} - \vec{r}_0) + 4\pi q \delta(\vec{r} - \vec{r}_i), \quad (59)$$

the superposition of two Laplacians for Coloumbian potentials produced by q and q_i images. The interest region is $z > 0$ then the second term in the right side is zero.

$$\nabla^2 \psi(x, y, z > 0) = -4\pi q \delta(\vec{r} - \vec{r}_0), \quad (60)$$

then (58) obeys the Poisson equation, the differential equation for the system and the boundary condition $\psi(x, y, 0) = 0$

Green's function in spherical coordinates

$$G(\vec{r}, \vec{r}') = \sum_{lm} \frac{4\pi}{2l+1} \frac{r_{<}^l}{r_{>}^{l+1}} Y_{lm}^*(\Omega) Y_{lm}(\Omega') = \frac{1}{|\vec{r} - \vec{r}'|} \quad (61)$$

Dipolar Electric moment

$$\vec{p} = \int d^3r \rho(\vec{r}) \vec{r} \quad (62)$$

Potential

$$\varphi(\vec{r}) = -\vec{p} \cdot \nabla \frac{1}{r} = \frac{\vec{p} \cdot \vec{r}}{r} \quad (63)$$

Electric field

$$\vec{E}(\vec{r}) = \frac{3\vec{p} \cdot \vec{r} \vec{r} - \vec{p} r^2}{r^5} \quad (64)$$

Charge distribution of an electric dipole

$$\rho(\vec{r}) = -\vec{p} \cdot \nabla \delta(\vec{r}) \quad (65)$$

Polarization

We define the polarization \vec{P} as the density of electric dipole moments in a continuous system with volume V

$$\vec{P} \equiv \frac{\Delta \vec{p}}{\Delta V} \quad (66)$$

Let a tiny volume ΔV at \vec{r}_0 have a dipole moment $\Delta \vec{p}$ and a charge density $\Delta \rho^{\text{int}}$, then

$$\Delta \rho^{\text{int}} = -\Delta \vec{p} \cdot \nabla \delta(\vec{r} - \vec{r}_0) = -P(\vec{r}) \cdot \nabla \delta(\vec{r} - \vec{r}_0) \quad (67)$$

Integrating over the volume we get

$$\rho^{\text{int}} = -\nabla \cdot \vec{P} \quad (68)$$

Macroscopic Maxwell equations

Baoundary conditions

Energy of a system of charges

$$U = \frac{1}{2} \sum_{i \neq j} \frac{q_i q_j}{r_{ij}} \quad (69)$$

$$= \sum_{i < j} \frac{q_i q_j}{r_{ij}} \quad (70)$$

Energy of a continuous charge distribution

$$U = \frac{1}{2} \int d^3r \rho(\vec{r}) \varphi(\vec{r}) \quad (71)$$

Energy of a dipole

$$U = \vec{p} \cdot \nabla \varphi = -\vec{p} \cdot \vec{E} \quad (72)$$

Density matrix

Pure state

$$\vec{\rho} = |\Psi\rangle\langle\Psi| \quad (73)$$

$$\langle \hat{O} \rangle = \langle \Psi | \hat{O} | \Psi \rangle = \sum_{nm} \langle \Psi | n \rangle \langle n | \hat{O} | m \rangle \langle m | \Psi \rangle \quad (74)$$

$$= \sum_{nm} \langle n | \hat{O} | m \rangle \langle m | \Psi \rangle \langle \Psi | n \rangle \quad (75)$$

$$= \sum_{nm} O_{nm} \rho_{mn} = \sum_{nmk} (O_{nm} \rho_{mk}) \delta_{km} \quad (76)$$

$$= \sum_{nk} (O \rho)_{nk} \delta_{kn} = \sum_n (\hat{O} \hat{\rho})_{nn} \quad (77)$$

$$= \text{tr}(\hat{O} \hat{\rho}) \quad (78)$$

$$\hat{\rho} = \sum_{\Psi_\alpha} p_\alpha |\Psi_\alpha\rangle \langle \Psi_\alpha| \quad (79)$$

Schroedinger equation Interaction Scheme

$$i\hbar \partial_t \hat{O}_I = i\hbar \frac{d\hat{O}}{dt} + [\hat{O}, \hat{H}] \quad (80)$$

Evolution of $\hat{\rho}$ in interaction scheme

Consider an interaction Hamiltonian term of the form $H_I = \zeta \hat{A}(t) \varphi(t)$. Then the Schrodinger equation reads

$$i\hbar \partial_t \hat{\rho}(t) = \zeta [\hat{A}(t), \hat{\rho}] \varphi \quad (81)$$

Integrating the equation

$$\hat{\rho}(t) = \frac{1}{i\hbar} \int_{-\infty}^t dt' \zeta [\hat{A}(t'), \hat{\rho}(t')] \varphi(t') + \hat{\rho}_0 \quad (82)$$

where $\hat{\rho}_0$ is the matrix density in equilibrium, and we have assumed that $\varphi(t \rightarrow -\infty) = 0$. Using perturbation theory we assume a solution of the form

$$\hat{\rho}(t) = \hat{\rho}_0 + \zeta \hat{\rho}^{(1)}(t) + \zeta^2 \hat{\rho}^{(2)}(t) + \dots \quad (83)$$

So we get a solution

$$\hat{\rho}^{(n+1)}(t) = \frac{1}{i\hbar} \int_{-\infty}^t dt' [\hat{A}(t'), \hat{\rho}^{(n)}(t')] \varphi(t') \quad (84)$$

Response

Consider a operator \hat{B} subject to an interaction H_I defined as before. Then at first approximation the expectation value of \hat{B} is

$$\langle \hat{B}(t) \rangle^{(1)} = \text{tr}(\hat{\rho}^{(1)}(t) \hat{B}(t)) \quad (85)$$

$$= \frac{1}{i\hbar} \int_{-\infty}^t dt' \text{tr}(\hat{B}(t) [\hat{A}(t'), \hat{\rho}_0]) \varphi(t') \quad (86)$$

Then we define the response as

$$\chi_{BA}(t - t') = \frac{1}{i\hbar} \text{tr}(\hat{B}(t) [\hat{A}(t'), \hat{\rho}_0]) \Theta(t - t') \quad (87)$$

Use $\text{tr}(O_1 [O_2, O_3]) = \text{tr}([O_1, O_2] O_3)$ to rewrite it as

$$\chi_{AB}(\tau) = \frac{1}{i\hbar} \langle [\hat{B}(\tau), \hat{A}(0)] \rangle_0 \Theta(\tau) \quad (88)$$

where the subscript 0 means that it is computed at equilibrium.

Polarizability

Suppose an electron that is interacting with an electric dipole with energy $H_I = -\hat{d}_I(t) \cdot \hat{E}(t) = e\hat{r}(t) \cdot \hat{E}$. We identify $\hat{A} \rightarrow -\hat{d} = e\hat{r}(t)$. And we define $d(t) = \langle \hat{d}_I(t) \rangle$. We define the polarizability α through

$$d^i = \int dt' \alpha^{ij}(t-t') E^j(t') \quad (89)$$

If we identify $\hat{B} \rightarrow \hat{d} = -e\hat{r}$, then

$$\alpha^{ij}(\tau) = -\frac{e^2}{i\hbar} \langle [\hat{r}_I^i(\tau), \hat{r}_I^j(0)] \rangle_0 \Theta(\tau) \quad (90)$$

Applying fourier transform

$$d_\omega^i = \alpha_\omega^{ij} E_\omega^j \quad (91)$$

$$\alpha_\omega^{ij} = \int_{-\infty}^{\infty} d\tau \alpha^{ij}(\tau) e^{i(\omega+i\eta)\tau} \quad (92)$$

$$= -\frac{e^2}{i\hbar} \int_0^{\infty} d\tau \langle [\hat{r}(\tau), \hat{r}(0)] \rangle_0 e^{i\tilde{\omega}\tau} \quad (93)$$

Classical model of polarizability

Consider an electron attached to a nucleus through a spring with spring constant $\kappa = m\omega_0^2$ and subject to an electric force $-e\vec{E}(t)$. Then the equation of motion is

$$m \frac{d^2 \vec{r}}{dt^2} = -e\vec{E}(t) - m\omega_0^2 \vec{r} \quad (94)$$

We write the electric field as $\vec{E}(t) = \text{Re } \vec{E}_0 e^{-i\omega t + \eta t}$. We look for solutions of the form $\vec{r}(t) = \text{Re } \vec{r}_0 e^{-i\tilde{\omega} t}$ where $\tilde{\omega} = \omega + i\eta$. Solving the equation of motion with this ansatz we get

$$\vec{r} = -\frac{e/m}{\omega_0^2 - \tilde{\omega}^2} \vec{E} \quad (95)$$

If $\vec{d} = -e\vec{r}$, then

$$\vec{d} = \frac{e^2/m}{\omega_0^2 - \tilde{\omega}^2} \vec{E} \quad (96)$$

since $\vec{d} = \alpha \vec{E}$, then

$$\alpha(\omega) = \frac{e^2/m}{\omega_0^2 - \tilde{\omega}^2} \quad (97)$$

Oscillator force

$$\alpha = \frac{e^2}{m} \sum_{ij} \frac{f_{ij}}{\omega_{ij}^2 - \tilde{\omega}^2} \quad (98)$$

where f_{ij} is known as the oscillator force.

$$f_{nm}^{ij} = \frac{2m}{\hbar} \rho_{nn} r_{nm}^i r_{mn}^j \omega_{mn} (1 - \rho_m) \quad (99)$$

Classius-Mousotte relation

$$\frac{\epsilon - 1}{\epsilon + 2} = \frac{4\pi}{3} n\alpha \quad (100)$$

Plasma Frequency

$$\omega_p^2 = \frac{4\pi n e^2}{m} \quad (101)$$

Pruede's Model

$$\epsilon = 1 - \frac{\omega_p^2}{\omega} \quad (102)$$

Plasmons

3 Magnetostatics

Biot-Savart Law

Dipolar magnetic moment

$$\vec{m} \equiv \frac{1}{2c} \int d^3\vec{r}' \vec{r}' \times \vec{j}(\vec{r}') \quad (103)$$

Current density of a magnetic dipole moment

$$\vec{j} = c\nabla\delta(\vec{r}) \times \vec{m} \quad (104)$$

Potential

$$\vec{A}(\vec{r}) = \nabla \frac{1}{r} \times \vec{m} = \frac{\vec{m} \times \vec{r}}{r^3} \quad (105)$$

Magnetic field

$$\vec{B}(\vec{r}) = \frac{3\vec{p} \cdot \vec{r} \vec{r} - \vec{m}r^2}{r^5} \quad (106)$$

Magnetization

We define the magnetization \vec{M} as the density of dipole magnetic moments in a continuous system with volume V

$$\vec{M} \equiv \frac{\Delta\vec{m}}{\Delta V} \quad (107)$$

Let a tiny volume ΔV at \vec{r}_0 have a dipole moment $\Delta\vec{m}$ and a current density $\Delta\vec{j}$, then

$$\Delta\vec{j} = -c\Delta\vec{m} \times \nabla\delta(\vec{r} - \vec{r}_0) = -c\vec{M}(\vec{r}) \times \nabla\delta(\vec{r} - \vec{r}_0) \quad (108)$$

Integrating over the volume we get

$$\vec{j} = c\nabla \times \vec{M} \quad (109)$$

Torque

$$\vec{\tau} = \vec{m} \times \vec{B} \quad (110)$$

Force on a dipole due to an external magnetic field

We consider a current density around the origin enclosed in a volume V . From the Lorentz force

$$\vec{F} = \frac{1}{c} \int_V d^3r \vec{j}(\vec{r}) \times \vec{B}(\vec{r}) \quad (111)$$

we consider the source of \vec{B} to be very far and V to be small in order for a Taylor series around 0 to be reliable.

$$\vec{B}(\vec{r}) = \vec{B}(0) + \vec{r} \cdot \nabla \vec{B}(0) + \dots \quad (112)$$

The first non zero term of the force is

$$\vec{F}^{(1)} = (\vec{m} \times \nabla) \times \vec{B}(0) = \nabla(\vec{m} \cdot \vec{B}(0)) \quad (113)$$

Then the energy is

$$U = -\vec{m} \cdot \vec{B}(0) \quad (114)$$

Dipole moment of a continuous system in terms of \vec{L}

Suppose a free moving system made by a continuous charge (mass) distribution ρ_q (ρ_m) made by chunks of mass m_q and charge q . Then its angular momentum is

$$\vec{L} = \int d^3r \vec{r} \times \rho_m(\vec{r}) \vec{v} \quad (115)$$

where $\rho_m(\vec{r}) = n(\vec{r})m_q$, $\rho_q(\vec{r}) = n(\vec{r})q = \rho_m(\vec{r})\frac{q}{m_q}$ and $n(\vec{r}) \equiv \frac{\Delta N}{\Delta V}$ is the number density. Then the dipole moment \vec{m} is

$$\vec{m} = \frac{1}{2c} \int d^3r' \vec{r}' \times \vec{j}(\vec{r}') = \frac{1}{2c} \int d^3r' \vec{r}' \times \rho_q \vec{v}(\vec{r}') = \frac{q}{2m_q c} \int d^3r \vec{r} \times \rho_m \vec{v} = \frac{q}{2m_q c} \vec{L} \quad (116)$$

Bohr's magneton

If we model an electron by a continuous mass/charge distribution, then since its angular momentum is quantized $L_z = m\hbar$. The component of the electron's dipole moment $\vec{\mu}$ in the z direction is

$$\mu_z = -m \frac{e\hbar}{2m_e c} \quad (117)$$

is known as the *Bohr's magneton*. And

$$\Delta\mu_z = -\frac{\hbar e}{2m_e c} \quad (118)$$

Change in Bohr's magneton due to an external magnetic field

We consider an electron orbiting a nucleus, and we apply a uniform magnetic field in the z direction. We choose \vec{A} to be

$$\vec{A} = \frac{1}{2} \vec{B} \times \vec{r} \quad (119)$$

The linear momentum of a charge subject to a vector potential \vec{A} is

$$\vec{p} = m\vec{v} + \frac{q}{c} \vec{A}(\vec{r}) \quad (120)$$

and the angular momentum is

$$\vec{L} = \vec{r} \times m\vec{v} + \frac{q}{c} \vec{r} \times \vec{A}(\vec{r}) \quad (121)$$

The change in \vec{L} due to changes in \vec{A} is 0, then

$$0 = \Delta L = \Delta \int d^3r \vec{r} \times \rho_m(\vec{r}) \vec{v} - \Delta \int d^3r \rho_P(\vec{r}) \vec{r} \times \vec{A}(\vec{r}) \quad (122)$$

from here we get

$$\Delta \vec{\mu} = -\frac{e^2}{4m_e c^2} \vec{B} \langle r_{\perp}^2 \rangle \quad (123)$$

The presence of a magnetic field creates a magnetic moment in the opposite direction of the field.

Diamagnetism

If we sum the contribution of all the changes of the dipole moment of every electron in the system and divide it by the volume we obtain a magnetization

$$\vec{M} = \frac{\Delta \vec{\mu}}{\Delta V} = n \vec{\mu} = -n \frac{e^2}{4m_e c^2} \langle r_{\perp}^2 \rangle \vec{B} = \chi \vec{B} \quad (124)$$

then

$$\vec{B} = \frac{\vec{H}}{1 - 4\pi\chi} \approx (1 + 4\pi\chi) \vec{H} = \mu \vec{H} \quad (125)$$

where μ is the magnetic permeability, thus

$$\mu = 1 + 4\pi\chi = 1 - n \frac{e^2}{4m_e c^2} \langle r_{\perp}^2 \rangle < 1 \quad (126)$$

A material where $\mu < 1$ is called **diamagnetic**. Diamagnetism is due to the orbital angular momentum of the electrons.

Paramagnetism

Now we model free electrons. Electrons have spin, $s_z = \pm \frac{\hbar}{2}$. If a magnetic field is on, the energy of the system is $U = -\mu_z B = \pm \mu_B s B$, where $s = \pm 1$. The electron has a probability of being in one of these orientations of spin given by $p \propto \exp(-\beta U)$, where $\beta = \frac{1}{k_B T}$. Now the expected value of the dipole moment

$$\langle \mu \rangle = -\mu_B \langle s \rangle = \mu_B \frac{-e^{-\beta \mu_B B} + e^{\beta \mu_B B}}{e^{-\beta \mu_B B} + e^{\beta \mu_B B}} = \mu_B \tanh \left(\frac{\mu_B B}{k_B T} \right) \quad (127)$$

then,

$$\vec{M} = n \vec{\mu}_B \tanh \left(\frac{\mu_B B}{k_B T} \right) \approx \frac{n \mu_B^2}{k_B T} \vec{B} \quad (128)$$

thus $\chi = \frac{n \mu_B^2}{k_B T}$, and $\mu = 1 + 4\pi\chi > 1$. That $\mu > 1$ is called **paramagnetism**. Paramagnetism is caused due to the spin of the electrons. The tendency of the electrons is to align with the external magnetic field. We need partially occupied states as in conductors. States with double occupation do not present paramagnetism.

Itinerant Paramagnetism

The degeneracy of a system with two identical atoms is broken if there are interactions between the atoms. with two energy levels. With a lot of atoms there are also a lot of energy levels which can be thought as a continuum instead of discrete. A measure of the number of states with energy $E < \mathcal{E} < E + \Delta E$ is

$$\rho(E) \Delta E \quad (129)$$

where $\rho(E)$ is called the density of states. In an insulator there are bands which are filled. There exist gaps between bands. It costs a lot of energy for an electron to go from a full band to an empty band.

In a conductor the bands are partially filled. It costs little energy for an electron to roam in the band. If we fill some electrons in the band, there is a maximum energy called the *Fermi energy*.

If we turn on a \vec{B} , the electrons with spins up force the energy to go up by $\Delta U = \mu B$, and the electrons with spin down force the energy to drop by ΔU . The system is unstable, electrons with higher energy (spin up) change their spin in order to reduce their energy.

Induction and Lenz's Law

We begin with an oriented closed loop. Then from Faraday's law we get

$$\mathcal{E} = -\frac{1}{c} \frac{d}{dt} \Phi_B, \quad (130)$$

where \mathcal{E} [Energy] is the *electromotive force* and $\Phi_B = \oint d\vec{a} \cdot \vec{B}$ is the *magnetic flux*. The minus sign in the equation is known as *Lenz's law*. We need energy to overcome the electromotive force and establish a current. Lenz's law makes a circuit behave as a system with inertia.

Inductance and inductors

The inductance L is the ability of a body to oppose a change in the current. And it is defined through

$$V = L \frac{dI}{dt} \quad (131)$$

An inductor is a device that stores energy in a magnetic field.

Total energy of an inductor

The change in energy of a charge δq going from a to b in a circuit with a battery and a coil, against a potential produced by the battery is

$$\delta U_q = \phi(b)\delta q - \phi(a)\delta q = -V\delta q. \quad (132)$$

The charge has lost energy, from conservation of energy, the energy has been transferred to the coil

$$\delta U_{\text{coil}} = V\delta q = L \frac{dI}{dt} \delta q \quad (133)$$

and the power

$$\mathcal{P} = \frac{\delta U_{\text{coil}}}{\delta t} = L \frac{dI}{dt} \frac{\delta q}{\delta t} = LI \frac{dI}{dt} = \frac{1}{2} L \frac{d}{dt} I^2. \quad (134)$$

The total energy in the coil is then

$$U = \int dt \mathcal{P} = \frac{1}{2} LI^2. \quad (135)$$

Hysteresis

Inductance of a coaxial cable


We know that $H = \frac{2I}{rc}$, thus the energy density is $u = \frac{\vec{B} \cdot \vec{H}}{8\pi} = \mu \frac{4I^2}{8\pi r^2 c^2}$. Then the total energy is

$$U = \int d^3r \mu \frac{4I^2}{8\pi r^2 c^2} = \mu \frac{4I^2}{8\pi r^2 c^2} l 2\pi \int_a^b dr r \frac{1}{r^2} = \mu \frac{I^2}{c^2} \log\left(\frac{b}{a}\right) = \frac{1}{2} LI^2 \quad (136)$$

Therefore

$$L = \frac{2l \log(b/a)}{c^2} \mu \quad (137)$$

LR circuit



$$\begin{aligned} L \frac{dI}{dt} + RI &= 0 \\ \frac{dI}{dt} &= -\frac{R}{L} I \\ I &= I(0)e^{-t/\tau}, \quad \tau = L/R \end{aligned} \quad (138)$$

Impedance

Suppose an electric circuit as in . The battery provides $V(t) = V_0 \cos \omega t = \text{Re}(V_0 e^{i\omega t})$. According to KVL, the equation for I is

$$V_0 e^{i\omega t} = L \frac{dI}{dt} + RI \quad (139)$$

We look for solution of the form $I = I_P + I_H$, where I_H is the solution for the homogenous equation and I_P is a particular solution. Suppose $I_P = I_0 e^{-i\omega t}$ then

$$V = (-i\omega L + R)I_P \quad (140)$$

This resembles Ohm's law with an imaginary resistance $-i\omega L$, which means that L behaves as a resistance. We call this imaginary resistance *impedance*

$$Z_L = -i\omega L \quad (141)$$

We have a nother impedance related to the capacitance

$$Z_C = \frac{i}{\omega c} \quad (142)$$

Kirchhoff's circuit Laws

Kirchhoff's Current Law (KCL)

The sum of the currents in a node is zero.
(Conservation of charge)

Kirchhoff's Voltage Law (KVL)


The sum of the potential differences in a closed loop is zero.
(Valid on regions where $\vec{B} = 0 \Rightarrow \oint d\vec{l} \cdot \vec{E} = 0$)

Dictionary between classical mechanics and electical circuits

Classical mechanics		Electrical circuits	
Position	x	Charge	q
Velocity	$v = \frac{dx}{dt}$	Current	$I = \frac{dq}{dt}$
Spring constant	κ	Capacitance ⁻¹	$1/C$
Mass	m	Inductance	L
Air resistance	γ	Resistance	R
Force	F	Voltage	V
Newton's second law		Kirchhoff's voltage law	

Transmission Line

Consider a long coaxial cable. Suppose the cable is the union of coaxial cables with length Δx which have a capacitance $C = \Delta x / (2 \log(b/a))$ and inductance $L = (2 \log(b/a)/c^2) \Delta x$. The equivalent electric circuit is



The potential difference of the n -th inductor is

$$-V_n = \varphi_{n-1} - \varphi_n = L \frac{dI_n}{dt}, \quad (143)$$

where $f_n = f(x_n, t)$. The potential difference in the n -th capacitor is $\varphi_n = \frac{Q_n}{C}$ and has a current $\frac{dQ_n}{dt}$. Taking the derivative of the potential and using KCL we get

$$\frac{d\varphi_n}{dt} = \frac{1}{C} \frac{dQ_n}{dt} = \frac{1}{C} (I_n - I_{n+1}) \quad (144)$$

thus

$$\frac{d\varphi_{n-1}}{dt} - \frac{d\varphi_n}{dt} = L \frac{d^2 I_n}{dt^2} \quad (145)$$

substituting, we get

$$\frac{1}{C} (I_{n-1} - I_n) - \frac{1}{C} (I_n - I_{n+1}) = L \frac{d^2 I_n}{dt^2} \quad (146)$$

Finally we arrive at

$$\frac{\partial^2 I}{\partial x^2} - \frac{1}{v^2} \frac{\partial^2 I}{\partial t^2} = 0 \quad (147)$$

where $v^2 = \frac{(\Delta x)^2}{LC} = c^2$.

Mutual inductance

Consider a cylinder made by a material with permeability μ , radius a and length $l \gg a$. Roll a cable around the cylinder with $N \gg 1$ turns and set a current I_1 , call this configuration circuit 1. Roll a second cable around the cylinder, but isolated from the first one, with $N_2 \gg 1$ turns and a current I_2 , call it circuit 2. Then the magnetic field is

$$B = B_1 + B_2 = \frac{4\pi}{c} \frac{N_1}{l} I_1 + \frac{4\pi}{c} \frac{N_2}{l} I_2 \quad (148)$$

and the energy density

$$u = \frac{BH}{8\pi} = \frac{\mu}{8\pi} \left(\frac{4\pi}{cl} \right)^2 (N_1 I_1 + N_2 I_2)^2 \quad (149)$$

integrating over the volume of the cylinder we get the total energy

$$U = \frac{2\pi}{c^2 l^2} \mu (\pi a^2 l) (N_1 I_1 + N_2 I_2)^2 = \frac{2\pi^2 a^2}{c^2 l} \mu (N_1^2 I_1^2 + 2N_1 N_2 I_1 I_2 + N_2^2 I_2^2) \quad (150)$$

Changing notation

$$U = \frac{1}{2} \sum_{ij} L_{ij} I_i I_j, \quad i, j = 1, 2 \quad (151)$$

where

$$L_{ii} = \frac{4\pi^2 a^2}{lc^2} \mu N_i \text{ (no implicit sum), } L_{12} = L_{21} = \frac{4\pi^2 a^2}{lc^2} \mu N_1 N_2 \quad (152)$$

We realize that $L_{ij} = L_{ji}$ with $i \neq j$. The power of the coil is the derivative w.r.t. time, then

$$\frac{dU}{dt} = \mathcal{P} = \sum_{ij} I_i L_{ij} \dot{I}_j \quad (153)$$

Also, the power is the work per unit time we do over the charges over I_1 plus, the work over I_2

$$\mathcal{P} = \sum_{ij} \frac{dq_i}{dt} L_{ij} I_j \quad (154)$$

If we move a charge from one point to another, it will lose energy which is gained by the coil, then

$$\mathcal{P} = \sum_i \frac{dq_i}{dt} V_i \quad (155)$$

Comparing, we get

$$V_i = \sum_j L_{ij} I_j \quad (156)$$

We call L_{ii} (no implicit sum) the *autoinductance*. And L_{ij} , with $i \neq j$ *mutual inductance*.

Transformers

Consider the circuit in figure ... Since it is an open circuit $I_2 = 0$ and using eq.

$$V_1 = L_{11} \frac{dI_1}{dt} \quad (157)$$

$$V_2 = L_{21} \frac{dI_2}{dt} \quad (158)$$

Then

$$\frac{V_2}{V_1} = \frac{L_{21}}{L_{11}} \Rightarrow V_2 = \frac{L_{21}}{L_{11}} V_1 \Rightarrow V_2 = \frac{N_2}{N_1} V_1 \quad (159)$$

Now we close the circuit with a resistor R_C . We suppose a voltage source that provides with an amplitude V_{10} and a frequency ω . We write the voltage and the current represented by the real part of

$$V_1(t) = V_{10} e^{-i\omega t} \quad (160)$$

$$I_1(t) = I_{10} e^{-i\omega t} \quad (161)$$

After some time we can also write the current in the second circuit as

$$I_2(t) = I_{20} e^{-i\omega t} \quad (162)$$

Using eq. and Ohm's law, we get the set of couple equations

$$V_1 = -i\omega L_{11} I_1 - i\omega L_{12} I_2 \quad (163)$$

$$V_2 = -i\omega L_{21} I_1 - i\omega L_{22} I_2 \quad (164)$$

$$V_2 = R_C I_2 \quad (165)$$

Now suppose R_C is large in order to I_2 to be really small. Then we can dismiss the terms containing I_2 so we can use eq.

$$I_2 \approx \frac{L_{21}}{L_{11}} \frac{V_1}{R_C} \quad (166)$$

Another situation is when $R_C = 0 \Rightarrow V_2 = 0$. Then circuit 2 is in short-circuit. This gives constraints, I_1 must not be too large in order for circuit 2 not to burn.

More on Conservation Laws

We know that $\vec{j} \cdot \vec{E}$ is the power per unit time per unit volume that the fields give to a mechanical system. Therefore $-\vec{j} \cdot \vec{E}$ is power the system gives to the fields. We can interpret this as the source of energy density of the fields. Then the energy density of the fields satisfy the a continuity equation with sources

$$-\vec{j} \cdot \vec{E} = \partial_t u + \nabla \cdot \vec{S} \quad (167)$$

where u is the energy density and \vec{S} is the flow of energy. We can substitute \vec{j} in the LHS using Ampere's law, then

$$-j_i E_i = -\frac{c}{4\pi} \epsilon_{ijk} (\partial_j B_k) E_i + \frac{1}{4\pi} (\partial_t E_i) E_i \quad (168)$$

$$= -\frac{c}{4\pi} \partial_j (\epsilon_{ijk} B_k E_i) + \frac{c}{4\pi} \epsilon_{ijk} B_k \partial_j E_i + \frac{1}{8\pi} \partial_t (E_i E_i) \quad (169)$$

$$= \partial_j \left(\frac{c}{4\pi} \epsilon_{ijk} E_i B_k \right) - \frac{c}{4\pi} (\epsilon_{kji} \partial_j E_i) B_k + \frac{1}{8\pi} \partial_t (E_i E_i) \quad (170)$$

$$= \partial_j \left(\frac{c}{4\pi} \epsilon_{ijk} E_i B_k \right) + \partial_t \frac{1}{8\pi} (E_i E_i + B_i B_i) \quad (171)$$

$$= \partial_t u + \nabla \cdot \vec{S} \quad (172)$$

where

$$u = \frac{1}{8\pi} (E^2 + B^2) \quad (173)$$

$$\vec{S} = \frac{c}{4\pi} \vec{E} \times \vec{B} \quad (174)$$

In materials, the energy must satisfy a continuity equation with sources, but now the current density is the current that does not come from the materials. We call it the external current density \vec{j}^{ex} . So

$$-\vec{j}^{\text{ex}} \cdot \vec{E} = \partial_t u + \nabla \cdot \vec{S} \quad (175)$$

With a similar analysis but now using Maxwell equations in materials we arrive at

$$u = \frac{1}{8\pi} (\vec{E} \cdot \vec{D} + \vec{B} \cdot \vec{H}) \quad (176)$$

$$\vec{S} = \frac{c}{4\pi} \vec{E} \times \vec{H} \quad (177)$$

However these relations are only valid in no dispersive regions of transparency where we can assume both the permeability and permittivity as constants.

Conservation of momentum

From Newton's 2nd law, the Lorentz force is $F_{\text{em}} = \dot{p}$ where F_{em} is the momentum per unit time that the fields give to a mechanical system, therefore $-F_{\text{em}}$ is the source of

momentum per unit time given to the fields. Then the fields satisfy a continuity equation of momentum with source

$$-\vec{f} = -\rho\vec{E} - \frac{1}{c}\vec{j} \times \vec{B} = \partial_t \vec{g} + \nabla \cdot (-T) \quad (178)$$

where \vec{g} is the momentum density and $-T$ is the flux of momentum. Using the component representation of the LHS and using Gauss' Law and Ampere's to substitute ρ and \vec{j} we arrive at

$$\vec{g} = \frac{1}{4\pi c} \vec{E} \times \vec{B} \quad (179)$$

$$T = \frac{1}{4\pi} \left(\vec{E}\vec{E} + \vec{B}\vec{B} - \frac{1}{2}(E^2 + B^2)I \right) \quad (180)$$

Stress Tensor

T is a symmetric tensor known as the *Stress Tensor*.

$-T_{ij}$: j component of momentum that flows over direction i per unit time per unit area. The stress tensor is useful to work with problems where the force is needed but there is a discontinuity of the fields. From fluid mechanics we know that

$$\vec{F}_V = \int d\vec{a} \cdot T \quad (181)$$

is the force acting in the interior of a volume V . Imagine a surface that experience a pressure p due to is immeresed in a fluid. A chunk of surface da feels a force that push it to the inside

$$d\vec{f} = -pd\vec{a} \quad (182)$$

thus

$$\vec{F} = - \int d\vec{a} p = \int d\vec{a} \cdot (-pI) \quad (183)$$

Therefore in a fluid $T = -pI$. This result let us interpret terms in the diagonal. Negative terms are pressures, and positive terms are tensions per unit area.

Force on the plates of a capacitor

Consider a capacitor where we glue a charge Q on the left and $-Q$ on the right. What force is acting on the left plaque? We know that the field is discontinous. Outside the plaques the field is 0 and inside is $E = \frac{4\pi Q}{A}$. We could argue that the force on the left plaque is $F = \frac{4\pi Q^2}{A}$. However, imagine that we move lightly to the left plaque 1 a distance l . In magnitude $F_{me} = F_{e-m}$, then we have produced work $W = F_{me}l = F_{e-m}l = \Delta U$, where ΔU is the change in potential energy and $\Delta U = \Delta V u = (Al)\frac{E^2}{8\pi}$, where ΔV is the change in the volume and u is the energy density of the electric field. Therefore $F_{e-m} = \frac{2\pi Q^2}{A}$. Which is differente from the previous one, which is the correct one?. We use instead the momentum conservation equation. In a stationary situation $\partial_t \vec{g} = 0$ and $-\vec{f} = -\nabla \cdot T$ so $\vec{F} = \int_{\partial V} d\vec{a} \cdot T$. The only non zero component of T is

$$T_{xx} = \frac{1}{8\pi} E^2 \quad (184)$$

thus

$$F_x = \frac{1}{8\pi} E^2 A = \frac{2\pi Q^2}{A} = \frac{1}{2} QE \quad (185)$$

Therefore, the plaques are in tension.

Force acting on the wires of a solenoid

Consider a long solenoid of radius a and length $l \gg a$. We know that $\vec{B} = \frac{4\pi}{c} \frac{N}{l} I \hat{z}$, where $r < a$ and $\vec{B} = 0$ where $r > a$. We have a discontinuity in the field

Conservation of angular momentum

We can consider $-\vec{r} \times \vec{f} = -\vec{\tau}$ to be the source of angular momentum per unit time per unit volume of the fields.

$$-\vec{r} \times \vec{f} = \vec{r} \times (\partial_t \vec{g} - \nabla \cdot \vec{T}) \quad (186)$$

$$-\vec{\tau}_i = \partial_t (\vec{r} \times \vec{g})_i + \epsilon_{ijk} r_j \partial_l T_{lk} \quad (187)$$

$$= \partial_t (\vec{r} \times \vec{g})_i + \partial_l (\epsilon_{ijk} r_j T_{lk}) - \epsilon_{ijk} (\partial_l r_j) T_{lk} \quad (188)$$

$$= \partial_t (\vec{r} \times \vec{g})_i + \partial_l (\epsilon_{ijk} r_j T_{lk}) - \epsilon_{ijk} T_{jk} \quad (189)$$

$$-\vec{\tau} = \partial_t \vec{l} + \nabla \cdot \vec{M} \quad (190)$$

where

$$\vec{l} = \vec{r} \times \vec{g} \quad (191)$$

$$M_{li} = \epsilon_{ijk} r_j T_{lk} \quad (192)$$

Particular solution to the wave equation in 3d with initial conditions

Suppose we have a function Ψ satisfying the wave equation with initial conditions

$$\Psi(\vec{r}, 0) = 0, \quad \partial_t \Psi(\vec{r}, 0) = V(\vec{r}) \quad (193)$$

We apply the fourier transform to the spatial component of Ψ , $\Psi(\vec{r}, t) = \int \frac{d^3 k}{(2\pi)^3} \Psi_{\vec{k}}(t) e^{i\vec{k} \cdot \vec{r}}$ and to the wave equation which turns $\nabla \rightarrow i\vec{k}$, therefore

$$-k^2 \Psi_{\vec{k}}(t) - \frac{1}{c^2} \partial_t^2 \Psi_{\vec{k}}(t) = 0 \Rightarrow \frac{d^2 \Psi_{\vec{k}}(t)}{dt^2} = -c^2 k^2 \Psi_{\vec{k}}(t) \quad (194)$$

so we get a solution

$$\Psi_{\vec{k}}(t) = a_{\vec{k}} \cos(\omega_k t) + b_{\vec{k}} \sin(\omega_k t), \quad \omega_k = kc \quad (195)$$

The initial conditions in fourier space are

$$\Psi_{\vec{k}}(0) = 0 \Rightarrow a_{\vec{k}} = 0, \quad \frac{d\Psi_{\vec{k}}}{dt}(0) = V_{\vec{k}} = \int d^3 r V(\vec{r}) e^{-i\vec{k} \cdot \vec{r}} \Rightarrow b_{\vec{k}} = \frac{V(\vec{k})}{kc} \quad (196)$$

thus

$$\Psi(\vec{r}, t) = \int d^3 r' V(\vec{r}') \int \frac{d^3 k}{(2\pi)^3} \frac{\sin(kct)}{kc} e^{i\vec{k} \cdot (\vec{r} - \vec{r}')} \quad (197)$$

To evaluate the integral we use spherical coordinates. We choose the z axis in the $\vec{r} - \vec{r}'$ direction, so $\vec{k} \cdot (\vec{r} - \vec{r}') = k|\vec{r} - \vec{r}'| \cos \theta$, we also write $\vec{R} = |\vec{r} - \vec{r}'| \hat{\Omega}$

$$\Psi(\vec{r}, t) = -\frac{2}{4c} \frac{1}{2\pi} \int d^3 r' \frac{V(\vec{r}')}{R} \left[\delta(\underbrace{R+ct}_{>0}) - \delta(R-ct) \right] \quad (198)$$

We are only considering times greater than $t = 0$, where the initial conditions are, so we can drop $\delta(R + ct)$. And

$$\Psi(\vec{r}, t) = \frac{1}{4\pi c} \int d^3 r' \frac{V(\vec{r}')}{|\vec{r} - \vec{r}'|} \delta(ct - |\vec{r} - \vec{r}'|) \quad (199)$$

We replace the integral over r' with R so

$$\Psi(\vec{r}, t) = \frac{t}{4\pi} \int d\hat{\Omega} V(\vec{r} + ct\hat{\Omega}) = t \langle V(\vec{r} + ct\hat{\Omega}) \rangle_{\hat{\Omega}} \quad (200)$$

General solution of the wave equation in 3d with initial conditions

Suppose we have a function Ψ satisfying the wave equation with initial conditions

$$\Psi(\vec{r}, 0) = I(\vec{r}), \quad \partial_t \Psi(\vec{r}, 0) = 0 \quad (201)$$

If we define the function $\varphi(\vec{r}, t) = t \langle I(\vec{r} + ct\hat{\Omega}) \rangle_{\hat{\Omega}}$, which satisfies the wave equation with initial conditions

$$\varphi(\vec{r}, 0) = 0, \quad \partial_t \varphi(\vec{r}, 0) = I(\vec{r}) \quad (202)$$

we verify that $\Psi(\vec{r}, t) = \partial_t \varphi(\vec{r}, t)$. Finally, if we have a function satisfying the wave equation with generic initial conditions

$$\varphi(\vec{r}, 0) = I(\vec{r}), \quad \partial_t \varphi(\vec{r}, 0) = V(\vec{r}) \quad (203)$$

we arrive at the solution

$$\Psi(\vec{r}, t) = t \langle V(\vec{r} + ct\hat{\Omega}) \rangle_{\hat{\Omega}} + \frac{\partial}{\partial t} t \langle I(\vec{r} + ct\hat{\Omega}) \rangle_{\hat{\Omega}} \quad (204)$$

Turning on a field adiabatically

The sources in Maxwell's equations are ones that have existed always. In reality, the sources are turn on at some time and the turned off at some later time. Consider a source $f(t) = \int \frac{d\omega}{2\pi} f_\omega e^{-i\omega t}$, this tells us that a source is a superposition of waves that have osicillated from $t \rightarrow -\infty$ and will keep oscillating to $t \rightarrow +\infty$ which is not realistic.

Spatial and time translational invariance in the Green's function

Greens' functions have a traslation invariance if we move rigidly the source and the point of measure. Also, they have time translational invariance since the effect at time t of a source produced at time $t' < t$ only depend on the time difference. In this sense

$$G(\vec{r}, \vec{r}', t, t') = G(\vec{r} + \vec{a}, t + t_0, \vec{r}' + \vec{a}, t' + t_0) \quad (205)$$

In particular we can choose $\vec{a} = -\vec{r}'$ and $t_0 = -t'$ so now the Green's function only depend on just two variables $\vec{r} - \vec{r}'$ and $t - t'$

$$G = G(\vec{r} - \vec{r}', t - t') \quad (206)$$

Solution to the wave equation with sources in 3d

Suppose $\Psi(\vec{r}, t)$ satisfies the wave equation with sources

$$\nabla^2 \Psi - \frac{1}{c^2} \partial_t^2 \Psi = -4\pi f \quad (207)$$

We guess a Green's function G to be the solution of a wave equation with a point source at \vec{r}' that only live at time t' (obviously not real since sources/charges are conserved)

$$\left(\nabla^2 - \frac{1}{c^2}\partial_t^2\right) G(\vec{r}, t, \vec{r}', t') = -4\pi\delta(\vec{r} - \vec{r}')\delta(t - t') \quad (208)$$

so that

$$\Psi(\vec{r}, t) = \int d^3r' \int dt' G(\vec{r} - \vec{r}', t - t') f(\vec{r}', t') \quad (209)$$

Now we do $\vec{r} - \vec{r}' \rightarrow \vec{r}$ and $t - t' \rightarrow t$ and the equation for G reads

$$\left(\nabla^2 - \frac{1}{c^2}\partial_t^2\right) G(\vec{r}, t) = -4\pi\delta(\vec{r})\delta(t) \quad (210)$$

Applying fourier transform and solving we get

$$G_{\vec{k}\omega} = \frac{4\pi}{k^2 - \frac{\omega^2}{c^2}} \quad (211)$$

Therefore

$$G(\vec{r}, t) = -4\pi c^2 \int \frac{d^3k}{(2\pi)^3} \int \frac{d\omega}{2\pi} \frac{1}{(\omega + kc)(\omega - kc)} e^{i(\vec{k}\cdot\vec{r} - \omega t)} \quad (212)$$

In order to evalute integral over ω we turn ω into a complex number $\omega = \omega' + i\omega''$ in this way we can use Cauchy's residue theorem. However we have two poles $\omega = \pm kc$ we need to avoid. We invoke the causality principle: sources now ($t = 0$) do not affect the field in the past ($t < 0$). This means that the integral at times $t < 0$ must vanish. One way to warranty that the reponse is causal, is to assure there are no singularities inside the countour. If we want integral over the semicircle to vanish at $R \rightarrow \infty$ we need to have $e^{\omega''t} < 1$ this means $\omega'' > 0$. If we turn the sources adiabatically, that is we write $\omega \rightarrow \omega + i\eta$, $\eta > 0$, $\eta \rightarrow 0^+$, then we would have moved the poles down to the imaginary axis by η , that is $\omega = \pm kc - i\eta$. From Cauchy's theorem the integral over the upper semicircle will vanish since there are no poles enclosed.

If $t > 0$ we use Cauchy's theorem with a countour as in figure ... Then we get

$$G(\vec{r}, t) = -4\pi c^2 \int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot\vec{r}} \frac{-2\pi i}{2\pi} \left[\frac{e^{-ikct}}{2kc} + \frac{e^{ikct}}{-2kc} \right] = 4\pi c^2 \int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot\vec{r}} \frac{\sin(kct)}{k} \quad (213)$$

Integrating over k using spherical coordinates, and considering times $t > 0$ we finally get

$$G(\vec{r}, t) = \frac{c}{r} \delta(r - ct) = \frac{1}{r} \delta\left(t - \frac{r}{c}\right) \quad (214)$$

Returning to the prevoius notation $t \rightarrow t - t'$ and $\vec{r} \rightarrow \vec{r} - \vec{r}'$

$$G(\vec{r} - \vec{r}', t - t') = \frac{1}{|\vec{r} - \vec{r}'|} \delta(t - t_r) \quad (215)$$

where $t_r = t - \frac{|\vec{r} - \vec{r}'|}{c}$ is the time light takes going from the source to the point of observation. The solution for the potentials in Lorenz gauge is then

$$\varphi(\vec{r}, t) = \int d^3r' \frac{\rho(\vec{r}', t_r)}{|\vec{r} - \vec{r}'|} \quad (216)$$

$$\vec{A}(\vec{r}, t) = \frac{1}{c} \int d^3r' \frac{\vec{j}(\vec{r}', t_r)}{|\vec{r} - \vec{r}'|} \quad (217)$$

Cross section

Consider a plane wave with intensity I_i [Power/Area] hits a molecule. The molecule will be polarized and will lose energy with power P . Only the light that hit the molecule will be scattered so we define the effective size of the molecule called the cross section given by

$$\sigma = \frac{P}{I_i} \quad (218)$$

Mean free path

Why the sky is blue?

Suppose a molecule of the atmosphere is hit by monochromatic light from the sun with intensity $I = \frac{c}{8\pi}|E|^2$. This molecule will polarize in the direction of the electric field $\vec{p} = \alpha\vec{E}$. This will lead the molecule to oscillate with a frequency ω equal to frequency of the incident light. Therefore it will lose energy

$$P = \frac{\omega^4}{3c^3}|p|^2 = \frac{\omega^4}{3c^3}|\alpha|^2|E|^2 \quad (219)$$

Antenna

Consider a current source $I(t) = I_0 \cos(\omega t)$ connected to two metal rods of length $L/2$ as in figure ... to form an antenna. Let $I(z, t)$ be the current in the antenna. The current between the two rods at $z = 0$ is equal to the current from the source, and at $z = \pm L/2$ is zero since there the wires end

$$I(0, t) = I(t) = I_0 \cos(\omega t), \quad I(\pm L/2, t) = 0 \quad (220)$$

We make an ansatz of the form of the current that satisfies this behaviour

$$I(z, t) = I_0 \cos(\omega t) \cos\left(\pi \frac{z}{L}\right) \quad (221)$$

This form of I tells us that at fixed time the lower part, at $z < 0$, of the antenna has the opposite charge of the upper half, at $z > 0$, so it forms a dipole in one dimension

$$p_z(t) = \int dz \lambda(z, t) z \quad (222)$$

where $\lambda(z, t)$ is the charge density in one dimension, which satisfy the continuity equation in one dimension

$$\frac{\partial \lambda}{\partial t} + \frac{\partial I}{\partial z} = 0 \quad (223)$$

so $\lambda(z, t) = \frac{\pi I_0}{\omega L} \sin(\omega t) \sin(\frac{\pi}{L} z)$. Therefore the instantaneous electric dipole moment is

$$p_z(t) = \frac{2I_0 L}{\omega \pi} \sin(\omega t) = \text{Re}(p_z e^{i\omega t}) \quad (224)$$

so the amplitude of the dipole is $p_z = \frac{2I_0 L}{\omega \pi} i$. And

$$\vec{p} = \frac{2I_0 L}{\omega \pi} i(0, 0, 1) \quad (225)$$

This vector and not the instantaneous is what we need to replace in Larmor's formula to get the power loss

$$P = \frac{\omega^4}{3c^3} |\vec{p}|^2 = \frac{4\omega^2 L^2}{3c^3 \pi^2} I_0^2 \quad (226)$$

assuming of course $\lambda \gg L$. In an oscillating electrical circuit the power loss is

$$P = \langle VI \rangle = \langle RI^2(t) \rangle = \frac{1}{2} RI_0^2 \quad (227)$$

So, in the perspective of an electrical circuit we can describe the antenna as a resistor with resistance

$$R_{\text{rad}} = \frac{8\omega^2 L^2}{3\pi^2 c^3} \quad (228)$$