

Pure SU(3) lattice gauge theory in equilibrium

José Antonio García Hernández

March 11, 2024

1 Introduction

We study the 4-dimensional SU(3) lattice gauge theory with the compact formulation. In this formulation we consider compact link variables $U_\mu(x) \in \text{SU}(3)$ rather than the Lie algebra-valued fields $A_\mu(x)$. The compact variables $U_\mu(x)$ become the fundamental fields to be integrated over the functional integral. One great advantage of the compact formulation is that we no longer require to fix the gauge.

Link variables are oriented, so they naturally live in the links between neighboring points on the lattice, with spacing a . Thus, the variable $U_\mu(x)$ is defined in the link between points x and $x + a\hat{\mu}$ in the positive μ direction, see Fig. 1a. We define the variable between points x and $x + a\hat{\mu}$ pointing in the negative direction via


$$U_{-\mu}(x + a\hat{\mu}) \equiv U_\mu^\dagger(x), \quad (1)$$

see Fig. 1b.

Under a gauge transformation the link variables transform as

$$U'_\mu(x) = \Omega(x)U_\mu(x)\Omega^\dagger(x + a\hat{\mu}), \quad (2)$$

where $\Omega(x) \in \text{SU}(3)$. We refer to [1] for a complete discussion on lattice gauge theory.



(a) Link variable at point x in the μ direction. a is the lattice spacing. (b) Link variable at point $x + a\hat{\mu}$ in the $-\mu$ direction.

For $SU(N)$, the lattice gauge action reads

$$S[U] = \frac{\beta}{N} \sum_x \sum_{\mu < \nu} \text{Re Tr} [\mathbb{1} - U_{\mu\nu}(x)], \quad (3)$$

where $\beta = \frac{2N}{g^2}$. The plaquettes are defined by

$$U_{\mu\nu}(x) = U_\mu(x) U_\nu(x + \hat{\mu}) U_\mu^\dagger(x + \hat{\nu}) U_\nu^\dagger(x) \quad (4)$$

Sum of plaquette variables

$$S_P = \frac{1}{N} \sum_x \sum_{\mu < \nu} \text{Re Tr} [U_{\mu\nu}(x)] \quad (5)$$

we define

$$E_P = \frac{\langle S_P \rangle}{DV} \quad (6)$$

where $V = L^d$ is the volume of the lattice and $D = \frac{d(d-1)}{2}$ is the number of planes of rotation.

Staples

$$\Sigma_\mu(x) = \sum_{\nu \neq \mu} [U_\nu(x) U_\mu(x + \hat{\nu}) U_\nu^\dagger(x + \hat{\mu}) + U_\nu^\dagger(x - \hat{\nu}) U_\mu(x - \hat{\nu}) U_\nu(x + \hat{\mu} - \hat{\nu})] \quad (7)$$

The change in the action by a local update when changing $U_\mu(x) \rightarrow U'_\mu(x)$ is

$$\Delta S = -\frac{\beta}{N} \text{Re Tr} [(U'_\mu(x) - U_\mu(x)) \Sigma_\mu^\dagger] \quad (8)$$

In $SU(3)$ the $SU(2)$ elements are of the form

$$R = \begin{pmatrix} r_{11} & r_{12} & 0 \\ r_{21} & r_{22} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad S = \begin{pmatrix} s_{11} & 0 & s_{12} \\ 0 & 1 & 0 \\ s_{21} & 0 & s_{22} \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 0 & 0 \\ 0 & r_{11} & r_{21} \\ 0 & r_{21} & r_{22} \end{pmatrix}. \quad (9)$$

2 Algorithms

2.1 Metropolis

1. Given some gauge field configuration, we go through all the lattice points in a lexicographic way. At each point x in the μ direction we generate a random $SU(2)$ matrix $U'_\mu(x)$.
2. We compute the sum of the staples and compute the change of the action according to eq. (8). We generate a uniform random number $r \in [0, 1)$ and accept the change if $r \leq p$, where p is

$$p = \min(1, \exp(-\Delta S)). \quad (10)$$

2.2 Heatbath

The heatbath algorithm was first implemented by Creutz for the SU(2) theory [5] and generalized to SU(N) theories by Cabbibo and Marinari [6]. The idea for implementing the algorithm in SU(N) is to iterate over a set of SU(2) subgroups in SU(N) and apply the heatbath algorithm to these SU(2) elements.

1. Given some field configuration at a point x and direction μ , we find the sum of staples $\Sigma_\mu(x)$ in eq. (7).
2. We compute $W = U_\mu(x)\Sigma_\mu^\dagger(x)$ and define W_2 as the 2×2 submatrix of W that has the same block structure as R (or S or T).
3. W_2 is a complex 2×2 matrix which is generally not an element of SU(2). As described in Ref. [7], we generate a matrix proportional to an SU(2) element from W_2 as follows

$$V = \frac{1}{2} \left[W_2 - W_2^\dagger + \mathbf{1} \text{Tr } W_2^\dagger \right] \quad (11)$$

4. We compute the determinant of V and take $\xi = \sqrt{\det V}$
5. We follow the original heatbath implementation of Creutz. We generate a uniform random number $r \in [\exp(-\frac{4}{3}\beta\xi), 1]$ and compute

$$x_0 = 1 + \frac{\log r}{\frac{2}{3}\beta\xi}. \quad (12)$$

Now we generate a uniform random number $u \in [0, 1)$ and accept x_0 if $u > 1 - \sqrt{1 - x_0^2}$. Repeat this step until a x_0 is accepted.

6. We now generate a vector $\vec{x} = (x_1, x_2, x_3)$ in the unit 2-sphere uniformly distributed. This may be accomplished as follows. We generate three random numbers $\vec{r} = (r_1, r_2, r_3)$ in the interval $[-1, 1]$ and accept them if they are inside the unit 3-sphere. Otherwise we generate other three numbers until they are accepted. Once they are accepted we take

$$\vec{x} = \sqrt{1 - x_0^2} \frac{\vec{r}}{|\vec{r}|}. \quad (13)$$

7. We have generated the elements of a SU(2) matrix X

$$X = \begin{pmatrix} x_0 + ix_1 & x_2 + ix_3 \\ -x_2 + ix_3 & x_0 - ix_1 \end{pmatrix} \quad (14)$$

8. We take

$$R_2 = X \frac{V^\dagger}{\xi}. \quad (15)$$

And transform this 2×2 matrix to the same block structure as the 3×3 matrix R (or S or T).

9. We go to step 2 and repeat the process to generate S (or T) but this time taking $W = RU_\mu(x)\Sigma_\mu(x)$ (or $W = SRU_\mu(x)\Sigma_\mu(x)$).

10. The new link element $U'_\mu(x)$ is

$$U'_\mu(x) = T S R U_\mu(x). \quad (16)$$

3 Results

We present simulation results for the 4-dimensional $SU(3)$ lattice gauge field theory. We implement two different local update algorithms, namely, Metropolis and Heatbath. Figure 2 shows the energy density E_P as a function of $\beta = \frac{6}{g^2}$. There is a maximum of the slope at $5 < \beta < 6$, however this is not an indication of a phase transition.

4 Acknowledgements

The author thanks Dr. Wolfgang Bietenholz for funding (in the form of meals) this research. Thanks to Dr. Urs Gerber for providing a recipe for implementing the heatbath algorithm for the $U(1)$ gauge theory at an early stage of this research. Finally, I want to thank Ariana Grande, The Weeknd and Polyphia for giving the right inspiration while working on this paper. Without their music I could have easily gone crazy.

References

- [1] C. Gattringer and C.B. Lang, *Quantum Chromodynamics on the Lattice: An Introductory Presentation*, Lect. Notes Phys. 788 (Springer, Berlin Heidelberg 2010).
- [2] N. Metropolis *et al.*, *Equation of State Calculations by Fast Computing Machines*, J. Chem. Phys. **21** (1953) 1087.

SU(3), d = 4, L = 6




Figure 2: Energy density vs. $\beta = \frac{6}{g^2}$. We simulated the SU(3) pure gauge theory in four dimensions using the Metropolis and Heatbath algorithms.

- [3] R.J. Glauber, *Time-Dependent Statistics of the Ising Model*, J. Math. Phys. **4** (1963) 294.
- [4] U. Gerber, *Heatbath Algorithm for the 2d U(1) Model*. Informal Notes. Universidad Nacional Autónoma de México, 2015.
- [5] M. Creutz, *Monte Carlo Study of the Quantized SU(2) Gauge Theory*. Phys. Rev. D **21** (1980) 2308-2315.
- [6] N. Cabbibo, E. Marinari, *A New Method for Updating SU(N) Matrices in Computer Simulations of Gauge Theories*, Phys. Lett. B **119** (1982) 387-390.
- [7] A.D. Kennedy, B.J. Pendleton, *Improved Heatbath Method for Monte Carlo Calculations in Lattice Gauge Theories*, Phys. Lett. **156B** (1985) 393-399.