

2d $\lambda\phi^4$ theory

This code simulates the 1-component $\lambda\phi^4$ theory.

Requisites

- A Fortran compiler: gfortran
- Makefile installed
- Use Linux

Theory

Let $\phi(x) \in \mathbb{R}$ be a field. The Lagrangian density in Euclidian time reads

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi(x) \partial_\mu \phi(x) + \frac{1}{2} m^2 \phi(x)^2 + \frac{1}{4} \lambda \phi(x)^4,$$

where $\lambda \geq 0$, for the potential to be bounded from below. The continuum action reads

$$S[\phi] = \int dx^d \mathcal{L}(\phi, \partial_\mu \phi).$$

With the lattice regularization

$$\begin{aligned} \phi(x) &\rightarrow \phi_x \\ \partial_\mu \phi(x) &\rightarrow \frac{\phi_{x+a\hat{\mu}} - \phi_x}{a} \\ \int dx^d &\rightarrow \sum_x a^d \end{aligned}$$

The unit vectors are defined as

$$\hat{1} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad \hat{2} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

On the lattice the action takes the form

$$S[\phi] = \sum_x a^d \left\{ \frac{1}{2} \sum_\mu \left(\frac{\phi_{x+a\hat{\mu}} - \phi_x}{a} \right)^2 + \frac{1}{2} m^2 \phi_x^2 + \frac{1}{4} \lambda \phi_x^4 \right\}.$$

In lattice units we set $a = 1$.

Expanding the action through the lattice

$$\begin{aligned}
S[\phi] &= \frac{1}{2} \sum_{\mu=1}^2 (\phi_{(1,1)+\hat{\mu}} - \phi_{(1,1)})^2 + \frac{1}{2} m^2 \phi_{(1,1)}^2 + \frac{1}{4} \lambda \phi_{(1,1)}^4 \\
&+ \frac{1}{2} \sum_{\mu=1}^2 (\phi_{(1,2)+\hat{\mu}} - \phi_{(1,2)})^2 + \frac{1}{2} m^2 \phi_{(1,2)}^2 + \frac{1}{4} \lambda \phi_{(1,2)}^4 \\
&\vdots \\
&+ \frac{1}{2} \sum_{\mu=1}^2 (\phi_{(i,j-1)+\hat{\mu}} - \phi_{(i,j-1)})^2 + \frac{1}{2} m^2 \phi_{(i,j-1)}^2 + \frac{1}{4} \lambda \phi_{(i,j-1)}^4 \\
&\vdots \\
&+ \frac{1}{2} \sum_{\mu=1}^2 (\phi_{(i-1,j)+\hat{\mu}} - \phi_{(i-1,j)})^2 + \frac{1}{2} m^2 \phi_{(i-1,j)}^2 + \frac{1}{4} \lambda \phi_{(i-1,j)}^4 \\
&+ \frac{1}{2} \sum_{\mu=1}^2 (\phi_{(i,j)+\hat{\mu}} - \phi_{(i,j)})^2 + \frac{1}{2} m^2 \phi_{(i,j)}^2 + \frac{1}{4} \lambda \phi_{(i,j)}^4 \\
&\vdots \\
&= \dots \\
&+ \frac{1}{2} \left[(\phi_{(i+1,j-1)} - \phi_{(i,j-1)})^2 + (\phi_{(i,j)} - \phi_{(i,j-1)})^2 \right] + \frac{1}{2} m^2 \phi_{(i,j-1)}^2 + \frac{1}{4} \lambda \phi_{(i,j-1)}^4 \\
&\vdots \\
&+ \frac{1}{2} \left[(\phi_{(i,j)} - \phi_{(i-1,j)})^2 + (\phi_{(i-1,j+1)} - \phi_{(i-1,j)})^2 \right] + \frac{1}{2} m^2 \phi_{(i-1,j)}^2 + \frac{1}{4} \lambda \phi_{(i-1,j)}^4 \\
&+ \frac{1}{2} \left[(\phi_{(i+1,j)} - \phi_{(i,j)})^2 + (\phi_{(i,j+1)} - \phi_{(i,j)})^2 \right] + \frac{1}{2} m^2 \phi_{(i,j)}^2 + \frac{1}{4} \lambda \phi_{(i,j)}^4 \\
&\vdots
\end{aligned}$$

If we update the field ϕ at site $x = (i, j)$, that is, $\phi_x \rightarrow \phi'_x$, the action becomes

$$\begin{aligned}
S[\phi'] = & \dots \\
& + \frac{1}{2} \left[\left(\phi_{(i+1,j-1)} - \phi_{(i,j-1)} \right)^2 + \left(\phi'_{(i,j)} - \phi_{(i,j-1)} \right)^2 \right] + \frac{1}{2} m^2 \phi_{(i,j-1)}^2 + \frac{1}{4} \lambda \phi_{(i,j-1)}^4 \\
& \vdots \\
& + \frac{1}{2} \left[\left(\phi'_{(i,j)} - \phi_{(i-1,j)} \right)^2 + \left(\phi_{(i-1,j+1)} - \phi_{(i-1,j)} \right)^2 \right] + \frac{1}{2} m^2 \phi_{(i-1,j)}^2 + \frac{1}{4} \lambda \phi_{(i-1,j)}^4 \\
& + \frac{1}{2} \left[\left(\phi_{(i+1,j)} - \phi'_{(i,j)} \right)^2 + \left(\phi_{(i,j+1)} - \phi'_{(i,j)} \right)^2 \right] + \frac{1}{2} m^2 \phi_{(i,j)}^2 + \frac{1}{4} \lambda \phi_{(i,j)}^4 \\
& \vdots
\end{aligned}$$

Therefore the change in the action $\Delta S = S[\phi'] - S[\phi]$ is

$$\begin{aligned}
\Delta S = & \frac{1}{2} \left[\left(\phi'_{(i,j)} - \phi_{(i,j-1)} \right)^2 - \left(\phi_{(i,j)} - \phi_{(i,j-1)} \right)^2 \right] \\
& + \frac{1}{2} \left[\left(\phi'_{(i,j)} - \phi_{(i-1,j)} \right)^2 - \left(\phi_{(i,j)} - \phi_{(i-1,j)} \right)^2 \right] \\
& + \frac{1}{2} \left[\left(\phi_{(i+1,j)} - \phi'_{(i,j)} \right)^2 - \left(\phi_{(i+1,j)} - \phi_{(i,j)} \right)^2 \right] \\
& + \frac{1}{2} \left[\left(\phi_{(i,j+1)} - \phi'_{(i,j)} \right)^2 - \left(\phi_{(i,j+1)} - \phi_{(i,j)} \right)^2 \right] \\
& + \frac{1}{2} m^2 \left(\phi_{(i,j)}'^2 - \phi_{(i,j)}^2 \right) + \frac{1}{4} \lambda \left(\phi_{(i,j)}'^4 - \phi_{(i,j)}^4 \right) \\
= & \frac{1}{2} \left[\left(\phi_{(i,j)}'^2 - \phi_{(i,j)}^2 \right) - 2\phi_{(i,j-1)} \left(\phi'_{(i,j)} - \phi_{(i,j)} \right) \right] \\
& + \frac{1}{2} \left[\left(\phi_{(i,j)}'^2 - \phi_{(i,j)}^2 \right) - 2\phi_{(i-1,j)} \left(\phi'_{(i,j)} - \phi_{(i,j)} \right) \right] \\
& + \frac{1}{2} \left[\left(\phi_{(i,j)}'^2 - \phi_{(i,j)}^2 \right) - 2\phi_{(i+1,j)} \left(\phi'_{(i,j)} - \phi_{(i,j)} \right) \right] \\
& + \frac{1}{2} \left[\left(\phi_{(i,j)}'^2 - \phi_{(i,j)}^2 \right) - 2\phi_{(i,j+1)} \left(\phi'_{(i,j)} - \phi_{(i,j)} \right) \right] \\
& + \frac{1}{2} m^2 \left(\phi_{(i,j)}'^2 - \phi_{(i,j)}^2 \right) + \frac{1}{4} \lambda \left(\phi_{(i,j)}'^4 - \phi_{(i,j)}^4 \right) \\
= & 2 \left(\phi_{(i,j)}'^2 - \phi_{(i,j)}^2 \right) \\
& - 2 \left(\phi'_{(i,j)} - \phi_{(i,j)} \right) \left(\phi_{(i-1,j)} + \phi_{(i,j-1)} + \phi_{(i+1,j)} + \phi_{(i,j+1)} \right) \\
& + \frac{1}{2} m^2 \left(\phi_{(i,j)}'^2 - \phi_{(i,j)}^2 \right) + \frac{1}{4} \lambda \left(\phi_{(i,j)}'^4 - \phi_{(i,j)}^4 \right)
\end{aligned}$$

Metropolis Algorithm

We generate a change in the field, $\phi_x \rightarrow \phi'_x$ and accept it with probability

$$p = \min(1, \exp(-\Delta S)).$$