

Supporting Information

The non-random assembly of functional motifs in plant-pollinator networks worldwide

Jose B. Lanuza | Alfonso Allen-Perkins | Ignasi Bartomeus

TABLES

Table S1 List of plant-pollinator network studies.

Table S2 Type of plant traits used in this study.

FIGURES

Figure S1 Over and under-estimated motif frequencies.

Figure S2 Over and under-estimated functional groups on motif positions.

Figure S3 Plant trait composition of functional groups.

Figure S4 Plant dendrogram with functional groups.

Table S1. List of studies ordered by author with the year of publication, number of contributed networks and digital object identifier

First author	Year	Number of networks	Country	DOI
Arroyo-Correa	2019	3	New Zealand	https://doi.org/10.1111/1365-2745.13332
Bartomeus	2008	6	Spain	https://doi.org/10.1007/s00442-007-0946-1
Bartomeus	2015	16	Spain	https://github.com/ibartomeus/BeeFunData
Bundgaard	2003	1	Denmark	Unpublished, Master thesis
Burkle	2013	1	United States	https://doi.org/10.1126/science.1232728
Dicks	2002	2	England	https://doi.org/10.1046/j.0021-8790.2001.00572.x
Dupont	2003	3	Denmark	https://doi.org/10.1111/j.1365-2656.2008.01501.x
Elberling	1999	1	Sweden	https://doi.org/10.1111/j.1600-0587.1999.tb00507.x
Fang	2008	1	China	https://doi.org/10.1111/1749-4877.12190
Inouye	1988	1	United States	https://doi.org/10.1111/j.1442-9993.1988.tb00968.x
Kaiser-Bunbury	2017	8	Seychelles	https://doi.org/10.1038/nature21071
Kaiser-Bunbury	2011	6	Seychelles	https://doi.org/10.1111/j.1365-2745.2010.01732.x
Kaiser-Bunbury	2010	2	Mauritius	https://doi.org/10.1016/j.ppees.2009.04.001
Lundgren	2005	1	Denmark (Greenland)	https://doi.org/10.1657/1523-0430(2005)037[0514:TDAHCW]2.0.CO;2
Olesen	2002	2	Mauritius and Portugal (Azores)	https://doi.org/10.1046/j.1472-4642.2002.00148.x
Peralta	2006	4	Argentina	https://doi.org/10.1111/ele.13510
Small	1976	1	Japan	/13960/t4km08d21
Souza	2017	1	Brazil	https://doi.org/10.1111/1365-2745.12978

Table S2. Traits used to delimit the different plant functional groups divided in quantitative and categorical traits.

Quantitative traits		Categorical traits	
Type	Traits	Type	Traits
Vegetative	Plant height (m)	Vegetative	Lifepan
Floral	Flower width (mm)	Vegetative	Life form
Floral	Flower length (mm)	Floral	Flower shape
Floral	Inflorescence width (mm)	Floral	Flower symmetry
Floral	Style length (mm)	Reproductive	Autonomous selfing
Floral	Ovules per flower	Reproductive	Compatibility system
Floral	Flowers per plant	Reproductive	Breeding system
Reproductive	Autonomous selfing (fruit set)		




Figure S1. Comparison of motif frequencies between empirical and simulated networks grouped by average path length (plots a, b, c and d) as determined in Simmons et al. (2020) without considering singletons. This is shown with the mean percentage of motif frequencies in empirical networks that were over the motif frequencies of the simulated ones (percentiles). This was done by network (light blue dots) and then averaged for all networks (black dots with error bars that correspond to the standard deviation).





Figure S2. Heatmap indicating under- and over- representation of pollinator and plant functional groups in the different motif positions after removing non-robust links (singletons). The different motif positions are divided by the average path length clasification by Simmons et al. (2020). The superior dendrogram indicates the differences across groups with the more separated groups showing larger differences.

Plant functional group composition

A) Qualitative variables



B) Quantitative variables




Figure S3. Plant functional group composition separated in qualitative and quantitative variables. Panel A) shows the percentage of the different categories within trait represented with different colours for each functional group. Plot B) shows the radar plot of the different quantitative variables standardized on the same scale also coloured with the same patterns of colours as qualitative variables per cluster or functional group. Group 1 corresponds to short-lived selfers; group 2 to small outcrossing perennials; group 3 to self-incompatible perennials with large flowers; group 4 to tall plants with small unisexual flowers; and, group 5 to short-lived outcrossers with long zygomorphic flowers.

Plant functional groups




Figure S4. Hierarchical clustering dendrogram with the branches coloured by the optimal number of clusters (5). The labels of the subgroup of species ($N = 524$) used in this study are coloured in black in order to show the evenness of the distribution of the species across clusters. The rest of species labels are omitted for visualization purposes ($N = 982$).