

working title Compatibility system and stigma size are the main predictors of heterospecific pollen effect

Jose B. Lanuza, Ignasi Bartomeus, Tia-Lynn Ashman, Romina Rader * ^{1,2,3}

¹US Environmental Protection Agency, Office of Research and Development, National Health and Environmental Effects Research Laboratory, Atlantic Ecology Division, 27 Tarzwell Drive Narragansett, RI, 02882, USA

²Big Name University, Department of R, City, BN, 01020, USA

³Estacion Biologica de Donana (EBD-CSIC), E-41092 Sevilla, Spain

* corresponding author: barragansljose@gmail.com

Pollinator sharing can have negative consequences for plant fitness with the arrival of foreign pollen. However, the costs of heterospecific pollen are not yet well understood. We conducted a glasshouse experiment to understand how phylogenetic relatedness and plant traits mediate the impacts of heterospecific pollen transfer. We conducted 4XXXX crosses by experimentally transferring pollen (50% and 100% ratio) with reciprocal crosses between 10 species belonging to three different families: Brassicaceae, Solanaceae and Convolvulaceae. Seed set was used as proxy of plant fitness. We found that for 65% of the treatments with 50% mix reduced seed set. Moreover, the reduction in seed set was dependent on the degree of relatedness and reproductive traits of the pollen recipient and not the pollen donor. Our results show that certain traits, particularly compatibility system, are critical in understanding the costs of heterospecific pollen.

Keywords: heterospecific pollen, plant reproduction, fitness, interspecific competition, phylogenetic distance.

INTRODUCTION

In most ecosystems, plant species normally coexist and share their floral visitors with other species Waser et al. (1996). From the plants' perspective, pollinator sharing can be positive for some plants Carvalheiro et al. (2014) or negative for others Pauw (2013), depending on the facilitation gradient. An increasing number of visits often correlates with higher chances of fertilization Engel and Irwin (2003). However this is not always the case, among these possible flower visitors there are also nectar robbers and pollen thieves Inouye (1980); Magrach et al. (2017). Receiving both sufficient quantity and quality deposited on the stigma is thus highly relevant to the pollination success of the plant Aizen and Harder (2007).

34 By visiting many plant species, many pollinators are responsible for conspecific pollen loss and the
35 transport of foreign pollen, both of which can have important detrimental effects on species fitness
36 Morales and Traveset (2008); Ashman and Arceo-Gómez (2013); Arceo-Gómez and Ashman (2016).
37 Foreign pollen arrival can play an important role in plant species fitness but outcomes are variable and
38 appear to be context dependent as there is not always a decrease in fitness Morales and Traveset (2008).
39 Some of this variation is likely due to the enormous variability of foreign pollen transferred across
40 systems ranging from 0 to 75 percent. However, most studies report ranges of heterospecific pollen
41 between 0 and 20 percent of the total pollen load Bartomeus et al. (2008) Montgomery and Rathcke
42 (2012); Ashman and Arceo-Gómez (2013); Fang and Huang (2013), yet even these relatively low
43 amounts of heterospecific pollen transferred can decrease fitness greatly Thomson et al. (1982). While
44 we now have some understanding of the impacts of heterospecific pollen quantity, we have less
45 understanding of other factors that could be driving the variation in impacts upon fitness. Ashman and
46 Arceo-Gómez (2013) postulated the first predictive framework that identifies a need to understand how
47 plant traits might mediate heterospecific pollen effect, whereby mating system and pollen size were
48 predicted to potentially mediate the impact of foreign pollen transfer on plant fitness. This concept is
49 supported by specific case studies, such as Tong and Huang (2016) that demonstrate an asymmetrical
50 effect in 6 species of *Pedicularis* whereby the pollen of long styled species was able to grow the full
51 length of the style on short styled species but not vice versa. While this suggests that the impacts of
52 heterospecific pollen may differ among pollen donor and recipient, few studies have been conducted to
53 ascertain whether this pattern is in fact a general trend or to identify the extent to which other plant
54 traits are critical to heterospecific pollen impacts.

55 Plant traits are crucial to understand heterospecific pollen effect but the multifactorial nature of the
56 traits that are involve in the pollen-pistil interaction make difficult to unravel what are the main traits
57 in driving the effect. These traits can be seen from a male perspective of both donor and recipient
58 where pollen size, pollen aperture number and pollen allelopathy are key components to understand the
59 outcome of foreign pollen arrival (REFS). In Ashman and Arceo-Gómez (2013) small pollen is
60 predicted to cause a greater fitness decrease, although this can be true there are also other possibilities
61 to consider which can obscure a predictive framework like big pollen can clogg small stigmas with fewer
62 pollen grains, bigger stigmas are less likely to be clogged by small pollen grains and bigger pollen can

63 outcompete smaller pollen grains due to faster pollen tube growth rate Williams and Rouse (1990).
64 Moreover, to understand the different mechanical or chemical effects of pollen also the female traits of
65 the pollen recipient have to be considered, from the literature these main traits are: stigma size, style
66 length, number of ovules, incompatibility system and finally a structural trait such as flower
67 morphology Montgomery and Rathcke (2012); Ashman and Arceo-Gómez (2013); Tong and Huang
68 (2016). For example, greater stigmatic area is positively correlated with greater heterospecific pollen
69 deposition Montgomery and Rathcke (2012) and therefore possibly with an increase in negative effect.
70 For species that are self-incompatible the barriers towards heterospecific pollen are stronger than
71 self-compatible species Ashman and Arceo-Gómez (2013). Nonetheless, an effect of foreign pollen is a
72 bit obscured by the variability within species, however species that are strong selfers or strong
73 outcrossers have less variability in mating systems and predictions of effect could be more realistic (see
74 figure 1 from Whitehead et al. (2018)). Although past research has progress in the understanding of
75 what traits can mediate the effect as we have shown here, there are multiple traits involved and
76 multiple possible scenarios still to be explored empirically for a full understanding of the importance of
77 heterospecific pollen effect in nature.

78 Closely related species are more likely to have similar traits (Refs? Brown and Mitchell (2001)
79 Arceo-Gómez et al. (2016) Tong and Huang (2016)). Several studies predict that the impact of HP
80 transfer is likely to be greater for closely related species (Ashman and Arceo-Gómez (2013)). Few
81 studies however, have focused on the impacts of heterospecific pollen of distantly related species
82 Thomson et al. (1982) Galen and Gregory (1989) Neiland and Wilcock (1999). Yet, most insects and
83 most stigmas have been found to carry multiple species of foreign pollen with little attention to degree
84 of relatedness (Arceo-Gómez and Ashman (2016); Fang and Huang (2013); also cite studies from pollen
85 transfer networks here such as...). Traditionally, studies have overlook the effect of closely related
86 species but in a natural context both can occur and different degree of relatedness should be considered.

87 Further, a majority of plant species are generalist and thus receive visits from multiple different
88 pollinators. Given these are generally the ones that receive greater loads of heterospecific pollen Fang
89 and Huang (2013) and unrelated species are more likely to coexist with other species due to less niche
90 overlap (Ref), understanding the role of foreign pollen from distantly related species thus deserves
91 greater attention in understanding coexistence blah blahXXXXX refs.. Notwithstanding, the effect of

heterospecific pollen of far and close related species at community level remains to be explored beyond single pairwise interactions. Moreover, heterospecific pollen studies in nature have the complexity added great environmental variability which can lead to confounding interpretations and diverse floral structures and some of them complex such as Asteraceae species, which involve tedious work to be able to study foreign pollen effect. For this reason, we investigated how floral reproductive traits and relatedness mediate the impact of heterospecific transfer by asking the following research questions : To what extent do (i) floral reproductive traits and (ii) relatedness, mediate the impacts of heterospecific pollen on seed set. We do this by creating an artificial co-flowering community with 10 species belonging to three different families with different traits.

-Would like to add here floral complexity

-Environmental variability of studies on the field

METHODS

The study was conducted in a glasshouse at University of New England (Armidale, Australia) from November 2017 to March 2018. Rooms were temperature controlled depending on the requirements of the species with day and night temperature differences. The species selected (**Table 1**) belonged to three different families, Solanaceae, Brassicaceae and Convolvulaceae. The criteria of species/family selection was based on close/distant related species (see phylogenetic tree for relatedness fig 1)I would explain more the bauty of our nested dessign to ensure close and far distance simultaneously, heterogeneous traits, low structural flower complexity and fast life cycle. For the purpose of the experiment all the species where considered as pollen recipient and as pollen donor (see interaction matrix, fig 2). Species were watered once or twice per day and fertilized weekly (NPK 23: 3.95: 14).

Table 1

Family	Genus	Species
Brassicaceae	Brassica	Brassica rapa
Brassicaceae	Brassica	Brassica oleracea

Family	Genus	Species
Brassicaceae	Eruca	Eruca versicaria
Brassicaceae	Sinapis	Sinapis alba
Convolvulaceae	Ipomoea	Ipomoea aquatica
Convolvulaceae	Ipomoea	Ipomoea purpurea
Solanaceae	Capsicum	Capsicum annuum
Solanaceae	Petunia	Petunia integrifolia
Solanaceae	Solanum	Solanum lycopersicum
Solanaceae	Solanum	Solanum melongena

114 Hand-pollination

115 Foreign pollen effect was studied through two different treatments, one with 50% conspecific pollen and
116 50% heterospecific pollen and a second one with 100% foreign pollen (N=10) this second I don't get,
117 maybe explain it's utility.. Therefore, 180 different combinations were performed with N=10 per
118 combination. Seed set was the proxy of effect for all our treatments. Moreover, hand cross pollination
119 (between individuals of the same species), hand self pollination, apomixis (bagged emasculated flowers)
120 and natural selfing were tested for each species (N=10). For the treatments with foreign pollen and
121 hand cross pollination, flowers were emasculated the day prior anthesis and hand pollinated next day
122 with a toothpick. Hand-pollination was conducted with 3-4 gentle touches on the stigma surface. The
123 mixes of pollen were realized on an eppendorf based on the pollen counts made with Neubauer chamber
124 (each anther was counted 4 times for 20 different anthers per species)-IB explain better and give a bit
125 more of detail. In order to confirm that the treatments applied were 50-50 percent pollen, for each focal
126 species the total stigmatic load of pollen was counted from one donor of each family (N=3).

127 Traits and evolutive distance

128 The traits measured for each species were pollen per anther, number of ovules, stigma width and length
129 and stigmatic area, style width and length, ovary width and length. Moreover stigma type explain was
130 tested. Pollen was counted for 20 anthers of each species with 4 replicates per sample with an

hemocytometer. Previously, anthers were squashed on a known solution with the pippete tip and
homogeneize with a vortex for 30 seconds. Ovule number was counted with the help of an
stereomicroscope and a small grid over a petri dish from 15 randomly selected flowers. The different
morphometrical traits were measured with a digital stereomicrospe. Levels of self incompatibility were
estimated by dividing the fruit set of hand self pollination by hand cross pollination Lloyd and Schoen
(1992).

Analysis

We used the statistical language R (R Core Team 2018) for all our analyses. Differences of seed set
between treatments and hand cross pollination for each species was tested through mixed linear models.
For the following analysis we scaled the values of seed production for all the species with mean 0 and sd
of 1. To test the effect of heterospecific pollen, we substracted to the seed set of hand cross pollination
the seed set of heterospecific pollen treatments. In order to see correlations between heterospecific
pollen effect and traits we performed Mantel test between the matrix of heterospecific pollen effect and
the distance matrix of each trait (euclidean distances). Moreover, Mantel test was also conducted
between heterospecific pollen effect and the square root of the matrix of phylogenetic distance due to
improvement in the statistical power (Letten & Cornwell 2014). all is here, but I would break it by
questions and give a bit more detail, to avoid overwhelm the reader We explored also the relations
between traits and heterospecific pollen effect through generalized mixed models where the response
variable was heterospecific pollen effect, the independent variable the different traits and the random
effects the different treatments per species [Here I think you should think if this controls for the non
independency of donors and recipients. I think not. Maybe look onto matrix regresions?). Moreover,
pairwise evolutive distances were calculated with MEGA7 for two kinds of markers: 1) Internal
transcribed spacer (ITS) and 2) ribulose-bisphosphate carboxylase (RBCL). The sequences of interest
were downloaded from NCBI GenBank and the phylogenetic tree constructed by maximum likelihood
with MEGA7. Make a section on how you contrsucted phylogeny.

I would explain three test. 0) treatment effects with GLM's, 1) Mantels: relative effects, 2) GLM's or
matrix models: Absolute effects and explain them in three independent paragraphs including a
rationale of why

RESULTS

Results of hand cross pollination, self hand pollination, natural selfing and apomixis are presented in **Table 2**. Heterospecific pollen reduced seed set significantly with the 50-50% heterospecific pollen treatments for 65% of the pairwise interactions $p < 0.05$. Across families we found a very similar effect but when species were looked at species level they responded differently even within the same family rephrase and maybe test statistically?, for instance for two species of the Brassicaceae family *Brassica oleracea* and *Eruca versicaria* we found very contrasting effects of foreign pollen where for the first one, all donors reduce seed set significantly and for the second, just two species did out of nine. The 100% foreign pollen treatments barely produced seeds or fruits and just for *Sinapis alba* we did not find significant differences between the hand cross pollination and one treatment with pollen from a confamilial- IB Unclear. Solanaceae species with berry fruit type developed small fruits or even normal fruits in some cases under which treatment. *S. lycopersicum* seems to produce small fruits (35% of the treatments) independently of pollen and pollen donor due to also apomictic treatments did, never normal size. *C. annuum* produced some fruits (9%) of both small and normal size and finally *S. melongena* produced seedless normal fruits with just confamilial pollen (3%), for both species seems that fruit formation was induced by pollen on the stigma because of lack of fruit production with treatments that tested for apomixis. clarify this descriptive statistics part- Also a figure with a summary of the treatments effect would be cool, or at least in the appendices

178 **Table 2.** Percentage of seeds produced per ovule for the ten species used in the experiment. The
179 treatments presented are hand cross pollination, hand self pollination, natural selfing and apomixis
180 (emasculated flowers). turn into a figure somehow?

Species	Cross	Self	Natural_selfing	Apomixis
Brassica oleracea	32.06897	0.0000000	0.00000	0
Brassica rapa	44.97041	0.0000000	0.00000	0
Eruca versicaria	23.75000	0.4166667	0.00000	0
Sinapis alba	43.33333	48.3333333	5.00000	15
Ipomoea aquatica	40.00000	30.0000000	20.00000	0
Ipomoea purpurea	31.66667	86.6666667	31.66667	0
Capsicum annuum	100.00000	66.2240664	23.48548	0
Petunia integrifolia	100.00000	24.7727273	0.00000	0
Solanum lycopersicum	90.38043	43.4782609	70.00000	0
Solanum melongena	60.47525	87.9702970	21.56436	0




Figure 1: Barplot with the different treatments that provide information of the reproductive biology of the ten species. The y axis is the proportion of ovules converted to seed in percentage. The different treatments (N=10) which are presented in the legend are, hand cross pollination, hand self pollination, natural selfing and apomixis. More information about these treatments can be found in Methods and Appendices.

Mantel test indicates that a possible?? It exists! correlation exist between heterospecific pollen effect and the evolutive relative distances, for ITS and RBCL markers we had r coefficients of 0.29 and 0.25 respectively $p < 0.05$ think on a figure - maybe using NMDS. Moreover, Mantel test indicates that also a possible?? correlation between stigma width and stigma type exist (stats??). Trait correlations were also explored with GLMM

I have done it at the moment just for Compatibility system Also I have to fix from mixed linear model to GLMM, just realize that




Figure 2: The effect of heterospecific pollen (scaled see set) is represented in function of the compatibility system (self/cross*100) for the the different species. Each coulored dot represents the interaction of a focal species with a different pollen donor.

188 [Compatibility index don't multiply per 100 from Lloyd](#)

189 **DISCUSSION**

190 Discussion

191 Herbs vs tress, annual vs perennial... Many flowers vs few flowered species; structural composition on
192 a system

193 What are the implications of the findings?

194 Ideas about pollen size in heterospecific pollen effect. (still have to develop it more...)

195 Let's classify pollen size in three groups in order to understand the interaction between pollen donor
196 and recipient: 1) Donor pollen size < Recipient pollen size 2) Donor pollen size = Recipient pollen size
197 3) Donor pollen size > Recipient pollen size

198 Now I try to develop each part

199 1) Donor pollen size < Recipient pollen size

200 Effect:

- 201 • Donor's pollen could clogg the stigma
- 202 • Chemical inhibition

203 Traits associated with bigger pollen of the recipient:

- 204 • Recipient's pollen have faster pollen tube growth (example with my data)
- 205 • Reduction in number of ovules (Also with my species)
- 206 • Big differences in pollen size can be traduced in low relatednes therefore less likely of pollen
207 germination on a far related stigma.

208 2) Donor pollen size = Recipient pollen size

- 209 • Very relatedness dependant this point

210 • Similar probabilities of taken space on the stigma

211 3) Donor pollen size > Recipient pollen size

212 Effect:

213 -In small stigmas big pollen grains can occupy great part of the stigmatic area.

214 -small pollen grains can get embedded

215 IB: Think also on using tree analysis to test if hp effect depends on complex trait combinations. Tree
216 analysis are great when two different strategies lead to the same outcome. This would never been pick
217 up by GLMs. The r package is party{} . You can see an example applied to birds is Sol et al 2010
218 Science. Ask me if you want more details or code examples.

219 CONCLUSIONS

220 ACKNOWLEDGEMENTS

221 REFERENCES

222 Aizen, M. A., and L. D. Harder. 2007. Expanding the limits of the pollen-limitation concept: Effects of
223 pollen quantity and quality. *Ecology* 88:271–281.

224 Arceo-Gómez, G., and T.-L. Ashman. 2016. Invasion status and phylogenetic relatedness predict cost
225 of heterospecific pollen receipt: Implications for native biodiversity decline. *Journal of Ecology*
226 104:1003–1008.

227 Ashman, T.-L., and G. Arceo-Gómez. 2013. Toward a predictive understanding of the fitness costs of
228 heterospecific pollen receipt and its importance in co-flowering communities. *American Journal of*

229 Botany 100:1061–1070.

230 Bartomeus, I., J. Bosch, and M. Vilà. 2008. High invasive pollen transfer, yet low deposition on native
 231 stigmas in a carpobrotus-invaded community. *Annals of Botany* 102:417–424.

232 Carvalho, L. G., J. C. Biesmeijer, G. Benadi, J. Fründ, M. Stang, I. Bartomeus, C. N.
 233 Kaiser-Bunbury, M. Baude, S. I. Gomes, V. Merckx, and others. 2014. The potential for indirect effects
 234 between co-flowering plants via shared pollinators depends on resource abundance, accessibility and
 235 relatedness. *Ecology letters* 17:1389–1399.

236 Engel, E. C., and R. E. Irwin. 2003. Linking pollinator visitation rate and pollen receipt. *American*
 237 *Journal of Botany* 90:1612–1618.

238 Fang, Q., and S.-Q. Huang. 2013. A directed network analysis of heterospecific pollen transfer in a
 239 biodiverse community. *Ecology* 94:1176–1185.

240 Inouye, D. W. 1980. The terminology of floral larceny. *Ecology* 61:1251–1253.

241 Lloyd, D. G., and D. J. Schoen. 1992. Self-and cross-fertilization in plants. i. functional dimensions.
 242 *International Journal of Plant Sciences* 153:358–369.

243 Magrach, A., J. P. González-Varo, M. Boiffier, M. Vilà, and I. Bartomeus. 2017. Honeybee spillover
 244 reshuffles pollinator diets and affects plant reproductive success. *Nature ecology & evolution* 1:1299.

245 Montgomery, B. R., and B. J. Rathcke. 2012. Effects of floral restrictiveness and stigma size on
 246 heterospecific pollen receipt in a prairie community. *Oecologia* 168:449–458.

247 Morales, C. L., and A. Traveset. 2008. Interspecific pollen transfer: Magnitude, prevalence and
 248 consequences for plant fitness. *Critical Reviews in Plant Sciences* 27:221–238.

249 Pauw, A. 2013. Can pollination niches facilitate plant coexistence? *Trends in ecology & evolution*
 250 28:30–37.

251 R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for

252 Statistical Computing, Vienna, Austria.

253 Thomson, J. D., B. J. Andrews, and R. Plowright. 1982. The effect of a foreign pollen on ovule
254 development in *diervilla lonicera* (caprifoliaceae). *New Phytologist* 90:777–783.

255 Tong, Z.-Y., and S.-Q. Huang. 2016. Pre-and post-pollination interaction between six co-flowering
256 *pedicularis* species via heterospecific pollen transfer. *New Phytologist* 211:1452–1461.

257 Waser, N. M., L. Chittka, M. V. Price, N. M. Williams, and J. Ollerton. 1996. Generalization in
258 pollination systems, and why it matters. *Ecology* 77:1043–1060.

259 Whitehead, M. R., R. Lanfear, R. J. Mitchell, and J. D. Karron. 2018. Plant mating systems often vary
260 widely among populations. *Frontiers in Ecology and Evolution* 6:38.

261 Williams, E., and J. Rouse. 1990. Relationships of pollen size, pistil length and pollen tube growth
262 rates in *rhododendron* and their influence on hybridization. *Sexual Plant Reproduction* 3:7–17.

264 **List of Figures**

265	1	Barplot with the different treatments that provide information of the reproductive biology	
266		of the ten species. The y axis is the proportion of ovules converted to seed in percentage.	
267		The different treatments (N=10) which are presented in the legend are, hand cross	
268		pollination, hand self pollination, natural selfing and apomixis. More information about	
269		these treatments can be found in Methods and Appendices.	10
270	2	The effect of heterospecific pollen (scaled see set) is represented in function of the	
271		compatibility system (self/cross*100) for the the different species. Each coulored dot	
272		represents the interaction of a focal species with a different pollen donor.	11