

working title Compatibility system and stigma size of pollen recipient as main predictors of heterospecific pollen effect

Jose B. Lanuza, Ignasi Bartomeus, Tia-Lynn Ashman, Romina Rader * ^{1,2,3}

¹US Environmental Protection Agency, Office of Research and Development, National Health and Environmental Effects Research Laboratory, Atlantic Ecology Division, 27 Tarzwell Drive Narragansett, RI, 02882, USA

²Big Name University, Department of R, City, BN, 01020, USA

³Estacion Biologica de Donana (EBD-CSIC), E-41092 Sevilla, Spain

* corresponding author: barragansljose@gmail.com

Pollinator sharing can have negative consequences for species fitness with the arrival of foreign pollen. However, the costs of heterospecific pollen are not yet well understood. For this reason, we have conducted a glasshouse experiment where we try to understand how phylogenetic relatedness and the different traits of these species are involved in this process. We experimentally crossed 10 species belonging to three different families: Brassicaceae, Solanaceae and Convolvulaceae. Overall, more than 4000 crosses were done and seed set and pollen tubes were considered as proxy of effect. We found that for all species foreign pollen (50% or less) reduced seed set. Moreover, the seed set reduction is not dependent on the degree of relatedness of the pollen donor. However, the effect is governed by the degree of relatedness and the traits of the species recipient. Our results show that the outcome of heterospecific pollen deposition is determined in greater degree by the traits of the pollen recipient than the pollen donor and that certain traits such as compatibility system are crucial to understand the costs of heterospecific pollen.

Keywords: heterospecific pollen, plant reproduction, fitness, interspecific competition, phylogenetic distance.

INTRODUCTION

Paragraph 1 General idea to our concept

In natural systems plant species normally coexist and share their floral visitors with other species Bascompte et al. (2003). This pollinator sharing from the plant perspective at the pre-pollination stage can be negative due to competition Pauw (2013) or positive due to facilitation Carvalheiro et al. (2014). Once the floral visitor has arrived to the flower, pollen deposition on the stigma can take place and hence ovule fertilization. An increasing number of visits generally correlates with higher chances of

35 fertilization Engel and Irwin (2003). However this is not always the case, among these possible flower
36 visitors we find also nectar robbers and pollen thieves Inouye (1980) and the quality of pollen that is
37 deposit on the stigma is also highly relevant to the pollination succes Aizen and Harder (2007).
38 Moreover, other less study issues in the pollination process are conspecific pollen loss and the arrival of
39 foreign pollen which can have important detrimental effects on species fitness Morales and Traveset
40 (2008) Ashman and Arceo-Gómez (2013).

41 **Paragraph 2** Introducing topic and knowledge gap

42 Recent studies have advanced in the ecological understanding of heterospecific pollen effect Morales and
43 Traveset (2008) Ashman and Arceo-Gómez (2013) Arceo-Gómez and Ashman (2016). A general
44 overview of foreign pollen arrival is that it can play an important role on species fitness but seems to be
45 context dependent and not always produce a decrease in fitness Morales and Traveset (2008). Part of
46 this unpredictability is due to the enormous variability of foreing pollen transferred in nature, where
47 levels between 0 and 75 percent are seen, but most commonly values ranges between 0 and 20 percent
48 of the total pollen load, being the generalist species the ones that receive greater loads of heterospecific
49 pollen Bartomeus et al. (2008) Montgomery and Rathcke (2012) Ashman and Arceo-Gómez (2013)
50 Fang and Huang (2013). Although heterospecific pollen quantity is fundamental to understand the
51 outcome of the interaction so is the different traits of both pollen donor and recipient. Ashman and
52 Arceo-Gómez (2013) postulated the first predictive framework for traits of heterospecific pollen effect,
53 where different traits such as compatibility system and pollen size among others seems to be crucial to
54 understand foreing pollen effect. Moreover, in Tong and Huang (2016) an assymetric effect was shown
55 in a crossing experiment between 6 species of the genus *Pedicularis* where the pollen of long styled
56 species was able to grow the full length of the style on short styled species but not viceversa. Despite
57 these recent caveats, we still lack empirical evidence to affirm what are the main traits that drive
58 heterospecific pollen effect for both pollen donor and recipient at seed production level. Interestingly, to
59 comprehend how these traits interact is also crucial to look at the phylogenetic relatedness of the
60 species. There is a considerable amount of literature of crosses between close related species Brown and
61 Mitchell (2001) Arceo-Gómez et al. (2016) Tong and Huang (2016) but few works focused on
62 heterospecific pollen of far related species. Although the effect of close related species is predicted to be
63 greater Ashman and Arceo-Gómez (2013) the presence of pollen of non related species on multiple

species Arceo-Gómez and Ashman (2016) and the higher chances to coexist with a species that has less niche overlap (Ref) make foreign pollen from far related species also an important subject of study in order to understand the importance of heterospecific pollen in natural systems. Notwithstanding, the effect of heterospecific pollen of far and close related species at community level remains to be explored.

Paragraph 3 Expanding ideas with examples

Interestingly, incompatibility system seems to play an important role in foreign pollen effect where species that are self incompatible would have stronger barriers towards heterospecific pollen than self compatible species. The type of incompatibility, sporophytic or gametophytic is related with the place of pollen recognition where the former take place at the stigma level and the latter occurs within the style, this last late acting pollen recognition mechanism is associated with greater negative effect (REF). Remarkably, there is a great variability in mating systems across populations Whitehead et al. (2018) and therefore predict an effect of foreign pollen is a bit obscured by the variability within species, however species that are strong selfers or strong outcrossers have less variability in mating systems and predictions of effect could be more realistic (see figure 1 from Whitehead et al. (2018)).

Paragraph 4 Introducing our experiment

The great environmental variability in natural systems and complexity of floral structures make heterospecific pollination studies a daunting task. Moreover, variation in sampling effort have been shown to be determinant to characterize pollen transfer interactions Arceo-Gómez et al. (2018). Although plant-pollinator network and pollen network studies can give a first picture of the importance of foreign pollen is necessary to address how its effect is shaped with both traits and relatedness of the species. For this reason, in this study we have created an artificial co-flowering community with 10 species belonging to three different families where we try to test the following questions: 1) Does heterospecific pollen reduce seed set, if so, 2) Does heterospecific pollen effect depend on the relatedness of the species, 3) Does heterospecific pollen effect depend on any floral trait?

88 METHODS

89 The study was conducted in a glasshouse at University of New England (Armidale, Australia) from
90 November 2017 to March 2018. Rooms were temperature controlled depending on the requirements of
91 the species with day and night temperature differences. The species selected (Table 1) belonged to
92 three different families, Solanaceae, Brassicaceae and Convolvulaceae. The criteria of species/family
93 selection was based on close/distant related species (see phylogenetic tree for relatedness fig 1),
94 heterogeneous traits, low structural flower complexity and fast life cycle. For the purpose of the
95 experiment all the species were considered as pollen recipient and as pollen donor (see interaction
96 matrix, fig 2). Species were watered once or twice per day and fertilized weekly (NPK 23: 3.95: 14).
97 Brown and Mitchell 2001 could be a good paper to explain why we pick seed set as a proxy and not
98 fruit set. We cannot see changes on it, losing information with it.

99 Hand-pollination

100 Foreign pollen effect was studied through two different treatments, one with 50% conspecific pollen and
101 50% heterospecific pollen and a second one with 100% foreign pollen (N=10). Seed set was the proxy of
102 effect (see Brown and Mitchell 2001, for differences in effect between seed set and fruit set) and “pollen
103 tubes”. Moreover, hand cross pollination, hand self pollination, apomixis (bagged emasculated flowers)
104 and natural selfing were tested (N=10). Flowers were emasculated the day prior anthesis and hand
105 pollinated next day with a toothpick. Hand-pollination was realized with 3-4 gentle touches on the
106 surface of the stigma. The mixes of pollen were performed on an eppendorf based on the pollen counts
107 made with Neubauer chamber (each anther was counted 4 times for 20 different anthers per species).

108 Evolutive distance

109 Two types of evolutive distances were calculated with MEGA7 for two kinds of markers: 1) Internal
110 transcribed spacer (ITS) and 2) ribulose-bisphosphate carboxylase (RBCL)

111 Traits

112 The traits measured for each species were pollen per anther, number of ovules, stigma width and length

113 and stigmatic area, style width and length, ovary width and length. Moreover stigma type was tested.
114 Pollen was counted for 20 anthers of each species with 4 replicates per sample with an hemocytometer.
115 Previously anthers were squashed on a known solution with the pippete tip and homogeneize with a
116 vortex for 30 seconds. Ovule number was counted with the help of an stereomicroscope and a small
117 grid over a petri dish from 15 randomly selected flowers. The different morphometrical traits were
118 measured with XXXX. Levels of self incompatibility were estimated by dividing the the fruit set of
119 hand self pollination by hand cross pollination

120 We used the statistical language **R** (R Core Team 2018) for all our analyses. These were implemented in
121 dynamic **rmarkdown** documents using **knitr** (Xie 2014, 2015, 2018) and **rmarkdown** (Allaire et al.
122 2018) packages. All the multilevel models were fitted with **lme4** (Bates et al. 2015).

123 **RESULTS**

124 **DISCUSSION**

125 Discussion

- 126 1. What are the implications of the findings?

127 CONCLUSIONS

128 ACKNOWLEDGEMENTS

129 REFERENCES

- 130 Aizen, M. A., and L. D. Harder. 2007. Expanding the limits of the pollen-limitation concept: Effects of
131 pollen quantity and quality. *Ecology* 88:271–281.
- 132 Allaire, J., Y. Xie, J. McPherson, J. Luraschi, K. Ushey, A. Atkins, H. Wickham, J. Cheng, and W.
133 Chang. 2018. Rmarkdown: Dynamic documents for r.
- 134 Arceo-Gómez, G., C. Alonso, T.-L. Ashman, and V. Parra-Tabla. 2018. Variation in sampling effort
135 affects the observed richness of plant–plant interactions via heterospecific pollen transfer: Implications
136 for interpretation of pollen transfer networks. *American journal of botany* 105:1601–1608.
- 137 Arceo-Gómez, G., and T.-L. Ashman. 2016. Invasion status and phylogenetic relatedness predict cost
138 of heterospecific pollen receipt: Implications for native biodiversity decline. *Journal of Ecology*
139 104:1003–1008.
- 140 Arceo-Gómez, G., R. A. Raguso, and M. A. Geber. 2016. Can plants evolve tolerance mechanisms to
141 heterospecific pollen effects? An experimental test of the adaptive potential in clarkia species. *Oikos*
142 125:718–725.
- 143 Ashman, T.-L., and G. Arceo-Gómez. 2013. Toward a predictive understanding of the fitness costs of
144 heterospecific pollen receipt and its importance in co-flowering communities. *American Journal of*
145 *Botany* 100:1061–1070.
- 146 Bartomeus, I., J. Bosch, and M. Vilà. 2008. High invasive pollen transfer, yet low deposition on native
147 stigmas in a carpobrotus-invaded community. *Annals of Botany* 102:417–424.
- 148 Bascompte, J., P. Jordano, C. J. Melián, and J. M. Olesen. 2003. The nested assembly of plant–animal

mutualistic networks. *Proceedings of the National Academy of Sciences* 100:9383–9387.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67:1–48.

Brown, B. J., and R. J. Mitchell. 2001. Competition for pollination: Effects of pollen of an invasive plant on seed set of a native congener. *Oecologia* 129:43–49.

Carvalho, L. G., J. C. Biesmeijer, G. Benadi, J. Fründ, M. Stang, I. Bartomeus, C. N. Kaiser-Bunbury, M. Baude, S. I. Gomes, V. Merckx, and others. 2014. The potential for indirect effects between co-flowering plants via shared pollinators depends on resource abundance, accessibility and relatedness. *Ecology letters* 17:1389–1399.

Engel, E. C., and R. E. Irwin. 2003. Linking pollinator visitation rate and pollen receipt. *American Journal of Botany* 90:1612–1618.

Fang, Q., and S.-Q. Huang. 2013. A directed network analysis of heterospecific pollen transfer in a biodiverse community. *Ecology* 94:1176–1185.

Inouye, D. W. 1980. The terminology of floral larceny. *Ecology* 61:1251–1253.

Montgomery, B. R., and B. J. Rathcke. 2012. Effects of floral restrictiveness and stigma size on heterospecific pollen receipt in a prairie community. *Oecologia* 168:449–458.

Morales, C. L., and A. Traveset. 2008. Interspecific pollen transfer: Magnitude, prevalence and consequences for plant fitness. *Critical Reviews in Plant Sciences* 27:221–238.

Pauw, A. 2013. Can pollination niches facilitate plant coexistence? *Trends in ecology & evolution* 28:30–37.

R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Tong, Z.-Y., and S.-Q. Huang. 2016. Pre-and post-pollination interaction between six co-flowering

- 172 pedicularis species via heterospecific pollen transfer. *New Phytologist* 211:1452–1461.
- 173 Whitehead, M. R., R. Lanfear, R. J. Mitchell, and J. D. Karron. 2018. Plant mating systems often vary
174 widely among populations. *Frontiers in Ecology and Evolution* 6:38.
- 175 Xie, Y. 2014. Knitr: A comprehensive tool for reproducible research in R. *in* V. Stodden, F. Leisch,
176 and R. D. Peng, editors. *Implementing reproducible computational research*. Chapman; Hall/CRC.
- 177 Xie, Y. 2015. *Dynamic documents with R and knitr*. 2nd editions. Chapman; Hall/CRC, Boca Raton,
178 Florida.
- 179 Xie, Y. 2018. Knitr: A general-purpose package for dynamic report generation in R.

