

working title Compatibility system and stigma size are the main predictors of heterospecific pollen effect

Jose B. Lanuza, Ignasi Bartomeus, Tia-Lynn Ashman, Romina Rader ^{*} 1,2,3

¹US Environmental Protection Agency, Office of Research and Development, National Health and Environmental Effects Research Laboratory, Atlantic Ecology Division, 27 Tarzwell Drive Narragansett, RI, 02882, USA

²Big Name University, Department of R, City, BN, 01020, USA

³Estacion Biologica de Donana (EBD-CSIC), E-41092 Sevilla, Spain

^{*} corresponding author: barragansljose@gmail.com

Pollinator sharing can have negative consequences for plant fitness with the arrival of foreign pollen. However, the costs of heterospecific pollen are not yet well understood. We conducted a glasshouse experiment to understand how phylogenetic relatedness and plant traits mediate the impacts of heterospecific pollen transfer. We conducted 4XXXX crosses by experimentally transferring pollen (50% and 100% ratio) with reciprocal crosses between 10 species belonging to three different families: Brassicaceae, Solanaceae and Convolvulaceae. Seed set was used as proxy of plant fitness. We found that for 65% of the treatments with 50% mix reduced seed set. Moreover, the reduction in seed set was dependent on the degree of relatedness and reproductive traits of the pollen recipient and not the pollen donor. Our results show that certain traits, particularly compatibility system, are critical in understanding the costs of heterospecific pollen.

Keywords: heterospecific pollen, plant reproduction, fitness, interspecific competition, phylogenetic distance.

INTRODUCTION

In most ecosystems, plant species normally coexist and share their floral visitors with other species Waser et al. (1996). From the plants' perspective, pollinator sharing can be positive for some plants Carvalheiro et al. (2014) or negative for others Pauw (2013), depending on the facilitation gradient. An increasing number of visits often correlates with higher chances of fertilization Engel and Irwin (2003). However this is not always the case, among these possible flower visitors there are also nectar robbers and pollen thieves Inouye (1980); Magrach et al. (2017). Receiving both sufficient quantity and quality deposited on the stigma is thus highly relevant to the pollination success of the plant Aizen and Harder (2007).

34 By visiting many plant species, many pollinators are responsible for conspecific pollen loss and the
35 transport of foreign pollen, both of which can have important detrimental effects on species fitness
36 Morales and Traveset (2008); Ashman and Arceo-Gómez (2013); Arceo-Gómez and Ashman (2016).
37 Foreign pollen arrival can play an important role in plant species fitness but outcomes are variable and
38 appear to be context dependent as there is not always a decrease in fitness Morales and Traveset (2008).
39 Some of this variation is likely due to the enormous variability of foreign pollen transferred across
40 systems ranging from 0 to 75 percent. However, most studies report ranges of heterospecific pollen
41 between 0 and 20 percent of the total pollen load Bartomeus et al. (2008) Montgomery and Rathcke
42 (2012); Ashman and Arceo-Gómez (2013); Fang and Huang (2013), yet even these relatively low
43 amounts of heterospecific pollen transferred can decrease fitness greatly Thomson et al. (1982). While
44 we now have some understanding of the impacts of heterospecific pollen quantity, we have less
45 understanding of other factors that could be driving the variation in impacts upon fitness. Ashman and
46 Arceo-Gómez (2013) postulated the first predictive framework that identifies a need to understand how
47 plant traits might mediate heterospecific pollen effect, whereby mating system and pollen size were
48 predicted to potentially mediate the impact of foreign pollen transfer on plant fitness. This concept is
49 supported by specific case studies, such as Tong and Huang (2016) that demonstrate an asymmetrical
50 effect in 6 species of *Pedicularis* whereby the pollen of long styled species was able to grow the full
51 length of the style on short styled species but not vice versa. While this suggests that the impacts of
52 heterospecific pollen may differ among pollen donor and recipient, few studies have been conducted to
53 ascertain whether this pattern is in fact a general trend or to identify the extent to which other plant
54 traits are critical to heterospecific pollen impacts.

55 Plant traits are crucial to understand heterospecific pollen effect but the multifactorial nature of the
56 traits that are involve in the pollen-pistil interaction make difficult to unravel what are the main traits
57 in driving the effect. On the one hand the main predictive traits of effect from the donor perspective
58 are pollen size, pollen allelopathy and pollen apertures (REFS). In Ashamn 2013 small pollen is
59 predicted to cause a greater fitness decrease, although this can be true there are also other possibilities
60 to consider which can obscure a predictive framework like big pollen can clogg small stigmas with fewer
61 pollen grains, bigger stigmas are less likely to be clogged by small pollen grains and bigger pollen can
62 outcompete smaller pollen grains due to faster pollen tube growth rate (REF). On the other hand the

main recipient traits that can drive heterospecific pollen effect are: stigma size, style length, number of ovules, incompatibility system and flower morphology (REFS). In first place, greater stigmatic area is related with greater amount of heterospecific pollen (REF) and therefore possibly a greater negative effect (REF). Species with shorter styles are more likely to be vulnerable to heterospecific pollen effect (REF). For species that are self-incompatible have stronger barriers to heterospecific pollen than self-compatible species Ashman and Arceo-Gómez (2013). Nonetheless, an effect of foreign pollen is a bit obscured by the variability within species, however species that are strong selfers or strong outcrossers have less variability in mating systems and predictions of effect could be more realistic (see figure 1 from Whitehead et al. (2018)). All in all, there is a need to deepen in the understanding of the relative importance of these traits, due to we still do not know what are the main traits in driving the effect with all these possible combinations that can occur in nature.

Comments on it

Species closely related are more likely to have similar traits XXXXXXXXXX. (Refs? Brown and Mitchell (2001) Arceo-Gómez et al. (2016) Tong and Huang (2016)). Several studies predict that the impact of HP transfer is likely to be greater for closely related species (Ashman and Arceo-Gómez (2013)). Few studies however, have focused on the impacts of heterospecific pollen of distantly related species Thomson et al. (1982) Galen and Gregory (1989) Neiland and Wilcock (1999). Yet, most insects and most stigmas have been found to carry multiple species of foreign pollen with little attention to degree of relatedness (Arceo-Gómez and Ashman (2016); Fang and Huang (2013) ; also cite studies from pollen transfer networks here such as...).

Further, a majority of plant species are generalist and thus receive visits from multiple different pollinators. Given these are generally the ones that receive greater loads of heterospecific pollen Fang and Huang (2013) and unrelated species are more likely to coexist with other species due to less niche overlap (Ref), understanding the role of foreign pollen from distantly related species thus deserves greater attention in understanding coexistence blah blahXXXXX refs.. Notwithstanding, the effect of heterospecific pollen of far and close related species at community level remains to be explored beyond single pairwise interactions. In this study we investigated how floral reproductive traits and relatedness mediate the impact of HP transfer by asking the following research questions : To what extent do (i)

floral reproductive traits and (ii) relatedness, mediate the impacts of heterospecific pollen on seed set. We do this by creating an artificial co-flowering community with 10 species belonging to three different families with different traits.

METHODS

The study was conducted in a glasshouse at University of New England (Armidale, Australia) from November 2017 to March 2018. Rooms were temperature controlled depending on the requirements of the species with day and night temperature differences. The species selected (**Table 1**) belonged to three different families, Solanaceae, Brassicaceae and Convolvulaceae. The criteria of species/family selection was based on close/distant related species (see phylogenetic tree for relatedness fig 1)I would explain more the bauty of our nested dessign to ensure close and far distance simultaneously, heterogeneous traits, low structural flower complexity and fast life cycle. For the purpose of the experiment all the species where considered as pollen recipient and as pollen donor (see interaction matrix, fig 2). Species were watered once or twice per day and fertilized weekly (NPK 23: 3.95: 14).

Table 1

Family	Genus	Species
Brassicaceae	Brassica	Brassica rapa
Brassicaceae	Brassica	Brassica oleracea
Brassicaceae	Eruca	Eruca versicaria
Brassicaceae	Sinapis	Sinapis alba
Convolvulaceae	Ipomoea	Ipomoea aquatica
Convolvulaceae	Ipomoea	Ipomoea purpurea
Solanaceae	Capsicum	Capsicum annuum
Solanaceae	Petunia	Petunia integrifolia
Solanaceae	Solanum	Solanum lycopersicum
Solanaceae	Solanum	Solanum melongena

105 **Hand-pollination**

106 Foreign pollen effect was studied through two different treatments, one with 50% conspecific pollen and
107 50% heterospecific pollen and a second one with 100% foreign pollen (N=10) this second I don't get,
108 maybe explain it's utility.. Therefore, 180 different combinations were performed with N=10 per
109 combination. Seed set was the proxy of effect for all our treatments. Moreover, hand cross pollination
110 (between individuals of the same species), hand self pollination, apomixis (bagged emasculated flowers)
111 and natural selfing were tested for each species (N=10). For the treatments with foreign pollen and
112 hand cross pollination, flowers were emasculated the day prior anthesis and hand pollinated next day
113 with a toothpick. Hand-pollination was conducted with 3-4 gentle touches on the stigma surface. The
114 mixes of pollen were realized on an eppendorf based on the pollen counts made with Neubauer chamber
115 (each anther was counted 4 times for 20 different anthers per species)-IB explain better and give a bit
116 more of detail. In order to confirm that the treatments applied were 50-50 percent pollen, for each focal
117 species the total stigmatic load of pollen was counted from one donor of each family (N=3).

118 **Traits and evolutive distance**

119 The traits measured for each species were pollen per anther, number of ovules, stigma width and length
120 and stigmatic area, style width and length, ovary width and length. Moreover stigma type explain was
121 tested. Pollen was counted for 20 anthers of each species with 4 replicates per sample with an
122 hemocytometer. Previously, anthers were squashed on a known solution with the pipette tip and
123 homogeneize with a vortex for 30 seconds. Ovule number was counted with the help of an
124 stereomicroscope and a small grid over a petri dish from 15 randomly selected flowers. The different
125 morphometrical traits were measured with a digital stereomicroscope. Levels of self incompatibility were
126 estimated by dividing the fruit set of hand self pollination by hand cross pollination Lloyd and Schoen
127 (1992).

128 **Analysis**

129 We used the statistical language R (R Core Team 2018) for all our analyses. Differences of seed set
130 between treatments and hand cross pollination for each species was tested through mixed linear models.
131 For the following analysis we scaled the values of seed production for all the species with mean 0 and sd

132 of 1. To test the effect of heterospecific pollen, we subtracted to the seed set of hand cross pollination
133 the seed set of heterospecific pollen treatments. In order to see correlations between heterospecific
134 pollen effect and traits we performed Mantel test between the matrix of heterospecific pollen effect and
135 the distance matrix of each trait (euclidean distances). Moreover, Mantel test was also conducted
136 between heterospecific pollen effect and the square root of the matrix of phylogenetic distance due to
137 improvement in the statistical power (Letten & Cornwell 2014). all is here, but I would break it by
138 questions and give a bit more detail, to avoid overwhelm the reader We explored also the relations
139 between traits and heterospecific pollen effect through generalized mixed models where the response
140 variable was heterospecific pollen effect, the independent variable the different traits and the random
141 effects the different treatments per species [Here I think you should think if this controls for the non
142 independency of donors and recipients. I think not. Maybe look onto matrix regresions?). Moreover,
143 pairwise evolutive distances were calculated with MEGA7 for two kinds of markers: 1) Internal
144 transcribed spacer (ITS) and 2) ribulose-bisphosphate carboxylase (RBCL). The sequences of interest
145 were downloaded from NCBI GenBank and the phylogenetic tree constructed by maximum likelihood
146 with MEGA7. Make a section on how you constructed phylogeny.

147 I would explain three test. 0) treatment effects with GLM's, 1) Mantels: relative effects, 2) GLM's or
148 matrix models: Absolute effects and explain them in three independent paragraphs including a
149 rationale of why

150 [Phylogenetic signal of traits?](#)

RESULTS

Results of hand cross pollination, self hand pollination, natural selfing and apomixis are presented in **Table 2**. Heterospecific pollen reduced seed set significantly with the 50-50% heterospecific pollen treatments for 65% of the pairwise interactions $p < 0.05$. Across families we found a very similar effect but when species were looked at species level they responded differently even within the same family rephrase and maybe test statistically?, for instance for two species of the Brassicaceae family *Brassica oleracea* and *Eruca versicaria* we found very contrasting effects of foreign pollen where for the first one, all donors reduce seed set significantly and for the second, just two species did out of nine. The 100% foreign pollen treatments barely produced seeds or fruits and just for *Sinapis alba* we did not find significant differences between the hand cross pollination and one treatment with pollen from a confamilial- IB Unclear. Solanaceae species with berry fruit type developed small fruits or even normal fruits in some cases under which treatment. *S. lycopersicum* seems to produce small fruits (35% of the treatments) independently of pollen and pollen donor due to also apomictic treatments did, never normal size. *C. annuum* produced some fruits (9%) of both small and normal size and finally *S. melongena* produced seedless normal fruits with just confamilial pollen (3%), for both species seems that fruit formation was induced by pollen on the stigma because of lack of fruit production with treatments that tested for apomixis. clarify this descriptive statistics part- Also a figure with a summary of the treatments effect would be cool, or at least in the appendices

169 **Table 2.** Percentage of seeds produced per ovule for the ten species used in the experiment. The
170 treatments presented are hand cross pollination, hand self pollination, natural selfing and apomixis
171 (emasculated flowers). turn into a figure somehow?

Species	Cross	Self	Natural_selfing	Apomixis
Brassica oleracea	32.06897	0.0000000	0.00000	0
Brassica rapa	44.97041	0.0000000	0.00000	0
Eruca versicaria	23.75000	0.4166667	0.00000	0
Sinapis alba	43.33333	48.3333333	5.00000	15
Ipomoea aquatica	40.00000	30.0000000	20.00000	0
Ipomoea purpurea	31.66667	86.6666667	31.66667	0
Capsicum annuum	100.00000	66.2240664	23.48548	0
Petunia integrifolia	100.00000	24.7727273	0.00000	0
Solanum lycopersicum	90.38043	43.4782609	70.00000	0
Solanum melongena	60.47525	87.9702970	21.56436	0




Figure 1: Barplot with the different treatments that provide information of the reproductive biology of the ten species. The y axis is the proportion of ovules converted to seed in percentage. The different treatments (N=10) which are presented in the legend are, hand cross pollination, hand self pollination, natural selfing and apomixis. More information about these treatments can be found in Methods and Appendices.

172 Mantel test indicates that a possible?? It exists! correlation exist between heterospecific pollen effect
 173 and the evolutive relative distances, for ITS and RBCL markers we had r coefficients of 0.29 and 0.25
 174 respectively $p < 0.05$ think on a figure - maybe using NMDS. Moreover, Mantel test indicates that also a
 175 possible?? correlation between stigma width and stigma type exist (stats??). Trait correlations were
 176 also explored with GLMM

177 I have done it at the moment just for Compatibility system Also I have to fix from mixed linear model
 178 to GLMM, just realize that




Figure 2: The effect of heterospecific pollen (scaled see set) is represented in function of the compatibility system (self/cross*100) for the the different species. Each coulored dot represents the interaction of a focal species with a different pollen donor.

179 [Compatibility index don´t multiply per 100 from Lloyd](#)

DISCUSSION

Discussion

What are the implications of the findings?

Ideas about pollen size in heterospecific pollen effect. (still have to develop it more...)

Let's classify pollen size in three groups in order to understand the interaction between pollen donor and recipient: 1) Donor pollen size < Recipient pollen size 2) Donor pollen size = Recipient pollen size 3) Donor pollen size > Recipient pollen size

Now I try to develop each part

1) Donor pollen size < Recipient pollen size

Effect:

- Donor's pollen could clog the stigma
- Chemical inhibition

Traits associated with bigger pollen of the recipient:

- Recipient's pollen have faster pollen tube growth (example with my data)
- Reduction in number of ovules (Also with my species)
- Big differences in pollen size can be traduced in low relatednes therefore less likely of pollen germination on a far related stigma.

2) Donor pollen size = Recipient pollen size

- Very relatedness dependant this point
- Similar probabilities of taken space on the stigma

3) Donor pollen size > Recipient pollen size

201 Effect:

202 -In small stigmas big pollen grains can occupy great part of the stigmatic area.

203 -small pollen grains can get embedded

204 IB: Think also on using tree analysis to test if hp effect depends on complex trait combinations. Tree
205 analysis are great when two different strategies lead to the same outcome. This would never been pick
206 up by GLMs. The r package is party{} . You can see an example applied to birds is Sol et al 2010
207 Science. Ask me if you want more details or code examples.

208 CONCLUSIONS

209 ACKNOWLEDGEMENTS

210 REFERENCES

- 211 Aizen, M. A., and L. D. Harder. 2007. Expanding the limits of the pollen-limitation concept: Effects of
212 pollen quantity and quality. *Ecology* 88:271–281.
- 213 Arceo-Gómez, G., and T.-L. Ashman. 2016. Invasion status and phylogenetic relatedness predict cost
214 of heterospecific pollen receipt: Implications for native biodiversity decline. *Journal of Ecology*
215 104:1003–1008.
- 216 Ashman, T.-L., and G. Arceo-Gómez. 2013. Toward a predictive understanding of the fitness costs of
217 heterospecific pollen receipt and its importance in co-flowering communities. *American Journal of*
218 *Botany* 100:1061–1070.
- 219 Bartomeus, I., J. Bosch, and M. Vilà. 2008. High invasive pollen transfer, yet low deposition on native
220 stigmas in a carpobrotus-invaded community. *Annals of Botany* 102:417–424.
- 221 Carvalheiro, L. G., J. C. Biesmeijer, G. Benadi, J. Fründ, M. Stang, I. Bartomeus, C. N.

222 Kaiser-Bunbury, M. Baude, S. I. Gomes, V. Merckx, and others. 2014. The potential for indirect effects
 223 between co-flowering plants via shared pollinators depends on resource abundance, accessibility and
 224 relatedness. *Ecology letters* 17:1389–1399.

225 Engel, E. C., and R. E. Irwin. 2003. Linking pollinator visitation rate and pollen receipt. *American*
 226 *Journal of Botany* 90:1612–1618.

227 Fang, Q., and S.-Q. Huang. 2013. A directed network analysis of heterospecific pollen transfer in a
 228 biodiverse community. *Ecology* 94:1176–1185.

229 Inouye, D. W. 1980. The terminology of floral larceny. *Ecology* 61:1251–1253.

230 Lloyd, D. G., and D. J. Schoen. 1992. Self-and cross-fertilization in plants. i. functional dimensions.
 231 *International Journal of Plant Sciences* 153:358–369.

232 Magrach, A., J. P. González-Varo, M. Boiffier, M. Vilà, and I. Bartomeus. 2017. Honeybee spillover
 233 reshuffles pollinator diets and affects plant reproductive success. *Nature ecology & evolution* 1:1299.

234 Montgomery, B. R., and B. J. Rathcke. 2012. Effects of floral restrictiveness and stigma size on
 235 heterospecific pollen receipt in a prairie community. *Oecologia* 168:449–458.

236 Morales, C. L., and A. Traveset. 2008. Interspecific pollen transfer: Magnitude, prevalence and
 237 consequences for plant fitness. *Critical Reviews in Plant Sciences* 27:221–238.

238 Pauw, A. 2013. Can pollination niches facilitate plant coexistence? *Trends in ecology & evolution*
 239 28:30–37.

240 R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for
 241 Statistical Computing, Vienna, Austria.

242 Thomson, J. D., B. J. Andrews, and R. Plowright. 1982. The effect of a foreign pollen on ovule
 243 development in *diervilla lonicera* (caprifoliaceae). *New Phytologist* 90:777–783.

244 Tong, Z.-Y., and S.-Q. Huang. 2016. Pre-and post-pollination interaction between six co-flowering

- 245 pedicularis species via heterospecific pollen transfer. *New Phytologist* 211:1452–1461.
- 246 Waser, N. M., L. Chittka, M. V. Price, N. M. Williams, and J. Ollerton. 1996. Generalization in
247 pollination systems, and why it matters. *Ecology* 77:1043–1060.
- 248 Whitehead, M. R., R. Lanfear, R. J. Mitchell, and J. D. Karron. 2018. Plant mating systems often vary
249 widely among populations. *Frontiers in Ecology and Evolution* 6:38.

251 List of Figures

252	1	Barplot with the different treatments that provide information of the reproductive biology	
253		of the ten species. The y axis is the proportion of ovules converted to seed in percentage.	
254		The different treatments (N=10) which are presented in the legend are, hand cross	
255		pollination, hand self pollination, natural selfing and apomixis. More information about	
256		these treatments can be found in Methods and Appendices.	9
257	2	The effect of heterospecific pollen (scaled see set) is represented in function of the	
258		compatibility system (self/cross*100) for the the different species. Each coulored dot	
259		represents the interaction of a focal species with a different pollen donor.	10