

# working title Compatibility system and stigma size are the main predictors of heterospecific pollen effect

Jose B. Lanuza, Ignasi Bartomeus, Tia-Lynn Ashman, Romina Rader \* <sup>1,2,3</sup>

<sup>1</sup>US Environmental Protection Agency, Office of Research and Development, National Health and Environmental Effects Research Laboratory, Atlantic Ecology Division, 27 Tarzwell Drive Narragansett, RI, 02882, USA

<sup>2</sup>Big Name University, Department of R, City, BN, 01020, USA

<sup>3</sup>Estacion Biologica de Donana (EBD-CSIC), E-41092 Sevilla, Spain

\* corresponding author: [barragansljose@gmail.com](mailto:barragansljose@gmail.com)

---

Pollinator sharing can have negative consequences for plant fitness with the arrival of foreign pollen. However, the costs of heterospecific pollen are not yet well understood. We conducted a glasshouse experiment to understand how phylogenetic relatedness and plant traits mediate the impacts of heterospecific pollen transfer. We conducted 4XXXX crosses by experimentally transferring pollen (50% and 100% ratio) with reciprocal crosses between 10 species belonging to three different families: Brassicaceae, Solanaceae and Convolvulaceae. Seed set was used as proxy of plant fitness. We found that for 65% of the treatments with 50% mix reduced seed set. Moreover, the reduction in seed set was dependent on the degree of relatedness and reproductive traits of the pollen recipient and not the pollen donor. Our results show that certain traits, particularly compatibility system, are critical in understanding the costs of heterospecific pollen.

---

**Keywords:** heterospecific pollen, plant reproduction, fitness, interspecific competition, phylogenetic distance.

## INTRODUCTION

In most ecosystems, plant species normally coexist and share their floral visitors with other species Waser et al. (1996). From the plants' perspective, pollinator sharing can be positive for some plants Carvalheiro et al. (2014) or negative for others Pauw (2013), depending on the facilitation gradient. An increasing number of visits often correlates with higher chances of fertilization Engel and Irwin (2003). However this is not always the case, among these possible flower visitors there are also nectar robbers and pollen thieves Inouye (1980); Magrach et al. (2017). Receiving both sufficient quantity and quality deposited on the stigma is thus highly relevant to the pollination success of the plant Aizen and Harder (2007).

34 By visiting many plant species, many pollinators are responsible for conspecific pollen loss and the  
35 transport of foreign pollen, both of which can have important detrimental effects on species fitness  
36 Morales and Traveset (2008); Ashman and Arceo-Gómez (2013); Arceo-Gómez and Ashman (2016).  
37 Foreign pollen arrival can play an important role in plant species fitness but outcomes are variable and  
38 appear to be context dependent as there is not always a decrease in fitness Morales and Traveset (2008).  
39 Some of this variation is likely due to the enormous variability of foreign pollen transferred across  
40 systems ranging from 0 to 75 percent. However, most studies report ranges of heterospecific pollen  
41 between 0 and 20 percent of the total pollen load Bartomeus et al. (2008) Montgomery and Rathcke  
42 (2012); Ashman and Arceo-Gómez (2013); Fang and Huang (2013), yet even these relatively low  
43 amounts of heterospecific pollen transferred can decrease fitness greatly Thomson et al. (1982). While  
44 we now have some understanding of the impacts of heterospecific pollen quantity, we have less  
45 understanding of other factors that could be driving the variation in impacts upon fitness. Ashman and  
46 Arceo-Gómez (2013) postulated the first predictive framework that identifies a need to understand how  
47 plant traits might mediate heterospecific pollen effect, whereby mating system and pollen size were  
48 predicted to potentially mediate the impact of foreign pollen transfer on plant fitness. This concept is  
49 supported by specific case studies, such as Tong and Huang (2016) that demonstrate an asymmetrical  
50 effect in 6 species of *Pedicularis* whereby the pollen of long styled species was able to grow the full  
51 length of the style on short styled species but not vice versa. While this suggests that the impacts of  
52 heterospecific pollen may differ among pollen donor and recipient, few studies have been conducted to  
53 ascertain whether this pattern is in fact a general trend or to identify the extent to which other plant  
54 traits are critical to heterospecific pollen impacts.

55 Plant traits are crucial to understand heterospecific pollen effect but the multifactorial nature of the  
56 traits that are involve in the pollen-pistil interaction make difficult to unravel what are the main traits  
57 in driving the effect. These traits can be seen from a male perspective of both donor and recipient  
58 where pollen size, pollen aperture number and pollen allelopathy are key components to understand the  
59 outcome of foreign pollen arrival Murphy and Aarssen (1995); Ashman and Arceo-Gómez (2013). In  
60 Ashman and Arceo-Gómez (2013) small pollen is predicted to cause a greater fitness decrease, although  
61 this can be true there are also other possibilities to consider which can obscure a predictive framework  
62 like big pollen can clogg small stigmas with fewer pollen grains, bigger stigmas are less likely to be

63 clogged by small pollen grains and bigger pollen can outcompete smaller pollen grains due to faster  
64 pollen tube growth rate Williams and Rouse (1990). Moreover, to understand the different mechanical  
65 or chemical effects of pollen also the female traits of the pollen recipient have to be considered, from  
66 the literature these main traits are: stigma size, style length, number of ovules, incompatibility system  
67 and also a structural trait such as flower morphology Montgomery and Rathcke (2012); Ashman and  
68 Arceo-Gómez (2013); Tong and Huang (2016). For example, greater stigmatic area is positively  
69 correlated with greater heterospecific pollen deposition Montgomery and Rathcke (2012) and therefore  
70 possibly with an increase in negative effect. For species that are self-incompatible the barriers towards  
71 heterospecific pollen are stronger than self-compatible species Ashman and Arceo-Gómez (2013).  
72 Nonetheless, an effect of foreign pollen is a bit obscured by the variability within species, however  
73 species that are strong selfers or strong outcrossers have less variability in mating systems and  
74 predictions of effect could be more realistic (see figure 1 from Whitehead et al. (2018)). Although past  
75 research has progress in the understanding of what traits can mediate the effect as we have shown here,  
76 there are multiple traits involved and multiple possible scenarios still to be explored empirically for a  
77 full understanding of the importance of heterospecific pollen effect in nature.

78 For the understanding at what level or intensity the interference of pollen can occur is important to  
79 consider the relatedness of the interacting species. Closely related species are more likely to have  
80 similar traits Letten and Cornwell (2015). The similarity in traits between closely related species can  
81 lead to higher chances of ovule usurpation/abortion (REF) and therefore studies predict and show a  
82 greater negative effect of closely related species (REFS). Few studies however, have focused on the  
83 impacts of heterospecific pollen on fitness of distantly related species Thomson et al. (1982) Galen and  
84 Gregory (1989) Neiland and Wilcock (1999). Yet, most insects and most stigmas have been found to  
85 carry multiple species of foreign pollen with little attention to degree of relatedness (Arceo-Gómez and  
86 Ashman (2016); Fang and Huang (2013). The relative amounts of closely/far related pollen that arrives  
87 to a stigma gives a first snapshot where pollen competition can occur and improves the understanding  
88 of how important the reduction of fitness can be from heterospecific pollen where closely related species  
89 can lead to greater detrimental effects Arceo-Gómez and Ashman (2016).

90 Further, a majority of plant species are generalist and thus receive visits from multiple different  
91 pollinators. Given these are generally the ones that receive greater loads of heterospecific pollen Fang

and Huang (2013) and unrelated species are more likely to coexist with other species due to less niche overlap (Ref), understanding the role of foreign pollen from distantly related species thus deserves greater attention in understanding coexistence blah blahXXXXX refs.. Notwithstanding, the effect of heterospecific pollen of far and close related species at community level remains to be explored beyond single pairwise interactions. Moreover, heterospecific pollen studies in nature have the complexity added great environmental variability which can lead to confounding interpretations and diverse floral structures and some of them complex such as Asteraceae species, which involve tedious work to be able to study foreign pollen effect. For this reason, we investigated how floral reproductive traits and relatedness mediate the impact of heterospecific transfer by asking the following research questions : To what extent do (i) floral reproductive traits and (ii) relatedness, mediate the impacts of heterospecific pollen on seed set. We do this by creating an artificial co-flowering community with 10 species belonging to three different families with different traits.

## METHODS

The study was conducted in a glasshouse at University of New England (Armidale, Australia) from November 2017 to March 2018. Rooms were temperature controlled depending on the requirements of the species with day and night temperature differences. The species selected (**Table 1**) belonged to three different families, Solanaceae, Brassicaceae and Convolvulaceae. The criteria of species/family selection was based on close/distant related species (see phylogenetic tree for relatedness fig 1)I would explain more the bauty of our nested dessign to ensure close and far distance simultaneously, heterogeneous traits, low structural flower complexity and fast life cycle. For the purpose of the experiment all the species where considered as pollen recipient and as pollen donor (see interaction matrix, fig 2). Species were watered once or twice per day and fertilized weekly (NPK 23: 3.95: 14).

**Table 1**

Family	Genus	Species
Brassicaceae	Brassica	Brassica rapa
Brassicaceae	Brassica	Brassica oleracea

Family	Genus	Species
Brassicaceae	Eruca	Eruca versicaria
Brassicaceae	Sinapis	Sinapis alba
Convolvulaceae	Ipomoea	Ipomoea aquatica
Convolvulaceae	Ipomoea	Ipomoea purpurea
Solanaceae	Capsicum	Capsicum annuum
Solanaceae	Petunia	Petunia integrifolia
Solanaceae	Solanum	Solanum lycopersicum
Solanaceae	Solanum	Solanum melongena

## 115 Hand-pollination

116 Foreign pollen effect was studied through two different treatments, one with 50% conspecific pollen and  
117 50% heterospecific pollen and a second one with 100% foreign pollen (N=10) this second I don't get,  
118 maybe explain it's utility.. Therefore, 180 different combinations were performed with N=10 per  
119 combination. Seed set was the proxy of effect for all our treatments. Moreover, hand cross pollination  
120 (between individuals of the same species), hand self pollination, apomixis (bagged emasculated flowers)  
121 and natural selfing were tested for each species (N=10). For the treatments with foreign pollen and  
122 hand cross pollination, flowers were emasculated the day prior anthesis and hand pollinated next day  
123 with a toothpick. Hand-pollination was conducted with 3-4 gentle touches on the stigma surface. The  
124 mixes of pollen were realized on an eppendorf based on the pollen counts made with Neubauer chamber  
125 (each anther was counted 4 times for 20 different anthers per species)-IB explain better and give a bit  
126 more of detail. In order to confirm that the treatments applied were 50-50 percent pollen, for each focal  
127 species the total stigmatic load of pollen was counted from one donor of each family (N=3).

## 128 Traits and evolutive distance

129 The traits measured for each species were pollen per anther, number of ovules, stigma width and length  
130 and stigmatic area, style width and length, ovary width and length. Moreover stigma type explain was  
131 tested. Pollen was counted for 20 anthers of each species with 4 replicates per sample with an

hemocytometer. Previously, anthers were squashed on a known solution with the pippete tip and  
homogeneize with a vortex for 30 seconds. Ovule number was counted with the help of an  
stereomicroscope and a small grid over a petri dish from 15 randomly selected flowers. The different  
morphometrical traits were measured with a digital stereomicrospe. Levels of self incompatibility were  
estimated by dividing the fruit set of hand self pollination by hand cross pollination Lloyd and Schoen  
(1992).

## Analysis

We used the statistical language R (R Core Team 2018) for all our analyses. Differences of seed set  
between treatments and hand cross pollination for each species was tested through mixed linear models.  
For the following analysis we scaled the values of seed production for all the species with mean 0 and sd  
of 1. To test the effect of heterospecific pollen, we substracted to the seed set of hand cross pollination  
the seed set of heterospecific pollen treatments. In order to see correlations between heterospecific  
pollen effect and traits we performed Mantel test between the matrix of heterospecific pollen effect and  
the distance matrix of each trait (euclidean distances). Moreover, Mantel test was also conducted  
between heterospecific pollen effect and the square root of the matrix of phylogenetic distance due to  
improvement in the statistical power (Letten & Cornwell 2014). all is here, but I would break it by  
questions and give a bit more detail, to avoid overwhelm the reader We explored also the relations  
between traits and heterospecific pollen effect through generalized mixed models where the response  
variable was heterospecific pollen effect, the independent variable the different traits and the random  
effects the different treatments per species [Here I think you should think if this controls for the non  
independency of donors and recipients. I think not. Maybe look onto matrix regresions?). Moreover,  
pairwise evolutive distances were calculated with MEGA7 for two kinds of markers: 1) Internal  
transcribed spacer (ITS) and 2) ribulose-bisphosphate carboxylase (RBCL). The sequences of interest  
were downloaded from NCBI GenBank and the phylogenetic tree constructed by maximum likelihood  
with MEGA7. Make a section on how you contrsucted phylogeny.

I would explain three test. 0) treatment effects with GLM's, 1) Mantels: relative effects, 2) GLM's or  
matrix models: Absolute effects and explain them in three independent paragraphs including a  
rationale of why



## RESULTS

Results of hand cross pollination, self hand pollination, natural selfing and apomixis are presented in **Table 2**. Heterospecific pollen reduced seed set significantly with the 50-50% heterospecific pollen treatments for 65% of the pairwise interactions  $p < 0.05$ . Across families we found a very similar effect but when species were looked at species level they responded differently even within the same family rephrase and maybe test statistically?, for instance for two species of the Brassicaceae family *Brassica oleracea* and *Eruca versicaria* we found very contrasting effects of foreign pollen where for the first one, all donors reduce seed set significantly and for the second, just two species did out of nine. The 100% foreign pollen treatments barely produced seeds or fruits and just for *Sinapis alba* we did not find significant differences between the hand cross pollination and one treatment with pollen from a confamilial- IB Unclear. Solanaceae species with berry fruit type developed small fruits or even normal fruits in some cases under which treatment. *S. lycopersicum* seems to produce small fruits (35% of the treatments) independently of pollen and pollen donor due to also apomictic treatments did, never normal size. *C. annuum* produced some fruits (9%) of both small and normal size and finally *S. melongena* produced seedless normal fruits with just confamilial pollen (3%), for both species seems that fruit formation was induced by pollen on the stigma because of lack of fruit production with treatments that tested for apomixis. clarify this descriptive statistics part- Also a figure with a summary of the treatments effect would be cool, or at least in the appendices



179 **Table 2.** Percentage of seeds produced per ovule for the ten species used in the experiment. The  
180 treatments presented are hand cross pollination, hand self pollination, natural selfing and apomixis  
181 (emasculated flowers). turn into a figure somehow?

Species	Cross	Self	Natural_selfing	Apomixis
Brassica oleracea	32.06897	0.0000000	0.00000	0
Brassica rapa	44.97041	0.0000000	0.00000	0
Eruca versicaria	23.75000	0.4166667	0.00000	0
Sinapis alba	43.33333	48.3333333	5.00000	15
Ipomoea aquatica	40.00000	30.0000000	20.00000	0
Ipomoea purpurea	31.66667	86.6666667	31.66667	0
Capsicum annuum	100.00000	66.2240664	23.48548	0
Petunia integrifolia	100.00000	24.7727273	0.00000	0
Solanum lycopersicum	90.38043	43.4782609	70.00000	0
Solanum melongena	60.47525	87.9702970	21.56436	0




Figure 1: Barplot with the different treatments that provide information of the reproductive biology of the ten species. The y axis is the proportion of ovules converted to seed in percentage. The different treatments (N=10) which are presented in the legend are, hand cross pollination, hand self pollination, natural selfing and apomixis. More information about these treatments can be found in Methods and Appendices.

182 Mantel test indicates that a possible?? It exists! correlation exist between heterospecific pollen effect  
 183 and the evolutive relative distances, for ITS and RBCL markers we had r coefficients of 0.29 and 0.25  
 184 respectively  $p < 0.05$  think on a figure - maybe using NMDS. Moreover, Mantel test indicates that also a  
 185 possible?? correlation between stigma width and stigma type exist (stats??). Trait correlations were  
 186 also explored with GLMM

187 I have done it at the moment just for Compatibility system Also I have to fix from mixed linear model  
 188 to GLMM, just realize that




Figure 2: The effect of heterospecific pollen (scaled see set) is represented in function of the compatibility system (self/cross\*100) for the the different species. Each coulored dot represents the interaction of a focal species with a different pollen donor.

189 [Compatibility index don't multiply per 100 from Lloyd](#)

## 190 DISCUSSION

### 191 Discussion

192 Herbs vs trees, annual vs perennial... Many flowers vs few flowered species; structural composition on  
193 a system

194 What are the implications of the findings?

195 Ideas about pollen size in heterospecific pollen effect. (still have to develop it more...)

196 Let's classify pollen size in three groups in order to understand the interaction between pollen donor  
197 and recipient: 1) Donor pollen size < Recipient pollen size 2) Donor pollen size = Recipient pollen size  
198 3) Donor pollen size > Recipient pollen size

199 Now I try to develop each part

200 1) Donor pollen size < Recipient pollen size

201 Effect:

- 202 • Donor's pollen could clog the stigma
- 203 • Chemical inhibition

204 Traits associated with bigger pollen of the recipient:

- 205 • Recipient's pollen have faster pollen tube growth (example with my data)
- 206 • Reduction in number of ovules (Also with my species)
- 207 • Big differences in pollen size can be translated in low relatedness therefore less likely of pollen  
208 germination on a far related stigma.

209 2) Donor pollen size = Recipient pollen size

- 210 • Very relatedness dependant this point

211 • Similar probabilities of taken space on the stigma

212 3) Donor pollen size > Recipient pollen size

213 Effect:

214 -In small stigmas big pollen grains can occupy great part of the stigmatic area.

215 -small pollen grains can get embedded

216 IB: Think also on using tree analysis to test if hp effect depends on complex trait combinations. Tree  
217 analysis are great when two different strategies lead to the same outcome. This would never been pick  
218 up by GLMs. The r package is party{} . You can see an example applied to birds is Sol et al 2010  
219 Science. Ask me if you want more details or code examples.

## 220 CONCLUSIONS

## 221 ACKNOWLEDGEMENTS

## 222 REFERENCES

223 Aizen, M. A., and L. D. Harder. 2007. Expanding the limits of the pollen-limitation concept: Effects of  
224 pollen quantity and quality. *Ecology* 88:271–281.

225 Arceo-Gómez, G., and T.-L. Ashman. 2016. Invasion status and phylogenetic relatedness predict cost  
226 of heterospecific pollen receipt: Implications for native biodiversity decline. *Journal of Ecology*  
227 104:1003–1008.

228 Ashman, T.-L., and G. Arceo-Gómez. 2013. Toward a predictive understanding of the fitness costs of  
229 heterospecific pollen receipt and its importance in co-flowering communities. *American Journal of*

- 230 Botany 100:1061–1070.
- 231 Bartomeus, I., J. Bosch, and M. Vilà. 2008. High invasive pollen transfer, yet low deposition on native  
232 stigmas in a carpobrotus-invaded community. *Annals of Botany* 102:417–424.
- 233 Carvalho, L. G., J. C. Biesmeijer, G. Benadi, J. Fründ, M. Stang, I. Bartomeus, C. N.  
234 Kaiser-Bunbury, M. Baude, S. I. Gomes, V. Merckx, and others. 2014. The potential for indirect effects  
235 between co-flowering plants via shared pollinators depends on resource abundance, accessibility and  
236 relatedness. *Ecology letters* 17:1389–1399.
- 237 Engel, E. C., and R. E. Irwin. 2003. Linking pollinator visitation rate and pollen receipt. *American*  
238 *Journal of Botany* 90:1612–1618.
- 239 Fang, Q., and S.-Q. Huang. 2013. A directed network analysis of heterospecific pollen transfer in a  
240 biodiverse community. *Ecology* 94:1176–1185.
- 241 Inouye, D. W. 1980. The terminology of floral larceny. *Ecology* 61:1251–1253.
- 242 Letten, A. D., and W. K. Cornwell. 2015. Trees, branches and (square) roots: Why evolutionary  
243 relatedness is not linearly related to functional distance. *Methods in Ecology and Evolution* 6:439–444.
- 244 Lloyd, D. G., and D. J. Schoen. 1992. Self-and cross-fertilization in plants. i. functional dimensions.  
245 *International Journal of Plant Sciences* 153:358–369.
- 246 Magrach, A., J. P. González-Varo, M. Boiffier, M. Vilà, and I. Bartomeus. 2017. Honeybee spillover  
247 reshuffles pollinator diets and affects plant reproductive success. *Nature ecology & evolution* 1:1299.
- 248 Montgomery, B. R., and B. J. Rathcke. 2012. Effects of floral restrictiveness and stigma size on  
249 heterospecific pollen receipt in a prairie community. *Oecologia* 168:449–458.
- 250 Morales, C. L., and A. Traveset. 2008. Interspecific pollen transfer: Magnitude, prevalence and  
251 consequences for plant fitness. *Critical Reviews in Plant Sciences* 27:221–238.
- 252 Murphy, S. D., and L. W. Aarssen. 1995. Reduced seed set in *elytrigia repens* caused by allelopathic

253 pollen from *phleum pratense*. *Canadian Journal of Botany* 73:1417–1422.

254 Pauw, A. 2013. Can pollination niches facilitate plant coexistence? *Trends in ecology & evolution*  
 255 28:30–37.

256 R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for  
 257 Statistical Computing, Vienna, Austria.

258 Thomson, J. D., B. J. Andrews, and R. Plowright. 1982. The effect of a foreign pollen on ovule  
 259 development in *diervilla lonicera* (caprifoliaceae). *New Phytologist* 90:777–783.

260 Tong, Z.-Y., and S.-Q. Huang. 2016. Pre-and post-pollination interaction between six co-flowering  
 261 *pedicularis* species via heterospecific pollen transfer. *New Phytologist* 211:1452–1461.

262 Waser, N. M., L. Chittka, M. V. Price, N. M. Williams, and J. Ollerton. 1996. Generalization in  
 263 pollination systems, and why it matters. *Ecology* 77:1043–1060.

264 Whitehead, M. R., R. Lanfear, R. J. Mitchell, and J. D. Karron. 2018. Plant mating systems often vary  
 265 widely among populations. *Frontiers in Ecology and Evolution* 6:38.

266 Williams, E., and J. Rouse. 1990. Relationships of pollen size, pistil length and pollen tube growth  
 267 rates in *rhododendron* and their influence on hybridization. *Sexual Plant Reproduction* 3:7–17.





269 **List of Figures**

270	1	Barplot with the different treatments that provide information of the reproductive biology	
271		of the ten species. The y axis is the proportion of ovules converted to seed in percentage.	
272		The different treatments (N=10) which are presented in the legend are, hand cross	
273		pollination, hand self pollination, natural selfing and apomixis. More information about	
274		these treatments can be found in Methods and Appendices. . . . .	10
275	2	The effect of heterospecific pollen (scaled see set) is represented in function of the	
276		compatibility system (self/cross*100) for the the different species. Each coulored dot	
277		represents the interaction of a focal species with a different pollen donor. . . . .	11